ETH zürich

Using nanogranitoids and phase equilibria modeling to unravel anatexis in the crustal footwall of the Ronda peridotites (Betic Cordillera, S Spain)

Journal Article

Author(s): Bartoli, Omar; Acosta-Vigil, Antonio; Tajčmanová, Lucie; Cesare, Bernardo; Bodnar, Robert J.

Publication date: 2016-07

Permanent link: https://doi.org/10.3929/ethz-a-010735914

Rights / license: In Copyright - Non-Commercial Use Permitted

Originally published in: Lithos 256-257, https://doi.org/10.1016/j.lithos.2016.03.016

Funding acknowledgement: 335577 - Interplay between metamorphism and deformation in the Earth's lithosphere (EC)

This page was generated automatically upon download from the <u>ETH Zurich Research Collection</u>. For more information, please consult the <u>Terms of use</u>.

1 Using nanogranitoids and phase equilibria modeling to unravel

2 anatexis in the crustal footwall of the Ronda peridotites (Betic

3 Cordillera, S Spain)

4 Omar Bartoli^{1*}, Antonio Acosta-Vigil^{1,2}, Lucie Tajčmanová³, Bernardo Cesare¹,

- 5 Robert J. Bodnar⁴
- 6
- ⁷ ¹Dipartimento di Geoscienze, Università di Padova, Padova, 35131, Italy
- 8 ²Instituto Andaluz de Ciencias de la Tierra, Consejo Superior de Investigaciones Científicas and
- 9 Universidad de Granada, Avda. de Las Palmeras nº 4, Armilla 18100, Granada, Spain
- ³Department of Earth Sciences, Swiss Federal Institute of Technology, Zurich, 8092, Switzerland
- ⁴Department of Geosciences, Virginia Tech, Blacksburg, Virginia 24061, U.S.A
- 12
- 13 * corresponding author, omar.bartoli@unipd.it
- 14 Phone: +39-0498279148
- 15
- 16

17 Abstract

Anatexis in the crustal footwall of Ronda peridotites (Betic Cordillera, S Spain) is apparently related to 18 the hot emplacement of this mantle slab over metasedimentary rocks. In this study we combine the 19 analysis of melt inclusions (MI) and phase equilibria calculations on guartzo-feldspathic mylonites 20 (former migmatites) occurring at the contact with the mantle rocks, in the region of Sierra Alpujata 21 (Ojén unit). The goal is to better characterize anatexis in these rocks, and to provide new constraints on 22 23 the geodynamic evolution of the crustal footwall. Such data are important for understanding the mechanisms of crustal emplacement of the mantle rocks. The guartzo-feldspathic mylonites are 24 characterized by the mineral assemblage Qtz+Pl+Kfs+Sil+Grt+Ilm+Bt±Ap±Gr. Clusters of MI are 25 26 observed both at the core and towards the rim of peritectic garnet. In each cluster, MI range from totally glassy to nanogranitoids, consisting of Qtz+Kfs+Bt+Ms+Pl aggregates. The trapped melt is 27 leucogranitic and peraluminous with variable Na₂O/K₂O values and low H₂O contents (\approx 2-4 wt%). 28 Phase equilibria modelling in the MnO-Na₂O-CaO-K₂O-FeO-MgO-Al₂O₃-SiO₂-H₂-TiO₂-O₂-C 29 (MnNCaKFMASHTOC) system with graphite-saturated fluid constrains the P-T conditions of melting 30 at ≈ 6 kbar, ≈ 820 °C. MI data support the fluid-absent character of melting. The investigated MI 31 represent the primary anatectic melts produced during prograde anatexis of the host rocks via biotite 32 dehydration melting. Field, compositional and textural observations indicate that mylonitic migmatites 33 represent strongly deformed former diatexites. The comparison between the new data and some 34 recently published information on migmatites located further from the contact with the peridotites and 35 36 towards the bottom of the crustal footwall, raises some important issues which question the previously proposed geodynamic models for this region. Among them, i) the crustal footwall at Sierra Alpujata 37 constitutes an inverted metamorphic sequence, not only in terms of temperature but also in pressure, 38 39 and ii) the Ojén unit does not seem to represent a coherent high-P portion of a continental subduction system. 40

41 Key words: crustal anatexis, melt inclusions, nanogranitods, phase equilibria modeling, Ronda
42 peridotites

43

44 **1. Introduction**

Progress in the investigation of high-grade, partially-melted crystalline basements has been made in the 45 last decade by studying melt and fluid inclusions (e.g., Cesare et al., 2009; Frezzotti and Ferrando, 46 2015), as well as through phase equilibria modelling based on large, internally consistent 47 thermodynamic datasets (e.g., Johnson et al., 2008; White et al., 2007, 2011). Of particular importance 48 49 is the discovery of primary melt inclusions (MI) hosted in peritectic minerals of regionally 50 metamorphosed crustal rocks, which has been described in recent papers (reviewed by Cesare et al., 2015 and Bartoli et al, 2016). Being trapped by growing peritectic phases at suprasolidus conditions, 51 52 these MI represent a window into the pre-peak anatectic history of partially-melted terranes (Acosta-Vigil et al., 2010), and may provide key microstructural and compositional information on crustal 53 anatexis, such as which mineral grew in the presence of melt, if a rock has melted, when a rock has 54 55 melted, the mechanisms and nature of the melting process, and the timeframes of melt production and segregation (Cesare et al., 2015). 56

The Ronda peridotites, the largest known exposure of subcontinental lithospheric mantle on the 57 Earth surface (≈300 km²; Obata, 1980), crop out in the Internal Zones of the Alpine Betic Cordillera 58 (southern Spain; Fig. 1a), primarily in the massifs of Sierra Bermeja, Sierra Alpujata and Carratraca 59 (Fig. 1b). The peridotites occur as km-thick slabs sandwiched between mostly metasedimentary crustal 60 rocks (metapelites and metagreywackes), which are in turn characterized by their increasing 61 metamorphic grade, degree of melting and intensity of deformation towards the mantle rocks (e.g., 62 Loomis, 1972; Westerhof, 1975; Tubía et al., 1997, 2013; Acosta-Vigil et al., 2001, 2014, 2016; Platt et 63 al., 2003; Esteban et al., 2008; Barich et al., 2014). Whereas partial melting of the metasedimentary 64

rocks above the peridotite has been related to decompression (Platt et al., 2003), in the crustal footwall 65 it has been related to the hot thrusting of the mantle slab over metasedimentary rocks, resulting in a 66 dynamothermal aureole (Tubía et al., 1997, 2013). Although several studies have examined in detail the 67 structural evolution and kinematics of the crustal footwall of the Ronda peridotites (Tubía et al., 1997, 68 2013; Cuevas et al., 2006; Esteban et al., 2008), anatexis throughout this complex crustal sequence is 69 still poorly characterized, particularly close to the contact with the mantle slice. The lack of modern 70 71 petrological studies on the crustal envelope, together with recent geochronological studies showing the existence of pre-Alpine mineral associations and fabrics in the western Betic Cordillera (e.g. Acosta-72 Vigil et al., 2014; Massonne, 2014; Sánchez-Navas et al., 2014), have renewed an old debate on the 73 74 geodynamic evolution of this important sector of the orogen characterized by the presence of subcontinental mantle slabs (e.g., Platt and Vissers, 1989; Zeck et al., 1992; Michard et al., 1997). 75 Melting in the crustal footwall is apparently related to the crustal emplacement of the mantle slab 76 (Westerhof, 1975; Torres-Roldán, 1983; Tubía et al., 1997). Therefore, a detailed characterization of 77 metamorphic conditions and the nature of anatexis is a key step to the constraining the mechanism of 78 emplacement of the Ronda peridotites and to the understanding of the tectono-metamorphic evolution 79 of the whole orogen. In this contribution, we report the occurrence and characteristics of MI hosted in 80 peritectic garnet of graphite-bearing, quartzo-feldspathic mylonites cropping out in the crustal footwall 81 82 of the Ronda peridotites (Sierra Alpujata, Ojén unit), close to the contact with the mantle rocks. By combining the microstructural and compositional investigation of MI with phase equilibria modelling 83 of the host rock we obtain new and robust data on anatexis near the contact with the peridotites. These 84 85 data are then compared with recently published information on MI in migmatites from the same unit though located further from the contact with the peridotites, in order to discuss the reliability of the 86 currently proposed models on the emplacement of the mantle slab. 87

88

89 2. Geological setting

The Betic Cordillera (southern Spain) represents the westernmost part of the peri-Mediterranean Alpine 90 orogen, formed during the N-S to NW-SE convergence of the African and Iberian plates from Late 91 Cretaceous to Early Neogene times (Dewey et al., 1989). The Internal Zone of this orogenic belt shows 92 mainly metamorphic rocks of Paleozoic to Paleogene age constituting several units distributed in two 93 main tectonic complexes, the Alpujárride Complex at the bottom and Maláguide Complex at the top 94 95 (e.g. Platt et al., 2013, and references therein) (Fig. 1b). The Ronda peridotites form the lower portion of the Los Reales unit, which is the structurally highest Alpujárride unit (Navarro-Vilá and Tubía, 96 1983; Tubía, 1988). The mantle rocks are emplaced over the Guadaiza and Ojén Alpujárride units (Fig. 97 98 1b), constituting the footwall of the Ronda peridotites (Navarro-Vilá and Tubía, 1983). The basal contacts between mantle and crustal rocks are HT shear zones formed by mylonites (Esteban et al., 99 2008; Tubía et al., 1997, 2013). The Guadaiza and Ojén units show differences in lithology, peak 100 pressure and, importantly, P-T path and kinematics, with the Ojén unit recording decompression from 101 \geq 15 kbar at \geq 730 °C and dominant top-to-the-ENE sense of shear, and the Guadaiza unit showing 102 heating to 700-800 °C at 4-6 kbar and top-to-the-NNW sense of shear (Tubía et al., 1997; Esteban et 103 al., 2008; Acosta-Vigil et al., 2014; and references therein). The relationship between these units is 104 unclear, and they may have been juxtaposed through the Albornoque strike-slip subvertical fault 105 106 (Tubía, 1988). In general isograds in these crustal sequences are roughly parallel to the lithological contacts and to the regional foliation (Westerhof, 1975; Tubía et al., 2013; Acosta-Vigil et al., 2014). 107 The rocks studied in this contribution are from the Ojén unit and consist of metasedimentary 108 109 quartzo-feldspathic mylonites from the metamorphic footwall of the Sierra Alpujata massif (Figs. 1, 2), located ~50 m below the contact with the peridotite (Fig. 2). The Ojén unit shows along-strike 110 variations which have been interpreted as large-scale boundin-like structures related to the late 111 exhumation stages of the Ronda peridotites (Tubía et al., 2013). The most complete sequence has a 112

thickness of \approx 700-800 m (Fig. 2). Below we describe this unit from the top to the bottom using field 113 and structural data reported in Tubía (1988) and Tubía et al. (1997, 2013). The sequence is made of 114 mylonites, migmatites, schists and marbles (Fig. 2). At 200-300 m from the contact with the peridotites, 115 the crustal rocks are strongly deformed and have been interpreted as HT quartzo-feldspathic mylonites. 116 These rocks are characterized by ENE-trending mineral lineations, and 40° to 60° S-dipping mylonitic 117 foliation (S_{myl}) subparallel to the contact between crustal and mantle rocks; sigmoidal shear bands and 118 S-C microstructures indicate a top-to-the-ENE shearing. According to Tubía et al. (1997), mylonites 119 have been intruded by granitic bodies up to 100 m think, roughly concordant with S_{mvl}. The central part 120 of the unit comprises migmatites, with diatexites on top of banded migmatites and migmatized 121 leucocratic gneisses. Decimetric to decametric amphibolite lenses that preserve eclogitic relicts. 122 recording peak conditions of >15 kbar and >730 °C, have been described towards the bottom of the 123 migmatitic sequence (Fig. 2; Tubía and Gil-Ibarguchi, 1991). A low-temperature shear zone (ENE-124 ward shearing) separates banded migmatites and Sil-bearing, amphibolite facies schists; here 125 migmatites are intensely retrogressed. In addition, isolated shear zones have also been described within 126 127 the crustal footwall, for example between amphibolite lenses and banded migmatites. The bottom of the crustal footwall is formed by amphibolite-facies marbles. 128

Regarding deformation, the oldest event recorded in the Ojén unit, D₁, is preserved as a residual 129 schistosity (S_1) within garnet from the schists; the subsequent deformation phase D_2 produced the main 130 schistosity S₂ observed in schists and, apparently, in the banded gneisses also (Fig. 2, Tubía, 1988). S₂ 131 can be affected by open folds as a result of a younger deformation event D₃ (Tubía, 1988). The melting 132 process responsible for the formation of anatectic melt and cordierite in the migmatitic sequence is 133 considered to be related to a younger, low-P post-D₃ static event (Tubía, 1988). Finally, mylonitization 134 135 at the contact with the peridoties and the penetrative mylonitic foliation (S_{mvl}) is connected to a subsequent deformation phase D₄ (Tubía, 1988). S_{mvl} is subparallel to the main foliation S₂ in banded 136

migmatites and schists, and axes of folds developed during D_3 are parallel to the mineral lineation observed in the mylonites (Tubía et al., 1997). Following Tubía (1988), migmatization occurred after inversion of the sequence and during the emplacement of the peridotites, and migmatization and mylonitization likely represent the continuation in time of the metamorphic and structural evolution recorded by the schists at the bottom of the sequence (D_1 through D_3).

The intrusion of the amphibolite/eclogite protolith into the metasedimentary rocks was dated 142 143 (U-Pb SHRIMP zircon) at ≈184 Ma (Sanchez-Rodrigez and Gebauer, 2000). Early Miocene deformed and undeformed granitoid dikes intrude the peridotite slab (Fig. 2; Priem et al., 1979; Tubía et al., 144 1997; Acosta, 1998; Sánchez-Rodríguez, 1998; Cuevas et al., 2006; Esteban et al., 2011a). The timing 145 146 of the high-temperature metamorphism and anatexis in the crustal footwall of the Ronda peridotites. however, is controversial, as it has been ascribed to either the Alpine and/or Variscan orogenies 147 (Acosta, 1998; Sánchez-Rodríguez, 1998; Sánchez-Rodríguez and Gebauer, 2000; Esteban et al., 148 2011a; Acosta-Vigil et al., 2014). 149

The limited *P*–*T* data available indicate that rocks from different levels of the Ojén unit record different decompression paths. Mylonites at the contact with the peridotites record an evolution from ≈ 8 kbar and ≈ 800 °C to 5.5 kbar and ≈ 685 °C (Westerhof, 1975, 1977; Tubía et al., 1997). The eclogite/amphibolite layers within the migmatites record decompression from ≥ 15 kbar and ≥ 730 °C to $\approx 5-8$ kbar and $\approx 700-750$ °C (Tubía et al., 1997). In addition, Bartoli et al. (2013c) have recently determined peak conditions of 4.5–5.0 kbar and 660-700 °C for quartzo-feldspathic metatexites at the base of migmatitic sequence (see below) and at the contact with the eclogite/amphibolite layers.

157

158 **3. Analytical techniques**

Back-scattered electron (BSE) imaging and semi-quantitative energy dispersive spectroscopy (EDS)
were carried out on crystallized and glassy melt inclusions using a CAM SCAN MX2500, equipped

with LaB₆ cathode, at the Dipartimento di Geoscienze, Università di Padova (Italy) and a Jeol JSM–
6500F thermal Field Emission Scanning Electron Microscope (FESEM), at INGV (Istituto Nazionale
di Geofisica e Vulcanologia), Rome, Italy. Elemental X–ray maps were acquired at 20 and 15kV
accelerating voltage and at variable magnifications, in the range 5000–6000X, depending on the MI
size, using the FEI Quanta 600 FEG equipped with a Bruker EDX–Silicon Drifted Detector, at the
Nanoscale Characterization and Fabrication Laboratory, Institute for Critical Technology and Applied
Science, Virginia Tech, USA.

168 The compositions of biotite, feldspar and glass were obtained using a Jeol JXA 8200 Superprobe at 169 the Dipartimento di Scienze della Terra, Università di Milano (Italy). Analytical parameters for minerals were: 15 kV accelerating voltage, 5 nA current, counting time of 30 s on peak and 10 s on 170 171 background. The analytical procedure for the analyses of glass followed the recommendations of 172 Morgan and London (1996, 2005); analytical parameters were: 15 kV accelerating voltage, 2 nA current, 1 µm beam diameter and a counting time of 10 s on peak and 2 s on background. Sodium, K, 173 Al and Si were analyzed first and concurrently. Owing to Na loss during electron microprobe analysis 174 of rhyolitic glasses with effects also on K, Al and Si, concentrations were corrected by analyzing 175 leucogranitic glass standards (Morgan and London, 1996, 2005). Details concerning the application of 176 correction factors and the composition of the standard glasses are given by Ferrero et al. (2012) and 177 Bartoli et al. (2013a, b). Garnet compositions were determined using the Cameca SX50 microprobe of 178 the C.N.R.-I.G.G. (Consiglio Nazionale delle Ricerche-Istituto di Geoscienze e Georisorse) at the 179 Dipartimento di Geoscienze, Università di Padova, Italy. Measurements were performed using a 20 kV 180 accelerating voltage, 20 nA beam current, and counting times of 10 s on peak and 5 s on background. 181 Natural and synthetic silicates and oxides were used as standards. 182

183

4. Sample description

185 *4.1 Field relationships*

Here we integrate the field description reported in Section 2 with our observations; this integration is 186 187 shown in Fig. 2. In detail, the mylonitic package beneath the Ronda peridotite at Sierra Alpujata is composed of ≈ 30 m of rutile-bearing metapelitic granulites at the contact with the mantle rocks, and 188 \approx 200 m of apparently rutile-free, and mostly metasedimentary quartzo-feldspathic mylonites, that grade 189 progressively downwards into metasedimentary diatexites, at the top of the migmatitic sequence, and 190 metatexites, at the bottom (Fig. 2). The metatexites would be equivalent to the banded migmatites and 191 migmatized gneisses of Tubía (1988) and Tubía et al. (1997) (see above). Based on mineral proportions 192 and whole-rock geochemistry, both morphological types of migmatites have been grouped into pelitic 193 and quartzo-feldspathic (Acosta, 1998; Acosta et al., 2001). Diatexites can be nebulitic, schlieric or 194 schollen rocks; nebulitic varieties contain fragments of paleosome rotated into different orientations 195 (see also Tubía, 1988), whereas a magmatic foliation (S_{mag}) defined by oriented elongated fragments of 196 paleosome, cm-thick leucosomes and K-feldspar megacrysts develops in the schlieric migmatites (Fig. 197 198 2; see also Acosta, 1998). Metatexites, located above the schists, mostly appear as stromatic migmatites with garnet-bearing undeformed leucosomes that define S_2 (Acosta, 1998; Bartoli et al., 2013c). 199 Locally, metatexites are folded and the main foliation is obliterated by axial-planar cordierite-bearing 200 201 and undeformed patches, representing cordierite-neosomes (Acosta, 1998; see below). S_{mvl}, S_{mag}, and S_2 are all subparallel (Fig. 2). 202

In this contribution we have investigated in detail the microstructures, MI and petrology of quartzo-feldspathic mylonites located ~50 m below the contact with the peridotite (Fig. 2). They occur as deformed, banded rocks composed of alternating fine- to medium-grained leucocratic bands (30-50 vol.%) and fine-grained mesocratic bands (50-70 vol.%) (Fig. 3a, b). The main foliation S_{myl} strikes N70°E and dips 60° to S, in agreement with what observed in other structural stations (Fig. 1c).

209 *4.2 Petrography*

The fine-grained matrix of the mesocratic bands (grain size of $\approx 20-200 \ \mu\text{m}$) is composed mostly by 210 211 Qtz+Kfs+Pl+Sil (mineral abbreviations after Kretz, 1983) and minor biotite and ilmenite, and includes frequent porphyroclasts of garnet (0.5-3 mm in diameter) and K-feldspar (up to 2 cm in size) (Fig. 3c-212 f). Some K-feldspar porphyroclasts have an augen-like appearance and may show simple twinning 213 214 (Fig. 3d). Rare irregularly shaped domains of quartz containing euhedral plagioclase and biotite may be present in the strain shadows of these crystals (Fig. 3e). Locally sillimanite may occur as $\approx 150-200 \,\mu m$ 215 prismatic crystals (Fig. 3f). Accessory phases are graphite, apatite, zircon and monazite. Garnet modal 216 proportion is \approx 5-10 vol.%, whereas that of biotite never exceeds 2-5 vol.%. The main foliation (S_{mvl}) is 217 defined by the alignment of sillimanite folia, ribbons of quartz, minor elongated crystals of biotite and 218 ilmenite, and the alternation of sillimanite-rich and leucocratic layers (Fig. 3b, c). Graphite (<1 vol.%) 219 is randomly distributed in the matrix, whereas apatite and ilmenite are generally associated with 220 sillimanite and biotie. Garnet and K-feldspar porhyroclasts may contain mineral inclusions of biotite, 221 222 quartz, plagioclase, sillimanite and graphite, often not oriented. Garnet frequently contains MI (see 223 below). In some garnet crystals, rare mineral inclusions together with abundant MI define an internal foliation having a sigmoidal to spiral-like shape (see below). Biotite often grew in the strain shadows 224 225 associated with garnet, partially to totally replacing it (Fig. 3f). In the fine-grained matrix, quartz shows cuspate-lobate boundaries and subgrains (Fig. 3g), sometimes with approximately square subgrains 226 resembling chessboard patterns (Fig. 3h), formed in response to grain boundary migration 227 228 recrystallization. Leucocratic bands may show a high lateral continuity with thicknesses of 1-50 cm, and are mainly 229 composed of Pl+Kfs+Qtz ranging in size from \approx 300 µm up to 2 mm. Sillimanite, garnet and biotite are 230

accessory phases. Crystals in these bands are often rounded and/or elongated, and mantled by a fine-

grained mesocratic matrix (Fig. 3 i). Locally, leucocratic bands may contain igneous microstructures,
such as feldspars displaying euhedral shapes with straight boundaries (Fig. 3j).

The apparently progressive change downwards observed in the field from mylonites at the contact 234 with the peridotites to the different morphological types of migmatites towards the lower part of the 235 sequence (Fig. 2), is also reflected in mineral proportions, mineral chemistry and microstructures of the 236 metasedimentary rocks. For instance, and considering similar bulk rock compositions (Table 1), the 237 238 above described microstructures in quartzo-feldspathic mylonites contrast with those in quartzofeldspathic stromatic metatexites towards the bottom of the sequence, which are characterized by i) 239 larger grain size of the matrix minerals (up to ≈ 1.5 mm), ii) higher proportions of biotite ($\approx 8-10$ 240 vol.%), iii) lower proportions of garnet (\approx 2-5 vol.%) occurring as small (50–200 µm in diameter) 241 crystals, iv) lower crystallinity of sillimanite which is always present as fibrolite, v) a foliation defined 242 by abundant oriented biotite generally clustered with sillimanite and vi) the presence of thin and 243 discontinuous garnet-bearing leucosomes parallel to the main foliation and typically with subhedral 244 igneous microstructures (Fig. 4a, b; Bartoli et al., 2013c). In addition the metatexites locally show 245 cordierite-neosomes which obliterate the main foliation and are discordant with respect to the garnet-246 bearing leucosomes (Fig. 4a, Acosta, 1998). Here apparently peritectic cordierite coexists with euhedral 247 crystals of feldspars and cuspate domains of quartz (Fig. 4c) suggesting that these portions are former 248 249 patches of anatectic melt (Sawyer, 2008).

250

251 *4.3 Mineral chemistry*

Biotite composition in quartzo-feldspathic mylonites is variable, particularly regarding Ti and X_{Mg} that range from 0.21 to 0.69 apfu and from 0.43 to 0.53, respectively (Table 1). This variability displays some systematic patterns as a function of the microstructural position. Biotite replacing garnet has higher X_{Mg} (0.47- 0.53) and lower Ti content (0.21-0.55 apfu) than biotite in the mesocratic matrix $(X_{Mg}=0.43-0.50; Ti=0.51-0.69 apfu)$. Inclusions of biotite in garnet and K-feldspar porphyroclasts are very similar in composition to biotite in the matrix. In addition, biotite replacing garnet shows higher F contents (up to 1.5 wt%) than biotite in the rock matrix (up to 1.0 wt%). Cl is low in all investigated crystals (0.2-0.6 wt%).

Garnet is an almandine-rich solid solution (Table 1) and no chemical variations were observed 260 between either MI-free and MI-bearing garnet, or garnet in mesocratic and leucocratic portions of the 261 262 rock. Garnet cores in the mesocratic matrix and leucocratic bands have a similar composition (Alm₇₂- $_{75}$ Prp₂₀₋₂₃Sps₀₂₋₀₃Grs₀₂₋₀₃; X_{Mg} =0.21-0.24). Most of the garnets are unzoned, with Fe, Mg, Mn and Ca 263 being fairly homogeneous throughout the crystal. Garnet rims have a composition of Alm72-76Prp20-264 24Sps02-03Grs02 (XMg=0.21-0.25). Only garnets in contact with biotite are zoned: almandine and 265 spessartine components increase from core to rim, whereas pyrope component decreases (Table 1). 266 Thus, garnet rims in contact with biotite display a composition of Alm76-79Prp15-20Sps03-05Grs02, with 267 $X_{Mg}=0.17-0.20$. Calcium content is always low and constant (CaO ≈ 0.7 wt%). 268 Plagioclase within the mesocratic matrix has a composition of $Ab_{66-73}An_{25-32}Or_{01-02}$, whereas crystals 269 slightly more albitic and richer in orthoclase component (Ab₇₀₋₇₆An₂₁₋₂₆Or₀₃₋₀₅) are present in the 270 leucocratic bands (Table 1). Plagioclase grains included in garnet porphyroclasts display compositions 271

272 $(Ab_{67-72}An_{27-31}Or_{02})$ that overlap with those of plagioclase in the matrix. No compositional differences

have been observed between K-feldspar porphyroclasts (Or₇₄₋₈₃Ab₁₇₋₂₆An₀₀₋₀₁) and K-feldspar crystals

274 in leucocratic bands ($Or_{75-82}Ab_{18-25}An_{00-01}$).

275 Compared to the investigated quartzo-feldspathic mylonites, biotite in the matrix of quartzo-

feldspathic metatexites shows clearly lower X_{Mg} (0.33-0.35), F (0.39), Cl (0.07) and, to lower extent, Ti

content (0.42-0.49 apfu), whereas garnet is enriched in Alm and Sps components having a composition

Alm₇₇₋₇₈Prp₁₁₋₁₃Sps₀₇₋₀₉Grs₀₃₋₀₄ (Bartoli, 2012; Bartoli et al., 2013c). These data are in accordance with

a lower T of equilibration of metatexites with respect to the mylonites, in accordance with phase

equilibria modeling (see below, and Bartoli et al., 2013c) and field and petrology data (see above;
Acosta, 1998).

282

283 5. Microstructural and chemical characterization of melt inclusions

Here we describe MI found in the quartzo-feldspathic mylonite ALP13 and, at the end of this section, their compositional features are briefly compared with those described in Bartoli et al. (2013c) for MI in quartzo-feldspathic metatexites.

MI-bearing garnet crystals show angular to rounded shapes both in the mesocratic matrix and in 287 leucocratic bands, and may contain clusters of several MI (Fig. 5). Melt inclusions clusters with a 288 289 subspherical geometry may occur both at the core and towards the rim of garnet (Figs. 3f, 5a). More rarely, clusters have a sigmoidal to spiral-like shape (Fig. 5b,c). Melt inclusions generally do not occur 290 close to the large mineral inclusions, similarly to what has been observed in garnets from anatectic 291 enclaves by Acosta-Vigil et al. (2007). Melt inclusions have isometric (regular) shapes and range from 292 approximately 2 to 10 µm in diameter. In transmitted light, most of the MI appear totally or partially 293 dark-brownish (Fig. 6a), and contain a polycrystalline aggregate of birefringent crystals under cross-294 polarized light. Other MI are transparent in plane-polarized light (Fig. 6b) and contain a homogeneous 295 isotropic phase, i.e. glass, often with an empty (no H_2O and CO_2 have been detected by Raman 296 297 spectroscopy) shrinkage bubble (Fig. 6b). In some garnets, dark-brownish MI mantle fibrolite needles (Fig. 6c). 298

When MI are investigated under the SEM, they appear typically facetted, and often with a welldeveloped negative crystal shape (Figs. 7, 8). In each cluster, MI show a variable degree of
crystallization ranging from totally crystallized MI (nanogranitoids; Cesare et al., 2015) to glassy (i.e.,
crystal-free) MI (Figs. 6a, 7a). No systematic difference in diameter between crystallized and glassy MI
is observed. Locally, decrepitation tails project from nanogranitoids into the surrounding host (Fig. 7b).

Crystallized inclusions contain aggregates of quartz, K-feldspar, biotite, muscovite, plagioclase (often
 modal Kfs>Pl) with equigranular, hypidiomorphic to allotriomorphic texture (Figs. 7b, 8). Crystal size
 ranges from hundreds of nm to a few µm.

307 Previous microstructural and experimental studies on other migmatitic rocks (see Ferrero et al.,

2012; Bartoli et al., 2013b) have found that, despite the different degree of crystallization, MI within

the same cluster have similar melt compositions. Therefore, we assume that nanogranitoids in

310 mylonites are likely to have similar compositions to the coexisting glassy inclusions. Electron

microprobe analyses have been performed on 39 totally glassy MI from three different thin sections of

sample ALP13. The composition of the trapped melt is leucogranitic (SiO₂ \approx 76 wt%,

FeO+MgO+MnO+TiO₂ < 2 wt%) and peraluminous [ASI =1.05-1.38; ASI = mol.

Al₂O₃/(CaO+Na₂O+K₂O)] (Table 2). The average maficity value (atomic Fe+Mg, Villaros et al., 2009)

is generally low, <0.03. The analyzed MI are highly variable in Na₂O and K₂O contents and, based on

the K# [=mol. $K_2O/(Na_2O+K_2O)$], can be grouped into two types: type I MI are characterized by K#

317 ≥ 0.6 (Na₂O/K₂O <0.5), whereas type II MI show K# ≤ 0.5 (Na₂O/K₂O >0.6) (Table 1). From all the

analyzed MI, only 8 (found in 2 garnets) correspond to the type II, whereas the rest (31) are type I MI

319 (Table 2). No compositional differences are observed between MI in mesocratic and leucocratic

320 portions. The average H_2O content estimated by difference (i.e. 100-EMP totals) is slightly but

321 consistently lower in type I MI (2.4 wt%) with respect to type II MI (3.6 wt%). In terms of CIPW

normative values, all the analyzed melts are corundum-normative. In the normative Qtz-Ab-Or

diagram, all MI plot in the Qtz field above the 5 kbar H₂O Qtz-Ab and Qtz-Or haplogranite cotectic

324 curves, and at some distance from the eutectic melt compositions (Fig. 9). Data from analyzed glassy

325 MI define two different clusters according to their K# (Fig. 9).

326 Interestingly, compositions of type II MI overlap those of analyzed both glassy and rehomogenized

327 MI from the quartzo-feldspathic metatexites at the base of the migmatitic sequence (Fig. 9; see also

Bartoli et al., 2013b). Glassy MI from metatexites, however, consistently show lower FeO (\approx 1.20 wt%) and higher concentrations of H₂O (5.4-9.1 wt%) and higher Na₂/K₂O when compared to MI in mylonites (Bartoli et al., 2014, 2015), in agreement with the lower temperatures of melting in their host rocks (\approx 700 °C; Bartoli et al., 2013c).

332

333 6. Phase equilibria modelling

Modeling the *P*–*T* conditions of melting and entrapment of the MI requires considering the presence 334 of graphite, implying the involvement of a graphite-saturated COH fluid. In such a system, the H₂O 335 activity is lowered below unity due to the presence of diluting carbonic species such as CH₄ and/or CO₂ 336 (Connolly and Cesare, 1993). Consequently, dehydration and partial melting reactions adjust their 337 position in the *P*–*T* space to accommodate this change of a_{H2O} (Thompson and Algor, 1977; Spear 338 1993). Under the assumption that the fluid in the rock is essentially produced by H₂O release from 339 phyllosilicates, the amount of H₂ and O₂ components in the fluid is constrained at a ratio 2:1. At this 340 initial condition (i.e. $X_0 = 1/3$ of Connolly, 1995), the fluid composition contains the maximum activity 341 of H₂O for a graphite-saturated COH fluid. 342

A phase diagram has been constructed for the mylonitic sample ALP13 collected from the 343 344 mesocratic portion of the outcrop of Fig. 3a and containing only a thin (\approx 1 cm) leucocratic band. Because the sample ALP13 displays a non-residual bulk rock composition that, in addition, is very 345 346 similar to those of quartzo-feldspathic undeformed diatexites at the top of the migmatitic sequence and 347 stromatic metatexites at the bottom of the migmatitic sequence (all corresponding to Ca-poor 348 peraluminous greywackes; Table 1), we infer that the bulk composition used in this modeling is not 349 affected by any significant gain or loss of melt. The model chemical system MnNCaKFMASHTOC 350 was used with the bulk rock composition obtained from XRF analysis (see the upper left inset of Fig. 10a). The amount of H₂O component involved in the calculation was assumed as the loss of ignition of 351

XRF analysis and thus represents the H₂O content available for equilibration of the observed mineral 352 assemblage. The amount of C was estimated from the modal proportion of graphite in the rock. 353 Although the Mn content of this rock is low (< 0.1 wt%), MnO was included in the modeling due to its 354 influence on the stability of garnet (Spear 1993; Tinkham et al., 2001). All calculations were done by 355 the Gibbs energy minimization using the Perple X software (Connolly, 2009) with the thermodynamic 356 database of Holland and Powell (1998, as revised in 2003). We used the solution model of melt from 357 358 White et al. (2007), of garnet from Holland and Powell (1998), of biotite from Tajčmanová et al. (2009), of white mica from Coggon and Holland (2002), of plagioclase from Newton et al. (1980) and 359 of K-feldspar from Thompson and Hovis (1979). An ideal model was used to account for the solution 360 of Mn in cordierite and ilmenite. 361

Based on the petrographic study, we deduce that the stable mineral assemblage associated with melt 362 (i.e., MI) in the studied mylonites corresponds to the quadrivariant field Grt-Bt-Sil-Pl-Kfs-Qtz-Gr-363 Ilm–Liq–COH (where COH means graphite-saturated COH fluid) in the upper right corner of the phase 364 diagram section (Fig. 10a). The relevant compositional isopleths for MI-bearing garnet cores (X_{Mg} = 365 0.21-0.24; $X_{\text{Grs}} = 0.02-0.03$; $X_{\text{Sps}} = 0.02-0.03$) cross consistently in this field at ≈ 820 °C and ≈ 6 kbar 366 (Fig. 10b). Similar temperature conditions are found considering the X_{Mg} of biotite (0.44-0.49) within 367 the mesocratic matrix (Fig. 10b). At the pressure of interest, i.e. 6 kbar, muscovite and biotite are 368 totally consumed at ≈700 and ≈840-850 °C, respectively (Fig. 10a). After crossing the Liq-in curve 369 $(\approx 700 \text{ °C})$ the modal proportion of melt and garnet increases, whereas that of biotite decreases (Fig. 370 10c). Rutile appears at P \geq 7-8 kbar. The X_{Mg} = 0.17-0.20 and X_{Sps} = 0.03 isopleths for garnet rim in 371 contact with biotite (\pm Pl, \pm Qtz) would overlap at \approx 800 °C and \approx 5 kbar in the same quadrivariant field 372 (Fig. 10b). However, the X_{Mg} of low-Ti biotite (0.47-0.53) does not match these conditions (see Section 373 7.2 for explanation). The proportion of grahite is nearly constant in the whole P-T space (≈ 0.13 vol. 374 %). The stability field of COH fluid is connected to another phase containing the O₂ component, which 375

is biotite (see Section 7.3). It is important to point out that the amount of peritectic garnet inferred at P-376 377 T conditions of interest from thermodynamic modeling (2-3 vol.%; Fig. 10c) is not consistent with petrographic observations which suggest instead modal amounts of \approx 5-10 vol.%. This is likely due to 378 the detrimental effect of Ti on the modal proportions of some phases. Bartoli et al. (2013c) have shown 379 that the involvement of Ti in the chemical system produces an unrealistic decrease of modal 380 proportions of some key Ti-free minerals (i.e. garnet) that have already relatively low modal proportion 381 382 in the rock. However, the investigated mylonites contain a small amount of biotite (i.e. these strongly deformed rocks reached temperatures close to the Bt-out line; Fig. 10). Because TiO₂ plays a 383 fundamental role in extending the stability field of biotite in P-T space (Tajčmanová et al., 2009 and 384 385 references therein), this component must be included in the phase equilibria modeling in order to obtain more realistic P-T estimates. The proportion of melt predicted by the modeling at 820 °C and 6 kbar is 386 ≈10-12 vol.%. 387

388

389 7. Discussion

390 7.1 Mylonites as former diatexites

391 The zonal arrangement of MI in garnet (Figs. 4, 5) indicates that they were trapped during host growth -i.e. they are primary MI (Roedder, 1984; Frezzotti, 2001). The investigated rocks are strongly 392 393 deformed and, therefore, the classic microstructures indicating the former presence of melt, such as 394 mineral pseudomorphs after melt films and pools (see Holness and Sawyer, 2008; Holness et al., 2011), are generally absent. Only rare crystallized pools of melt have survived in strain shadows (Fig. 3e). 395 396 However, the presence of abundant primary MI sheltered by the host garnet, which is resistant to deformation, clearly indicates the former occurrence of melt in the investigated mylonites. The quartzo-397 feldspathic mylonites represent, therefore, former migmatites. The leucocratic bands, showing a coarser 398 399 grain size, mainly composed of a granitic assemblage of plagioclase, K-feldspar and quartz and

containing some igneous microstructures such as euhedral minerals (Fig. 3j), likely represent former 400 401 anatectic leucosomes. The occurrence of a slightly more albitic plagioclase in these leucocratic portions with respect to the mesocratic domains is consistent with their crystallization from a melt (e.g. Sawyer, 402 2001). We interpret that the quartzo-feldspathic mylonites represent former diatexites, based on the 403 following observations and arguments: i) the similarity in bulk rock composition between these 404 mylonites, the underlying quartzo-feldspathic diatexites, and stromatic metatexites at the bottom of the 405 406 migmatitic sequence (Table 1); ii) their structural position above the diatexitic migmatites and at the contact between the migmatitic sequence and the peridotites (Fig. 2); iii) the presence of simply 407 twinned K-feldspar porphyroclasts (Fig. 3d) resembling the large K-feldspar observed in the diatexites 408 409 but absent in the metatexites (Acosta, 1998); iv) the abundance of leucocratic bands (30-50 vol.%) likely representing former leucosomes (Fig. 3a); and v) the apparently continuous evolution in terms of 410 petrography, mineral proportions and chemistry, and MI compositions across the migmatitic sequence 411 (Sections 4 and 5). 412

In this perspective, however, it can be noted that the maximum amount of melt predicted by 413 phase equilibria modeling, ≈10-12 vol.% (Fig. 10c), is neither enough to form a diatexite (i.e., a melt-414 supported structure) nor consistent with field observations suggesting \approx 30-50 vol.% of leucosomes 415 (Fig. 3a; see also Acosta, 1998). In anatectic terranes, the passage from metatexite to diatexite 416 417 migmatite is commonly transitional, and schollen diatexites are expected to form at relatively low to moderate melt fractions of ≈ 25 vol.%, and in some cases even at values as low as 0.16 (see Figure 1 in 418 Sawyer, 2008). Assuming that the protolith of the studied mylonites is similar to that of quartzo-419 420 feldspathic metatexites at the bottom of the migmatitic sequence (as field, bulk rock composition, mineral and MI data strongly suggest), and considering the difference in the amount of biotite between 421 the stromatic metatexites (see Bartoli et al., 2013c) and the mylonite (Fig. 10c), mass-balance 422 considerations indicate that ≈15-20 vol.% of a H₂O-undersaturated melt should have been produced 423

from the protolith of the mylonites (H₂O in the melt \approx 3 wt% as suggested by MI). This indicates that the thermodynamic model may somehow underestimate the proportion of melt produced in the mylonites.

Two additional explanations for the difference in melt proportion predicted by thermodynamic 427 models versus indicated by the volume of leucosomes, are the influx of a hydrous fluid enhancing melt 428 production (White et al., 2005) or the infiltration of melt from external sources (Hasalová et al., 2008). 429 Water-fluxed melting is not supported by different lines of evidence described in section 7.2 (see 430 below). On the other hand, Tubía et al. (1997) have reported (though not documented) the intrusion of 431 granitic melts into the mylonitic package, as roughtly concordant granite bodies parallel to S_{myl}. 432 433 Summarizing, field, compositional and textural arguments support that mylonites represent strongly deformed former diatexites; the discrepancy between calculated and observed amount of melt can be 434 ascribed in part to the limitations of thermodynamic modeling (see discussion in Section 7.3) and to the 435 infiltration of an external melt. 436

437

438 7.2 *P*-*T* estimates of melting and mylonitization close to the peridotites

439 P-T conditions at which the MI were trapped and, in turn, the quartzo-feldspathic mylonites previously melted, are constrained at \approx 820 °C and \approx 6 kbar (Fig. 10b) by the intersection among the 440 441 relevant compositional isopleths for MI-bearing garnet cores. The presence of sillimanite as trapped phase within MI proves that peritectic garnet and melt were produced in the field of sillimanite, in 442 agreement with petrographic observations and phase equilibria modelling. Biotite replacing garnet (Fig. 443 444 3g) likely represents the product of melt-consuming reactions (Kriegsman and Hensen, 1998). In particular, the concomitant decrease and increase of X_{Mg} in garnet rims and adjacent biotite grains, 445 respectively, indicate the progress of retrograde Fe-Mg exchange by diffusion ("ReERs" of Kohn and 446 Spear 2000) between garnet and biotite in mutual contact (Fig. 3g, h). Because their compositions have 447

been modified by occurrence of ReERs, their isopleths cannot be used for inferring the retrograde path. 448 After anatexis, the investigated rocks were strongly deformed down to subsolidus conditions 449 (phase D₄ of Tubía, 1988). According to Stipp et al. (2002) who studied natural deformation 450 microstructures of guartz over a temperature range of ≈ 500 °C, the formation of chessboard subgrains 451 (Fig. 3h) may have occurred at 600-650 °C. It is important to note that quartz deformation 452 microstructures are not only temperature dependent, and additional factors (presence of fluid, strain 453 454 rate and strain partitioning) may have an important effect (Mancktelow and Pennacchioni, 2004; Peternell et al., 2010). For example, the presence of H₂O may decrease the estimated temperature of 455 \approx 100 °C (Little et al., 2013). The CPO (crystallographic preferred orientation) patterns of quartz from 456 457 quartzo-feldspathic mylonites of the Ojén unit have been studied in detail by Tubía et al. (2013). Although they did not discuss the role of other variables in the development of quartz CPO (compare 458 Stipp et al., 2002, and Peternell et al., 2010), they concluded that the observed quartz fabrics support 459 that the mylonitization of the quartzo-feldspathic rocks occurred at HT (>500-600 °C). Pressure, 460 however, is not well constrained. 461

Because subsolidus penetrative deformation affected the entire rock, it is important to 462 understand its role on the documented microstruture and composition of MI. Some nanogranitoids 463 display diametrically opposite decrepitation tails (Fig. 7b), resembling the microstructures shown by 464 465 fluid inclusions experimentally deformed by deviatoric stress (see Tarantola et al., 2010). Decrepitation cracks are usually filled with the same minerals observed within nanogranitoids, indicating that 466 decrepitation took place before the beginning of melt crystallization (Ferrero et al., 2012). 467 468 Decrepitation may have induced fluid leakage from nanogranitoids (Cesare et al., 2011) and may have triggered crystallization by causing a pressure drop and, in turn, a drop in solubility within MI (Ferrero 469 et al., 2012). However, the coexisting glassy MI do not show evidence of decrepitation, but rather a 470 well-developed regular shape (Fig. 7a). The absence of microstructures indicative of decrepitation and 471

472 overheating in glassy MI (cf. Figure 9 in Cesare et al., 2015) suggests that their EMP analyses reported
473 in Table 2 can be considered as reliable compositions of the primary melt formed during the prograde
474 melting of the host rocks.

475

476 *7.3 Considerations on the thermodynamic modeling of anatexis*

Phase equilibria modeling predicts that after crossing the solidus and with increasing 477 temperature, the modal proportions of melt and garnet increase whereas the amount of biotite decreases 478 (Fig. 10c). This indicates that partial melting occurred by a continuous melting reaction consuming 479 biotite up to the Bt-out curve (Fig. 10) and also supports the peritectic nature of MI-bearing garnet. An 480 481 intriguing aspect of our phase equilibria modeling is that a small amount (≤0.01 vol.%) of COH fluid is present as long as biotite is stable above the solidus, and disappears in Bt-free assemblages (Fig. 10a). 482 Such a pattern on the stability of COH fluids is due to an artifact of the modeling under the imposed 483 bulk chemical constraints. In fact, since the O₂ component is involved in the thermodynamic 484 calculations, and since biotite is allowed to contain 10-15% of iron as Fe³⁺ (Tajčmanová et al., 2009), 485 in order to conserve mass Perple X removes from the fluid phase the amount of oxygen required for 486 the Fe₂O₃ component of biotite, leading to the attainment of the general conditions $X_0 < 1/3$ –i.e. an 487 imperceptible amount of a CH₄-rich fluid (≤ 0.01 vol.%) is forced to be present in Bt-bearing 488 489 supersolidus assemblages. These conditions persist until biotite is totally consumed. At this point, since the melt model used does not account for the solubility of carbonic species, and since biotite is the only 490 carrier of Fe^{3+} in the phase equilibria modeling, X_0 returns to the input value of 1/3, maximizing the 491 492 amount of H₂O dissolved in the melt and resulting in the consumption of the all free H₂O available in the system and in the precipitation of a small amount of graphite. Although such a process may seem 493 petrologically tenable, the presence of a CH₄-rich fluid predicted by this modeling is clearly in contrast 494 495 with the common occurrence of CO₂-rich fluids in high-grade graphitic metamorphic terranes (e.g.,

Touret, 2009; Hollister, 1988; Cesare et al., 2007; Ferrero et al., 2011; La Madrid et al., 2014; Santosh 496 497 and Omori, 2008). On the contrary, CH₄ has been detected in a very few cases (e.g., Lamb et al., 1991). There are, however, independent petrologic lines of evidence suggesting that partial melting of 498 the investigated rocks largely occurred in the absence of a fluid phase. They include: i) the H₂O content 499 of melt, estimated by difference from the EMP analyses of the glassy MI, is very close to the values 500 predicted for H₂O-undersaturated melting at \approx 820 °C and \approx 6 kbar (\approx 4 wt%; Holtz et al., 2001); ii) the 501 502 average ASI of the investigated MI is 1.20, and corresponds to the value predicted by equilibrium experiments for H₂O-undersaturated melts derived from fluid-absent incongruent melting of biotite 503 (Acosta-Vigil et al., 2003); iii) no primary fluid inclusions were found coexisting with the MI in the 504 505 investigated rocks; iv) the lack of Na- and Ca-rich melts typically produced by H₂O-fluxed melting of metasedimentary rocks (Weinberg and Hasalová, 2015). For example, MI showing tonalitic-506 trondhjemitic-granodioritic compositions with H₂O contents between ≈ 8 and ≈ 15 wt%, and produced 507 by H₂O-present melting have been recently documented in migmatites at the base of the Greater 508 Himalayan Sequence (Himalaya, Nepal) and in the Jubrique sequence (Betic Cordillera, Spain) (Carosi 509 et al., 2015; Acosta-Vigil et al., 2016). We therefore conclude that the extent melt + fluid field in Fig. 510 10a has been overestimated by the thermodynamic modeling. 511

Owing to the above problems in the modeling and in order to verify the reliability of the P-T512 513 estimates obtained in the C-bearing system, a new P-T section for the former diatexite of Fig. 10a has been calculated in a C-free system (Fig. 11). The displacement of Ms- and Bt-out curves is negligible 514 and the phase assemblage Grt-Bt-Sil-Pl-Kfs-Qtz-Ilm-Liq observed in the sample still corresponds to 515 516 the quadrivariant field in the upper right corner of the phase diagram section (Fig. 11). Instead, the shift of the solidus towards lower temperatures is significant at P < 5kbar (≈ 60 °C at 4 kbar and ≈ 100 °C at 2 517 kbar; Fig. 11), as expected in non-graphitic systems (Cesare et al., 2003). The P-T conditions of 518 equilibration during melting of the investigated rocks, however, do not differ in C-bearing and C-free 519

systems (\approx 820 °C and \approx 6 kbar; Fig. 11 and Fig. 1 in online supporting material). The most important difference between C-free and C-bearing systems is the extension of the *P*-*T* area of coexistence of melt + fluid. In the C-free phase diagram, melt and H₂O coexist only on the H₂O-saturated solidus. After crossing the solidus, the system almost instantaneously evolves towards a fluid-absent state in which further melting proceeds by fluid-absent (i.e., hydrate-breakdown) melting reactions, supporting the inference that melting largely occurred in absence of a fluid phase.

526 In Figure 12, we compared the compositions of MI with the calculated melt composition obtained from the thermodynamic modeling at the *P*-*T* conditions of interest. MI generally show higher 527 FeO+MgO and #K, and lower CaO and Al₂O₃. The inconsistency between calculated and measured 528 529 melt composition was already pointed out by Bartoli et al. (2013c) for stromatic metatexites from Ojén unit and by Grant (2009) and White et al. (2011) for experimentally remelted metapelites and 530 metagreywackes, and has been ascribed to the current melt model which needs some improvements to 531 reproduce properly natural processes (see discussion in Bartoli et al., 2013c). In the same figure, we 532 also plotted experimental glasses produced by melting of metagreywackes at 810-850 °C, 5-7 kbar. 533 Experimental melt compositions only partly overlap those of MI from Ojén metagreywackes and 534 formed at similar *P*-*T* conditions (Fig. 12). Because the composition of the source and conditions of 535 anatexis play a primary control on melt chemistry (Neogi et al., 2014), only MI, rather than 536 537 experiments and thermodynamic modeling, can make accessible the precise melt composition for the specific rock and the specific P-T- X_{H2O} investigated (Bartoli et al., 2013, 2016; Cesare et al., 2015). 538

539

540 7.4 P-T estimates and evolution of the Ojén unit: open questions

541 Tubía (1988) and Tubía et al. (1997, 2013) have documented a structural continuity throughout Ojén

unit, in terms of: (i) close orientation of foliations (strike from N10°E to N90°E and dip gently to

543 moderately south), stretching lineations (trends varying from NNE-SSW to ENE-WSW) and fold axes

parallel to the stretching lineation in the several rock types and at different structural levels; and (ii) similar evolution in time of the kinematics and localization of shear zones throughout the sequence of mylonites and migmatites, i.e. earlier shear zones are higher T, top-to-the ENE, and located at the contact with the peridotites, whereas later shear zones are lower T, top-to-the ENE and top-to-the-NNW and located at or towards the contact between the anatectic sequence and schists.

Field observations indicate a progressive evolution in the morphological type of migmatite, from 549 550 stromatic and fold-structured migmatites at the bottom of the sequence, to diatexite migmatites towards the middle-upper part of the section, where schollen, nebulitic and schlieric diatexitic migmatites have 551 been described (Fig. 2; Acosta, 1998). Mylonites at the top of the sequence represent former diatexites 552 553 (this study; see also Tubía, 1988; Acosta, 1998). Comparing quartzo-feldspathic rocks of similar bulk rock composition but varying metamorphic degree (Table 1), and in the frame of the phase equilibria 554 modeling (section 6, Figs. 10, 11; see also Bartoli et al., 2013c), variations in mineral proportions, 555 microstructures (see Section 4.1; compare Figs. 3 and 4), mineral and MI compositions (see Sections 556 4.2 and 5) and in melting reactions (from fluid-present, muscovite- and biotite-consuming to fluid-557 absent biotite dehydration melting; Bartoli et al., 2013c, 2015; this study) indicate an increase in T 558 towards the top of the anatectic sequence (Fig. 13, and references in the caption). 559

Considering the metasedimentary crustal rocks, which constitute most of the sequence of Ojén 560 561 (Fig. 2), it is clear also that P increases towards upper structural levels, (Fig. 13). Thus, a pressure of \approx 8-9 kbar reported for the mylonitic gneisses at the contact with the peridotites (Westerhof, 1975, 562 1977; Tubía et al., 1997) is in agreement with the occurrence of rutile trapped within MI in garnet 563 564 found in these rocks (Bartoli et al., 2015), indicating the presence of anatectic melt in the rutile stability field. By contrast, lower pressures of 4.5-5 kbar at the bottom of the migmatitic sequence are reflected 565 by the presence of peak MI-bearing Ilm crystals within the quartzo-feldspathic metatexites (Bartoli et 566 al., 2015). This is consistent with an increase towards upper structural levels of the Grt/Crd modal ratio 567

in rocks of similar bulk rock composition (Acosta, 1998). In addition, petrology and microstructures 568 also indicate that after reaching the peak *P*-*T* conditions, crustal rocks both at the contact with the 569 peridotite and at the bottom of the migmatitic sequence experienced a decompression path (Fig. 13). 570 This is clearly indicated, on the one hand, by formation of cordierite after garnet in pelitic granulites 571 and previously reported decompression P-T paths for mylonites and amphibolites/eclogites (Tubía et 572 al., 1997); and, on the other hand, by the presence of cordierite-bearing neosomes obliterating the main 573 574 foliation in the stromatic metatexites and postdating formation of garnet -bearing leucosomes (Figs. 2, 4, Section 4.1; see Acosta, 1998). The distribution and evolution of the *P*-*T* conditions documented at 575 the different levels of the Ojén unit is difficult to explain considering a metamorphic continuity 576 577 throughout the sequence and current petrogenetic models associated with the emplacement of a hot peridotite slab during continental subduction (Tubía et al., 1997, 2013; Mazzoli and Martín-Algarra, 578 2011, 2014). 579

Another important problem is that pressures ≥ 15 Kbar recorded in eclogitic boudins at the base of 580 the anatectic sequence (Tubía and Gil-Ibarguchi, 1991) contrast with both the much lower values 581 recorded in their host metatexites (Fig. 2), and the upward increase in P shown by rocks in the crustal 582 sequence (Fig. 13). These mafic rocks form rounded- to lenticular-shaped, cm-to-dm, or even 100-m 583 thick, bodies parallel to the main foliation (Tubía, 1998), and their presence has been used by Tubía et 584 585 al. (1997, 2013) and Mazzoli and Martín-Algarra (2011, 2014) to argue that the Ojén unit represents a coherent high-P portion of an Alpine continental subduction system. The presence of pods, boudins 586 and lenses of eclogites within host metasedimentary rocks equilibrated at lower P-T conditions is a 587 588 common occurrence in orogenic belts worldwide (e.g. Carswell, 1990). These eclogite lenses have been explained either as (i) foreign tectonic slices brought together with the crustal rocks late in their 589 metamorphic evolution, or (ii) rocks that, having been assembled together with the crustal rocks early 590 and having experienced the same tectonometamorphic evolution, can record earlier metamorphic stages 591

592 593

that are erased in their crustal hosts. In the latter case, although lower than in eclogites, pressures recorded in the metasediments are generally high as well (\geq 15-20 kbar; Li et al., 2015).

In the case of the Ojén unit, the presence of eclogite relicts, together with the decompression paths 594 recorded by the crustal rocks (Fig. 13), might suggest that maximum pressures recorded in the crustal 595 rocks represent retrograde pressures at which rocks equilibrated at some point during a decompression 596 path from HP peak values (see Tubía et al., 1997). Several lines of evidence, however, are not in 597 accordance this hypothesis. (1) Pressures exceeding 9 kbar have never been reported in the Ojén 598 metasedimentary sequence, particularly close to the eclogite lenses (Fig. 13). (2) Metatexites are Ca-599 poor, Si-rich peraluminous metagreywacke (Table 1), very similar in composition to peraluminous 600 601 granites, for which phase equilibria modeling and petrographic observations indicate mineral associations made of Ca-rich Grt+Ky+Rt at the *P*-*T* recorded in the eclogites of Ojén (e.g., Tajčmanová 602 et al., 2006; O'Brien, 2008; Massonne, 2009; Nahodilová et al., 2011). These minerals have never 603 been described or reported in these rocks, whereas HP metagranitic rocks (i.e., felsic granulites) that 604 experienced decompression down to 4-5 kbar and subsequent re-equilibration commonly preserve 605 relics of the original HP mineral assemblage (e.g., Liu and Zhong, 1997; O'Brien and Rotzler, 2003; 606 Tajčmanová et al., 2011). (3) The host metatexites have been well characterized in terms of 607 petrography, petrology and P-T conditions (Acosta, 1998; Bartoli, 2012; Bartoli et al., 2013b, c; this 608 609 work), and all these detailed studies indicate that mineralogy and microstructures in these rocks correspond to those generated during the prograde history and at peak conditions, and not to retrograde 610 associations after HP conditions. (4) Melt inclusions in garnet of rocks from the migmatitic sequence 611 612 where trapped during prograde melting reactions, and there is a systematic evolution in the composition of the MI from the metatexites to the mylonites (Bartoli et al., 2015; this work), which is in accordance 613 with the prograde *P*-*T* history inferred from mineralogy, microstructures and comparison with phase 614 equilibria modeling. In addition, in metatexites close to mafic boudins zircon overgrowths, interpreted 615

to have crystallized from anatectic melts, have been dated at ≈ 290 Ma (sample NDA / in S	Sanchez-
---	----------

Rodríguez, 1998), whereas the eclogite protolith formed at \approx 184 Ma (Sánchez-Rodríguez, 1998;

618 Sánchez-Rodríguez and Gebauer, 2000).

Because Tubía et al. (2013) have described late metamorphic mylonitic bands close to the amphibolite/eclogite lenses, from all the above considerations we suggest that the high–P mafic rocks and low–P migmatites may have been assembled together in the crustal footwall after their peak metamorphism. Similarly, Štípská et al. (2008) demonstrated that the contrasting peak pressures of eclogite lenses and host migmatites from Bohemian Massif (Czech Republic) are the result of the tectonic mixing of rocks originally coming from different depths, rather than the exhumation of a coherent HP terrane affected by heterogeneous retrogression.

Additional problems in the area relate to the available *P*-*T* data, showing a large variability and several inconsistencies such as: i) *P*-*T* paths that cross each other at lower P, e.g. eclogites and pelitic granulites, (ii) peak *P*-*T* values in pelitic granulites being \approx 50-100 °C lower and 2-3 kbar higher with respect to those in the underlying quartzo-feldspathic mylonites, and iii) *P*-*T* estimates for schists that are similar to, or higher than, those for the overlying migmatitic metatexites (Fig. 13). All these considerations raise doubts about the reliability of several of the thermobarometric estimates currently available in the literature for these rocks.

633

634 8. Regional implications

Most of the previous models have explained the tectono-metamorphic evolution of the Ojén and
Guadaiza units as due to the oblique subduction of a continental margin in a transpressional setting in
the early Miocene (i.e., the emplacement of Ronda peridotites occurred during continental subduction
as a result of oblique plate convergence; Tubía et al., 1997, 2013; Mazzoli and Martín-Algarra, 2011,
2014). Accordingly, Ojén would have been subducted to deeper levels (≥50 Km) beneath the

subcontinental mantle represented today by the Ronda peridotites, acquiring the high pressure
metamorphic stage recorded in the eclogites. Conversely, the Guadaiza unit would have remained at
mid-to-low crustal level depths. Alternatively, taking into account the petrologic and geochronological
constraints proposed by Esteban et al. (2008, 2011a) for the Guadaiza unit at Sierra Bermeja, Précigout
et al. (2013) provided numerical results supporting the emplacement of Ronda peridotites in a back-arc
basin during the earliest Miocene (see also Garrido et al., 2011).

646 This study on the Ojén unit has raised some important issues which are not easy to explain by some of these previously proposed models. Among them, i) the crustal footwall at Sierra Alpujata represents 647 an inverted metamorphic sequence, not only in terms of temperature but also in pressure, and ii) the 648 Ojén unit does not seem to represent a coherent high–pressure portion of a continental subduction 649 system. Regarding the generation and preservation of the inverted temperature gradients of the Ojén 650 unit, Tubía et al. (1997) called for a combination of high temperatures in the overlying peridotites and 651 rapid exhumation, as argued for many metamorphic soles of ophiolites. However, this model is rather 652 general and does not provide a detailed explanation of the occurrence of lower peak pressure values 653 654 towards the bottom of the sequence. Moreover, a very recent study of the Guadaiza unit has shown that a rather large portion of this unit records Variscan or older mineral assemblages and structures (Acosta-655 Vigil et al., 2014). Recent advances in the thermomechanical modeling (Moulas et al., 2014; 656 657 Tajčmanová et al., 2015) documenting the pressure build-up in and around bodies with a high viscous contrast, such as peridotites and migmatites studied here, might explain the pressure variation towards 658 the top of the sequence. In addition, the zone of high strain rate may result in temperature increase due 659 to viscous heating (e.g. Schmalholz and Podladchikov, 2013). 660

It follows that we are still far from understanding the origin, and geodynamic implications, of the tectono-metamorphic evolution of the western Alpujárride units, and particularly those located beneath the Ronda peridotite slab. In order to constrain the timing and mechanism of the crustal

- emplacement of the Ronda peridotites, future research on crustal rocks of the western Alpujárrides 664
- should focus on detailed petrological and geochronological studies of individual rock types within 665
- these crustal sequences, with particular emphases on those techniques that permit establishing a clear 666
- link between *P*-*T* conditions and ages. 667
- 668

Acknowledgements 669

670 This research benefitted from funding from the Italian Ministry of Education, University, Research

(Progetto SIR RBSI14Y7PF to OB and grant PRIN 2010TT22SC to BC), from Padova University 671

(Progetti per Giovani Studiosi 2013 to OB and Progetto di Ateneo CPDA107188/10 to BC), from 672 Società Italiana di Mineralogia e Petrologia (grant for a research stav abroad to OB), from the

- 673
- Ministerio de Ciencia e Innovación of Spain (grant CGL2007-62992 and a Ramón y Cajal research 674 contract to AAV), from MADE-IN-EARTH ERC starting grant (n. 335577) to LT and from the U.S.
- 675 National Science Foundation (Grant No. EAR-1019770) to RJB. The research leading to these results 676
- has received funding from the European Commission, Seventh Framework Programme, under Grant 677

Agreement nº 600376. We are grateful to F. Rossetti and an anonymous reviewer for constructive 678

- comments. We also thank A Risplendente (Milan University) for assistance during EMP analyses. A. 679
- Cavallo (INGV, Rome) for help with SEM observations and S. McCartney (ICTAS, Virginia Tech) for 680 assistance during elemental mapping. J.A.D. Connolly and G. Pennacchioni are also acknowledged for 681
- their helpful discussion on phase equilibria modeling and quartz microstructures, respectively. 682
- 683 684

685 References

- Acosta, A., 1998. Estudio de los fenómenos de fusión cortical y generación de granitoides asociados a 686 las peridotitas de Ronda. Unpublished PhD Thesis, Universidad de Granada, p. 305. 687
- Acosta-Vigil, A., Buick, I., Hermann, J., Cesare, B., Rubatto, D., London, D., Morgan VI, G.B., 2010. 688 Mechanisms of crustal anatexis: a geochemical study of partially melted metapelitic enclaves and 689 690 host dacite, SE Spain. Journal of Petrology 51, 785-821.
- Acosta-Vigil, A., Cesare, B., London, D., Morgan, G.B. VI, 2007. Microstructures and composition of 691 melt inclusions in a crustal anatectic environment, represented by metapelitic enclaves within El 692 Hoyazo dacites, SE Spain. Chemical Geology 235, 450–465. 693
- Acosta-Vigil, A., London, D., Morgan VI, G.B., Dewers, T.A., 2003. Solubility of excess alumina in 694 hydrous granitic melts in equilibrium with peraluminous minerals at 700–800 °C and 200 MPa, and 695

applications of the aluminium saturation index. Contributions Mineralogy Petrology 146, 100–119. 696

- Acosta-Vigil, A., Pereira, M.D., Shaw, D.M., London, D., 2001. Contrasting behaviour of boron during 697 698 crustal anatexis. Lithos 56, 15-31.
- Acosta-Vigil, A., Rubatto, D., Bartoli, O., Cesare, B., Meli, S., Pedrera, A., Azor, A., Tajčmanová, L., 699 2014. Age of anatexis in the crustal footwall of the Ronda peridotites, S Spain. Lithos 210-211, 700 147-167. 701
- Acosta-Vigil, A., Barich, A., Bartoli, O., Garrido, C.J., Cesare, B., Remusat, L., Poli, S., Raepsaet, C., 702 2016. The composition of nanogranitoids in migmatites overlaying the Ronda peridotites (Betic 703
- Cordillera, S Spain): the anatectic history of a polymetamorphic basement. Contributions to 704

- 705 Mineralogy and Petrology 171:24.
- Barich, A., Acosta-Vigil, A., Garrido, C.J., Cesare, B., Tajčmanová, L., Bartoli O., 2014.
- Microstructures and petrology of melt inclusions in the anatectic sequence of Jubrique (Betic
 Cordillera, S Spain): Implications for crustal anatexis. Lithos, 206–207, 303–320.
- Bartoli, O., 2012. When the continental crust melts: a combined study of melt inclusions and classical
 petrology on the Ronda migmatites. Unpublished PhD Thesis, Università degli Studi di Parma, 128
 pp.
- Bartoli, O., Cesare, B., Poli, S., Acosta-Vigil, A., Esposito, R., Turina, A., Bodnar, R.J., Angel, R.J.,
 Hunter, J., 2013a. Nanogranite inclusions in migmatitic garnet: behavior during piston cylinder remelting experiments. Geofluids 13, 405–420.
- Bartoli, O., Cesare, B., Poli, S., Bodnar, R.J., Acosta-Vigil, A., Frezzotti, M.L., Meli, S., 2013b.
 Recovering the composition of melt and the fluid regime at the onset of crustal anatexis and S-type granite formation. Geology 41, 115–118.
- Bartoli, O., Tajčmanová, L., Cesare, B., Acosta-Vigil, A., 2013c. Phase equilibria constraints on
 melting of stromatic migmatites from Ronda (S. Spain): insights on the formation of peritectic
 garnet. Journal of Metamorphic Geology 31, 775–789.
- Bartoli, O., Cesare, B., Remusat, L., Acosta-Vigil, A., Poli, S., 2014. The H₂O content of granite
 embryos. Earth and Planetary Science Letters 395, 281–290.
- Bartoli, O., Acosta-Vigil, A., Cesare, B., 2015. High-temperature metamorphism and crustal melting:
 working with melt inclusions. Periodico di Mineralogia84, 3B, 591–614..
- Bartoli, O., Acosta-Vigil, A., Ferrero, S., Cesare, B., 2016. Granitoid magmas preserved as melt
 inclusions in high-grade metamorphic rocks. American Mineralogist doi:10.2138/am-2016 5541CCBYNCND
- Becker, A., Holtz, F., and Johannes, W., 1998. Liquidus temperatures and phase compositions in the
 system Qz-Ab-Or at 5 kbar and very low water activities. Contributions to Mineralogy and
 Petrology 130, 213–224.
- Carswell, D.A., 1990. Eclogites and eclogite facies: definitions and classifications. In: Carswell, D.A.
 (Ed.), Eclogite Facies Rocks. Blackie and Son Ltd, New York, 1–13.
- Carosi, R., Montomoli, C., Langone, A., Turina, A., Cesare, B., Iaccarino, S., Fascioli, L., Visonà, D.,
 Ronchi, A., Man Rai, S., 2015. Eocene partial melting recorded in peritectic garnets from kyanite-
- gneiss, Greater Himalayan Sequence, central Nepal. In: Mukherjee, S., Carosi, R., van der Beek, P.
 A., Mukherjee, B. K. & Robinson, D. M. (Eds.), Tectonics of the Himalaya. Geological Society,
 London, Special Publications 412, pp. 111–129.
- Cesare, B., Acosta-Vigil, A., Ferrero, S., Bartoli, O., 2011. Melt Inclusions in migmatites and
 granulites. In: Forster, M.A., Fitz Gerald, J.D. (Eds.), The Science of Microstructure Part II.
 Journal of Virtual Explorer, (ISSN 1441-8142) 38, paper 2, Electronic edition.
- Cesare, B., Acosta-Vigil, A., Bartoli, O., Ferrero, S., 2015. What can we learn from melt inclusions in
 migmatites and granulites?. Lithos 239, 186–216.
- Cesare, B., Ferrero, S., Salvioli-Mariani, E., Pedron, D., Cavallo, A., 2009. Nanogranite and glassy
 inclusions: the anatectic melt in migmatites and granulites. Geology 37 627–630.
- Cesare, B., Maineri, C., Baron Toaldo, A., Pedron, D. & Acosta-Vigil, A., 2007. Immiscibility between
 carbonic fluids and granitic melts during crustal anatexis: a fluid and melt inclu- sion study in the
 enclaves of the Neogene Volcanic Province of SE Spain. Chemical Geology 237, 433–449.
- 748 Cesare, B., Marchesi, C., Hermann, J., Gomez-Pugnaire, M.T., 2003. Primary melt inclusions in
- andalusite from anatectic graphitic metapelites: implications for the position of the Al₂SiO₅ triple
- 750 point. Geology 31, 573–576.

- Coggon, R., Holland, T.J.B., 2002. Mixing properties of phengitic micas and revised garnet–phengite
 thermobarometers. Journal of Metamorphic Geology 20, 683–696.
- Connolly, J.A.D., 1995. Phase diagrams for graphitic rocks. Contributions to Mineralogy and Petrology
 119, 94–116.
- Connolly, J.A.D., 2009. The geodynamic equation of state: what and how. Geochemistry, Geophysics,
 Geosystems 10, Q10014.
- Connolly, J.A.D., Cesare, B., 1993. C-O-H-S fluid composition and oxygen fugacity in graphitic
 metapelites. Journal of Metamorphic Geology 11, 379–388.
- Cuevas, J., Esteban, J.J., Tubía, J.M., 2006. Tectonic implications of the granite dyke swarm in the
 Ronda peridotites (Betic Cordilleras, Southern Spain). Journal of the Geological Society 163, 631–
 640.
- Dewey, J.F., Helman, M.L., Turco, E., Hutton, D.H.W., Knott, S.D., 1989. Kinematics of the Western
 Mediterranean. In: Coward, M.P., Dietrich, D., Park, R.G. (Eds.), Alpine Tectonics. Special
 Publication, Geological Society of London, pp. 265–283.
- Esteban, J.J., Cuevas, J., Vegas, N., Tubía, J.M., 2008. Deformation and kinematics in a melt-bearing
 shear zone from the Western Betic Cordilleras (Southern Spain). Journal of Structural Geology 30,
 380–393.
- Esteban, J.J., Cuevas, J., Tubía, J.M., Sergeev, S., Larionov, A., 2011a. A revised Aquitanian age for
 the emplacement of the Ronda peridotites (Betic Cordilleras, southern Spain). Geological Magazine
 148, 183–187.
- Esteban, J.J., Tubía, J.M., Cuevas, J., Vegas, N., Sergeev, S., Larionov, A., 2011b. Peri-Gondwanan
 provenance of pre- Triassic metamorphic sequences in the western Alpujarride nappes (Betic
 Cordillera, southern Spain). Gondwana Research 20, 443–449.
- Ferrero, S., Bodnar, R.J., Cesare, B., Viti, C., 2011. Reequilibration of primary fluid inclusions in
 peritectic garnet from metapelitic enclaves, El Hoyazo, Spain. Lithos, 124, 117–131.
- Ferrero, S., Bartoli, O., Cesare, B., Salvioli-Mariani, E., Acosta-Vigil, A., Cavallo, A., Groppo, C.,
 Battiston, S., 2012. Microstructures of melt inclusions in anatectic metasedimentary rocks. Journal
 of Metamorphic Geology 30, 303–322.
- Frezzotti, M.L., 2001. Silicate melt inclusions in magmatic rocks: applications to petrology. Lithos 55,
 273–99.
- Frezzotti, M.L., Ferrando, S., 2015. The chemical behavior of fluids released during deep subduction
 based on fluid inclusions. American Mineralogist 100, 352–377.
- Garrido, C.J., Gueydan, F., Booth-Rea, G., Precigout, J., Hidas, K., Padron-Navarta, J.A., Marchesi, C.
 2011. Garnet lherzolite and garnet- spinel mylonite in the Ronda peridotite: vestiges of Oligocene
 backarc mantle lithospheric extension in the western Mediterranean. Geology 39, 927–930.
- Grant, J.A., 2009. THERMOCALC and experimental modeling of melting of pelite, Morton Pass,
 Wyoming. Journal of Metamorphic Geology 27, 571–578.
- Hasalová, P., Štípská, P., Powell, R., Schulmann, K., Janoušek, V., Lexa, O., 2008. Transforming
 mylonitic metagranite by open-system interactions during melt flow. Journal of Metamorphic
 Geology 26, 55–80.
- Holland, T.J.B., Powell, R., 1998. An internally consistent thermodynamic data set for phases of
 petrological interest. Journal of Metamorphic Geology 16, 309–343.
- Holland, T.J.B., Powell, R., 2003. Activity-composition rela- tions for phases in petrological
 calculations: an asymmetric multicomponent formulation. Contributions to Mineralogy and
 Petrology 145, 492–501.
- Hollister, L.S., 1988. On the origin of CO₂-rich fluid inclusions in migmatites. Journal of Metamorphic
 Geology 6, 467–474.

- Holness, M.B., Sawyer, E.W., 2008. On the pseudomorphing of melt-filled pores during the
 crystallization of migmatites. Journal of Petrology 49, 1343–1363.
- Holness, M.B., Cesare, B., Sawyer, E.W., 2011. Melted rocks under the microscope: microstructures
 and their interpretation. Elements, 7, 245–250.
- Holtz, F., Johannes, W., Tamic, N., Behrens, H., 2001. Maximum and minimum water contents of
 granitic melts gener- ated in the crust: a re-evaluation and implications. Lithos, 56, 1–14.
- Johnson, T.E., White, R.W., Powell, R., 2008. Partial melting of metagreywacke: a calculated mineral
 equilibria study. Journal of Metamoprhic Geology 26, 837–853.
- Kohn, M.J., Spear, F., 2000. Retrograde net transfer reaction insurance for pressure-temperature
 estimates. Geology 28/12, 1127-1130.
- Kretz, R., 1983. Symbols for rock-forming minerals. American Mineralogist 68, 277–279.
- Kriegsman, L.M., Hensen, B.J., 1998. Back reaction between restite and melt: Implications for
 geothermobarometry and pressure-temperature paths. Geology 26, 1111–1114.
- Lamadrid, H.M., Lamb, W.M., Santosh, M., Bodnar, R.J., 2014. Raman spectroscopic characterization
 of H₂O in CO₂-rich fluid inclusions in granulite facies metamorphic rocks. Gondwana Research 26,
 301–310.
- Lamb, W.M., Brown, P.E., Valley, J.W., 1991. Fluid inclusions in Adirondack granulites: implications
 for the retrograde P–T path. Contributions to Mineralogy and Petrology 107, 472–483.
- Li, J-L., Klemd, R., Gao, J., Jiang, T., Song, Y-H., 2015. A common high-pressure metamorphic
 evolution of interlayered eclogites and metasediments from the 'ultrahigh-pressure unit' of the
 Tianshan metamorphic belt in China. Lithos 226, 169–182.
- Little, T.A., hacker, B.R., brownlee, S.J., Seward, G., 2013. Microstructures and quartz lattice preferred orientations in the eclogite-bearing migmatitic gneisses of the D'Entrecasteaux Islands,
 Papua New Guinea. Geochemistry Geophysics Geosystems 14/6, doi:10.1002/ggge.20132.
- Liu, Y., Zhong, D., 1997. Petrology of high-pressure granulites from the eastern Himalayan syntaxis.
 Journal of Metamorphic Geology 15. 451–466.
- Loomis, T.P., 1972. Diapiric emplacement of the Ronda high-temperature ultramafic intrusion,
 southern Spain. Geological Society of America Bulletin 83, 2475–2496.
- Lundeen, M.T., 1978. Emplacement of the Ronda peridotite, Sierra Bermeja, Spain. Geological Society
 of America Bulletin 89, 172–180.
- Mancktelow, N.S., Pennacchioni, G., 2004. The influence of grain boundary fluids on the microstructure of quartz-feldspar mylonites. Journal of Structural Geology 26, 47–69.
- Massonne, H.-J., 2009. Hydration, dehydration, and melting of metamorphosed granitic and dioritic
 rocks at high- and ultrahigh-pressure conditions. Earth and Planetary Science Letters 288, 244–254.
- Massonne, H.-J., 2014. Wealth of P-T-t information in medium-high grade metapelites: example from
 the Jubrique Unit of the Betic Cordillera, S Spain. Lithos 208–209, 137–157.
- Mazzoli, S., Martín Algarra, A., 2011. Deformation partitioning during transpressional emplacement of
 a 'mantle extrusion wedge': the Ronda peridotites, western Betic Cordillera, Spain. Journal of the
 Geological Society, London 168, 373–382.
- Mazzoli, S., Martín Algarra, A., 2014. Comment on: "Localization of deformation and kinematic shift during the hot emplacement of the Ronda peridotites (Betic Cordilleras, southern Spain)" by J.M.
 Tubía, J. Cuevas, and J.J. Esteban, Journal of Structural Geology 50 (2013), 148–160. Journal of Structural Geology 60, 97–101.
- Michard, A., Goffé, B., Bouybaouène, M.L., Saddiqi, O., 1997. Late Hercynian–Mesozoic thinning in
 the Alboran domain: metamorphic data from the northern Rif, Morocco. Terra Nova 9, 171–174.
- Montel, J.M., Vielzeuf, D., 1997. Partial melting of metagreywackes, Part II. Compositions of minerals
- and melts. Contributions to Mineralogy and Petrology 128, 176–196.

- Morgan, G.B., London, D., 1996. Optimizing the electron microprobe analysis of hydrous alkali
 aluminosilicate glasses. American Mineralogist, 81 1176–1185.
- Morgan, G.B., London, D., 2005. Effect of current density on the electron microprobe analysis of alkali
 aluminosilicate glasses. American Mineralogist 90, 1131–1138.
- Moulas, E., Burg, J.P., Podladchikov, Y., 2014. Stress field associated with elliptical inclusions in a
 deforming matrix: mathematical model and implications for tectonic overpressure in the lithosphere.
 Tectonophysics 631, 37–49.
- Nahodilová, R., Faryad, S.W., Dolejš, D., Tropper, P., Konzett, J., 2011. High-pressure partial melting
 and melt loss in felsic granulites in the Kutná Hora complex, Bohemian Massif (Czech Republic).
 Lithos 125, 641–658.
- Navarro-Vilá, F., Tubía, J.M., 1983. Essai d'une nouvelle différenciation des Nappes Alpujárrides dans
 le secteur occidental des Cordillères Bétiques (Andalousie, Espagne). Comptes Rendus de
 l'Académie des Sciences Paris 296, 111–114.
- Neogi, S., Bolton, E.W., Chakraborty, S., 2014. Timescales of disequilibrium melting in the crust:
 constraints from modelling the distribution of multiple trace elements and a case study from the
 Lesser Himalayan rocks of Sikkim. Contributions to Mineralogy and Petrology 168:1020.
- Newton, R.C., Charlu, T.V. & Kleppa, O.J., 1980. Thermo- chemistry of high structural state
 plagioclases. Geochimica et Cosmochimica Acta, 44, 933–941.
- Obata, M., 1980. The Ronda peridotite: garnet-, spinel-, and plagioclase-lherzolite facies and the P-T
 trajectories of a high temperature mantle intrusion. Journal of Petrology, 21, 533–572.
- O'Brien, P.J., 2008. Challenges in high-pressure granulite metamorphism in the era of pseudosections:
 reaction textures, compositional zoning and tectonic interpretation with examples from the
 Bohemian Massif. Journal of Metamorphic Geology 26, 235–251.
- O'Brien, P.J., Rötzler, J., 2003. High-pressure granulites: formation, recovery of peak conditions and
 implications for tectonics. Journal of Metamorphic Geology 21, 3–20.
- Patiño Douce, A.E., Beard, J.S., 1996. Effects of P, *f*(O₂) and Mg/Fe ratio on dehydration melting of
 model metagreywackes. Journal of Petrology 37, 999–1024.
- Pattison, D.R.M., 1992. Stability of andalusite and sillimanite and the Al_2SiO_5 triple point, constraints from the Ballachulish aureole, Scotland. Journal of Geology 100, 423–446.
- Peternell, M., Hasalová, P., Wilson, C.J.L., Piazolo, S., Schulmann, K., 2010. Evaluating quartz
 crystallographic preferred orientations and the role of deformation partitioning using EBSD and
 fabric analyser techniques. Journal of Structural Geology 32, 803–817.
- Platt, J.P., Argles, T.W., Carter, A., Kelley, S.P., Whitehouse, M.J., Lonergan, L., 2003. Exhumation of
 the Ronda peridotite and its crustal envelope: Constraints from thermal modelling of a P–T–time
 array. Journal of the Geological Society of London 160, 655–676.
- Platt, J.P., Behr, W.M., Johanesen, K., Williams, J.R., 2013. The Betic-Rif arc and its orogenic
 hinterland: A review. Annual Review of Earth and Planetary Sciences 41, 14.1–14.45.
- Platt, J.P., and Vissers, R.L.M., 1989. Extensional collapse of thickened conti- nental lithosphere: A
 working hypothesis for the Alboran Sea and Gibraltar arc. Geology 17, 540–543.
- Précigout, J., Gueydan, F., Garrido. C.J., Cogné, N., Booth-Rea, G., 2013. Deformation and
 exhumation of the Ronda peridotite (Spain). Tectonics 32, 1–15.
- Priem, H.N.A., Boelrijk, N.A.I.M., Hebeda, E.H., Oen, I.S., Verdurmen, E.A.Th., Verschure, R.H.,
 1979. Isotopic dating of the emplacement of the ultramafic masses in the Serranía de Ronda,
- 888 Southern Spain. Contributions to Mineralogy and Petrology 70, 103–109.
- Roedder, E., 1984. Fluid inclusions. Mineralogical Society of America, Reviews in Mineralogy, 12,
 644.
- 891 Sánchez-Navas, A., García-Casco, A., Martín-Algarra, A., 2014. Pre-Alpine discordant granitic dikes

- in the metamorphic core of the Betic Cordillera: tectonic implications. Terra Nova 26/6, 477–486.
- Sánchez-Rodríguez, L., 1998. Pre-Alpine and Alpine evolution of the Ronda Ultramafic Complex and
 its country-rocks (Betic chain, southern Spain): U–Pb SHRIMP zircon and fission-track dating.
 (PhD Thesis) ETH, Zürich, p. 170.
- Sánchez-Rodríguez, L., Gebauer, D., 2000. Mesozoic formation of pyroxenites and gabbros in the
 Ronda area (southern Spain), followed by Early Miocene subduction metamorphism and
 emplacement into the middle crust: U–Pb sensitive high-resolution ion microprobe dating of zircon.
 Tectonophysics 316, 19–44.
- Santosh, M., Omori, S., 2008. CO₂ flushing: a plate tectonic perspective. Gondwana Research 13, 86– 102.
- Sawyer, E.W., 2001. Melt segregation in the continental crust: distribution and movement of melt in
 anatectic rocks. Journal of Metamorphic Geology, 19, 291–309.
- Sawyer, E.W., 2008. Atlas of migmatites. Mineralogical Association of Canada. The Canadian
 Mineralogist Special Publication 9, Quebec City.
- Schmalholz, S.M., Podladchikov, Y.Y., 2013. Tectonic overpressure in weak crustal-scale shear zones
 and implications for the exhumation of high-pressure rocks. Geophysical Research Letters 40,
 1984–1988.
- Spear, F.S., 1993. Metamorphic Phase Equilibria and Pressure–Temperature–Time Paths.
 Mineralogical Society of America, Washington DC, 799 pp.
- Stipp, M., Stünitz, H., heilbronner R., Schmid, S.M. 2002. The eastern Tonale fault zone: a 'natural laboratory' for crystal plastic deformation of quartz over a temperature range from 250 to 700 °C.
 Journal of Structural Geology 24, 1861–1884.
- Štípská, P., Schulmann, K., Powell, R., 2008. Contrasting metamorphic histories of lenses of highpressure rocks and host migmatites with a flat orogenic fabric (Bohemian Massif, Czech
 Republic): a result of tectonic mixing within horizontal crustal flow?. Journal of Metamorphic
 Geology 26, 623–646.
- Tajčmanová, L., Abart, R., Neusser, G., Rhede, D., 2011. Growth of decompression plagioclase rims
 around metastable kyanite from high-pressure felsic granulites (Bohemian Massif). Journal of
 Metamorphic Geology, 29, 1003–1018.
- Tajčmanová, L., Conolly, J.A.D., Cesare, B., 2009. A thermodynamic model for titanium and ferric
 iron solution in biotite. Journal of Metamorphic Geology 27, 153–165.
- Tajčmanová, L., Konopásek, J., Schulmann, K., 2006. Thermal evolution of the orogenic lower crust
 during exhumation within a thickened Moldanubian root of the Variscan belt of Central Europe.
 Journal of Metamorphic Geology 24,119–134.
- Tajčmanová, L., Vrijmoed, J. Moulas, E., 2015. Grain-scale pressure variations in metamorphic rocks:
 implications for the interpretation of petrographic observations. Lithos 216-217, 338–351.
- Tarantola, A., Diamond, L.W., Stünitz, H., 2010. Modification of fluid inclusions in quartz by
 deviatoric stress I: experimentally induced changes in inclusion shapes and microstructures.
 Contributions to Mineralogy and Petrology 160, 825–843.
- Thompson, A.B., Algor, J.R., 1977. Model systems for anatexis of pelitic rocks. I. Theory of melting
 reactions in the system KAlO₂-NaAlO₂-Al₂O₃-SiO₂-H₂O. Contributions to Mineralogy and
 Petrology 63, 247–269.
- Thompson, J.B., Hovis, G.L., 1979. Entropy of mixing in sanidine. American Mineralogist 64, 57–65.
- Tinkham, D.K., Zuluaga, C.A., Stowell, H.H., 2001. Metapelite phase equilibria modeling in
 MnNCKFMASH: the effect of variable Al₂O₃ and MgO/(MgO+FeO) on mineral stability.
- 937 Geological Materials Research 3, 1–42.
- 938 Torres-Roldán, R.L., 1983. Fractionated melting of metapelite and further crystal-melt equillibria. The

- example of the Blanca Unit migmatite complex, north of Estepona (southern Spain). Tectonophysics
 96, 95–123.
- Touret, J.L.R., 2009. Mantle to lower-crust fluid/melt transfer through granulite metamorphism.
 Russian Geology and Geophysics 50, 1052–1062.
- Tubía, J.M., 1988. Estructura de los Alpujárrides occidentales: cinemática y condiciones de
 emplazamiento de las peridotitas de Ronda. Publicaciones Especiales del Boletín Geológico y
 Minero de España. 99, p. 124.
- Tubía, J.M., Gil-Ibarguchi, J.I., 1991. Eclogites of the Ojen nappe: a record of subduction in the
 Alpujarride complex (Betic Cordilleras, southern Spain). Journal of the Geological Society 148,
 801–804.
- Tubía, J.M., Cuevas, J., Gil-Ibarguchi, J.I., 1997. Sequential development of the metamorphic aureole
 beneath the Ronda peridotites and its bearing on the tectonic evolution of the Betic Cordillera.
 Tectonophysics 279, 227–252.
- Tubía, J.M., Cuevas, J., Esteban, J.J., 2013. Localization of deformation and kinematics shift during the
 hot emplacement of the Ronda peridotites (Betic Cordilleras, southern Spain). Journal of Structural
 Geology 50, 148–160.
- Villaros, A., Stevens, G., Moyen, J..F, Buick, I.S. 2009. The trace element compositions of S-type
 granites: evidence for disequilibrium melting and accessory phase entrainment in the source.
 Contributions to Mineralogy and Petrology 158, 543–61.
- Weinberg, R.F., Hasalovà, P., 2015. Water-fluxed melting of the continental crust: a re- view. Lithos
 212–215, 158–188.
- Westerhof, A.B., 1975. Genesis of a magnetite ore near Marbella, Southern Spain: Formation by
 oxidation of silicates in polymetamorphic gedrite-bearing and other rocks. PhD Thesis, University
 of Amsterdam, G.U.A. papers of Geology 6, 216 pp.
- Westerhof, A.B., 1977. On the contact relations of high-temperature peridotites in the Serrania de
 Ronda, southern Spain. Tectonophysics 39, 579–591.
- Wilke, S., klahn, C., Bolte T., Almeev, R., Holtz, F., 2015. Experimental investigation of the effect of
 Ca, Fe and Ti on cotectic compositions of the rhyolitic system. European Journal of Mineralogy
 27/2, 147–159.
- White, R.W., Pomoroy, N.E., Powell. R., 2005. An in situ metatexite–diatexite transition in upper
 amphibolite facies rocks from Broken Hill, Australia. Journal of Metamorphic Geology 23, 579–
 602.
- White, R.W., Powell, R., Holland, T.J.B., 2007. Progress relating to calculation of partial melting
 equilibria for metapelites. Journal of Metamorphic Geology 25, 511–527.
- White, R.W., Stevens, G., Johnson, T.E., 2011. Is the crucible reproducible? Reconciling melting
 experiments with thermodynamic calculations. Elements 7, 241–246
- Zeck, H.P., Monie[´], P., Villa, I., Hansen, B.T., 1992. Very high rates of cooling and uplift in the Alpine
 belt of the Betic Cordilleras, southern Spain. Geology, 20, 79–82.
- 977 978
- 979 Captions
- 980

Fig. 1. (a) Location map of the study area on the Iberian Peninsula in southern Spain. (b) Simplified

- geological map of the western sector of the Betic Cordillera (modified after Esteban et al., 2011b). (c)
- 983 Geological map of the Sierra Alpujata massif. Blue and yellow stars show the location of the studied
- mylonites (N 36°36'31.9", W4°48'21.7") and of the stromatic metatexites investigated by Bartoli et al.
 (2013c), respectively.

986

Fig. 2. Schematic cross-section based on work of Tubía (1988) and Acosta (1998) of the crustal footwall of the Ronda peridotites at Sierra Alpujata showing the location of the studied mylonitic samples (blue star as in Figure 1). The photomicrographs (4.2x2.2 mm) show the microstructural evolution of migmatites as a function of distance from the bottom of the Ronda peridotite slab. Red arrows show the location of peritectic Grt. Yellow lines show the traces of the main foliation defined by biotite and/or sillimanite folia.

- 993
- 994

995 **Fig. 3.** Outcrop views (a, b) and photomicrographs (c-l) of the investigated rocks. (a) Quartzofeldspathic mylonites outcropping at Sierra Alpujata with leucocratic bands (interpreted as former 996 leucosomes) parallel to main foliation. Coin = 2 cm. (b) Close-up of garnet-bearing leucosome (with tip 997 of pen pointing to the garnet). (c) Mesocratic matrix showing fabric-forming sillimanite, biotite and 998 ilmenite (Sil>>Bt>Ilm). White arrows: graphite lamellae. (d) K-feldspar porphyroclast showing simple 999 twinning. (e) Quartz with irregular outlines containing euhedral Pl and Bt. This domain has probably 1000 crystallized from a pool of melt (see Holness et al., 2011). (f) Garnet shows embayment and 1001 replacement by Bt. Red dotted square: MI cluster. (g, h) Characteristic deformation microstructures of 1002 1003 quartz formed by grain boundary migration recrystallization and showing cuspate-lobate boundaries and irregular subgrains (g) or approximately square (i.e., chessboard) subgrains (h). (k, l) 1004 Photomicrographs of two leucosomes. In (g) feldspars and quartz are deformed and elongated, and 1005 scattered in a fine-grained matrix, whereas in (h) there is a dense framework of euhedral feldspars 1006 1007 (white arrows) touching along faces.

1007

Fig. 4. Outcrop view (a) and photomicrographs (b, c) of the quartzo-feldspathic metatexites collected
towards the bottom of the migmatitic sequence (see yellow star in Figure 2). (a) Small cordieritebearing leucogranitic patches obliterate the main foliation. (b) The main foliation is defined by
abundant oriented biotite grains, and garnet occurs as small crystals. (c) Cordierite-bearing leucocratic
domains have microstructures dominated by crystallization of the melt, such as quartz with a interstitial
morphology (white asterisk) surrounding euhedral grains of feldspars (white arrows).

Fig. 5. Photomicrographs of the host garnets showing the different arrangements of MI clusters (red dotted lines). (a) MI cluster with a subspherically (equant) shape located near the garnet core. (b) MI cluster with a sigmoidal shape. (c) MI cluster with a spiral-like shape.

Fig. 6. (a) Cluster of crystallized (white arrows) and preserved glassy (red arrows) melt inclusions. (b)
Glassy MI containing a shrinkage bubble (red arrow). (c) Crystallized MI with a Sil needle (white arrow) that is likely to have favored the entrapment of melt.

Fig. 7. SEM-BSE images of melt inclusions in garnet with the typical negative crystal shape. (a) Coexistence of nanogranitoids (white arrows) and preserved glassy MI (gray arrows) in the same cluster. (b) Fully crystallized inclusions (i.e. nanogranitoids). Black arrows: decrepitation cracks.

Fig. 8. Elemental EDS X-rays map of a nanogranitoid inclusion in mylonite ALP13.

Fig. 9. CIPW normative compositions of analyzed MI, shown on the Qtz-Ab-Or diagram (see text for details). *Dashed lines*: outline fields defined by the distributions of the retrieved compositions. *Black dots:* mean compositions. Compositions of MI from metatexites (*gray area*) are plotted for comparison

(data from Bartoli et al., 2013b). Black triangle and lines show eutectic point and cotectic lines for the 1033 subaluminous haplogranite system at 0.5 GPa and $a_{H2O} = 1$; black stars are eutectic points at $a_{H2O} = 0.6$ 1034 and 0.4 (Becker et al. 1998). The involvement of Fe, Ti and Ca moves eutectic points and cotectic 1035 curves toward more quartz-rich, albite-poor compositions (Wilke et al., 2015). 1036 1037 Fig. 10. (a) P–T section for mylonite ALP13calculated in MnNCaKFMASHTOC system. (b) Contours 1038 for grossular and spessartine components and X_{Mg} value of garnet, and for Ti content and X_{Mg} value of 1039 biotite. Yellow ellipse: inferred P-T conditions for formation of MI-bearing garnet cores. (c) Isopleths 1040 of modal proportions of biotite, garnet and melt. 1041 1042 1043 **Fig. 11.** *P*-*T* section in the system MnNCaKFMASHT for the mylonite composition. Important curves in the C-bearing system (Figure 10) are superimposed: dotted yellow line for Ms-out curve, dotted blue 1044 line for Bt-out curve and dotted red line for Lig-in curve. The gray field reflects the region of melt + 1045 1046 fluid coexistence in the C-bearing system. Yellow ellipse: inferred P-T conditions for MI-bearing garnet cores (see Fig. 1 in supporting online material). 1) Bt-Crd-Pl-Kfs-Ilm-Qtz-H₂O; 2) Bt-Crd-Pl-1047 Kfs-Ilm-And-Otz-H2O; 3) Bt-Kfs-Crd-Pl-Otz-Ilm-Sil-H2O 4) Bt-Pl-Kfs-Sil-Grt-Otz-H2O; 5) Bt-Kfs-1048 Crd-Pl-Qtz-Ilm-Sil-Liq. 1049 1050 Fig. 12. FeO+MgO vs. CaO and FeO+MgO vs. K# diagrams comparing compositions of melt 1051 inclusions, calculated melts and experimental glasses. K#=mol. K₂O/(Na₂O+K₂O). Light grey areas 1052 show the compositional domains corresponding to experimental melts produced by partial melting of 1053 metagreywackes at 810-850 °C, 5-7 kbar (data from Patiño Douce and Beard, 1996; Montel and 1054 Vielzeuf, 1997). Note that these glasses show CaO content up to 1.5 wt%; not shown in (a). See text for 1055 details. 1056 1057 **Fig. 13.** (a) Various *P*–*T* estimates derived from different rocks in the crustal footwall of the Ronda 1058 1059 peridotites at Sierra Alpujata (Ojén unit). P–T conditions for the quartzo-feldspathic mylonite are from this study. T97 refers to Tubía et al. (1997), TI91 to Tubía and Gil-Ibarguchi (1991), W77 to Westerhof 1060 (1977), B13c to Bartoli et al. (2013c), P92 to Pattison (1992). Areas with continuous lines refer to P 1061 and/or T peak conditions for each rock type. Mylonitic gneisses described by Tubía et al. (1997) 1062 correspond to pelitic granulites of Figure 2 and Section 4.1. (b) Simplified section reported in Figure 2, 1063 showing the relative stratigraphic position of the rocks reported in (a). See text for explanation. 1064 1065 Fig. 1. (supporting online material). (a) Contours for grossular and spessartine components and X_{Mg} 1066 value of garnet, and for Ti content and X_{Mg} value of biotite in the *P*-*T* section calculated in 1067 MnNCaKFMASHT system (see Fig. 10). Yellow ellipse: inferred *P*-*T* conditions for MI-bearing 1068 1069 garnet cores. (b) Isopleths of modal proportions of biotite, garnet and melt. 1070 **Tab. 1.** Electron microprobe analyses (wt%) of minerals from mylonite ALP13 and bulk rock 1071 composition of quartzo-feldspathic migmatites. 1072 1073 Tab. 2. Major element composition (wt%) of glassy melt inclusions. Numbers in parentheses refer to 1074

1075 1σ standard deviation.