


Cobalt-Catalyzed Cyclization of Unsaturated N-Acyl Sulfonamides: a Diverted Mukaiyama Hydration Reaction

Journal Article

Author(s):

Fischer, David M.; Balkenhohl, Moritz; [Carreira, Erick M.](#) 

Publication date:

2022-05-23

Permanent link:

<https://doi.org/10.3929/ethz-b-000553186>

Rights / license:

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#)

Originally published in:

JACS Au 2(5), <https://doi.org/10.1021/jacsau.2c00186>

Funding acknowledgement:

833540 - Development of Stereoselective Olefin Functionalization Methods (EC)

Cobalt-Catalyzed Cyclization of Unsaturated *N*-Acyl Sulfonamides: a Diverted Mukaiyama Hydration Reaction

David M. Fischer, Moritz Balkenhohl, and Erick M. Carreira*



Cite This: *JACS Au* 2022, 2, 1071–1077



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: The cycloisomerization of β -, γ -, and δ -unsaturated *N*-acyl sulfonamides to *N*-sulfonyl lactams and imidates is reported. This transformation is effected in the presence of a Co^{III} (salen) catalyst using *t*-BuOOH or air as the oxidant. The method shows good functional group tolerance (alkyl, aryl, heteroaryl, ether, *N*-Boc) and furnishes an underexplored class of cyclic building blocks. The strong solvent dependence of the transformation is investigated, and the synthetic versatility of the *N*-sulfonyl imidate product class is highlighted.


KEYWORDS: cycloisomerization, *N*-acyl sulfonamides, *N*-sulfonyl imidates, cobalt catalysis, diverted Mukaiyama hydration



Hydrofunctionalization of unactivated olefins using inexpensive first-row transition metals, such as Fe, Mn, Co, and Ni, has received considerable attention.^{1,2} Early examples of their application include earth-abundant metal-mediated olefin hydration³ and hydroperoxidation,⁴ pioneered by Drago⁵ and Mukaiyama,⁶ respectively. Since then there has been considerable evolution of olefin hydrofunctionalization to include oximation,⁷ cyanation,⁸ hydrazination,⁹ azidation,^{8b,10} amination,¹¹ and halogenation.^{8b,12} More recently, Shigehisa and others have demonstrated intermolecular olefin hydroalkoxylation,¹³ hydrofluorination,^{13b} intramolecular hydroarylation,¹⁴ hydrothioetherification,¹⁵ and hydroamination.^{13e,16} Herein we report the Co-catalyzed cycloisomerization of unsaturated *N*-acyl sulfonamides, which provides ready access to a wide range of cyclic *N*-sulfonyl imidates, an underexplored functional group (Scheme 1). In addition to examination of conditions with *t*-BuOOH, a salient feature of the process is our discovery that the reaction can be performed with air as a convenient and safe oxidant. Additionally, over the course of this study we observed a notable solvent effect that governs the product distribution, which we attribute to solvent donor ability and the attendant electronic character of the cobalt metal complex.

Imidates are a synthetically versatile class of compounds that act as activated amide equivalents through their dual behavior as both electrophiles and nucleophiles. Thus, they provide access to a wide range of structural motifs, including oxazolines, indazoles, and isoquinolines.¹⁷ Acyclic *N*-sulfonyl imidates have been investigated as surrogates for azaheterenes in steroidal antiproliferative agents.¹⁸ They have also been examined in drug discovery as prodrugs for esters and sulfonamides, addressing challenges associated with bioavailability and metabolism.¹⁹ While probenazole, a member of the class, has found applications as an antifungal agent against rice

Scheme 1. Cyclization of Alkenyl *N*-Acyl Sulfonamides



blast fungus and leaf blight, erythromycin derivatives modified through the incorporation of a cyclic imidate showed 4-fold increased activity against *Streptococcus pneumoniae*.²⁰ Cyclic *N*-sulfonyl imidates are less well studied, despite the fact that they could be useful in a prodrug approach as precursors toward lactones.

Cycloisomerizations of alkenyl *N*-acyl sulfonamides to *N*-sulfonyl lactams have been described, mediated by Ph_3PAuCl ,²¹ Ir catalysts,²² NbCl_5 ,²³ and $\text{Zn}(\text{OTf})_2/\text{TfOH}$.²⁴

Received: March 22, 2022


Revised: April 26, 2022

Accepted: April 28, 2022


Published: May 6, 2022



Prior Work

A. Cyclization of *N*-acyl sulfonamides

B. Classical reactivity: Mukaiyama Hydration



This Work

C. Key Observations




Figure 1. Background and initial observations.

(Figure 1A). Cyclic *N*-sulfonyl imidates have generally been accessed through electrophilic cyclization of *N*-acyl sulfonyl alkenes mediated by iodine,²⁵ *m*-CPBA,²⁶ Ph₂Se₂,²⁷ or TsN₃.^{28,29,26} The direct cycloisomerization of alkenyl *N*-acyl sulfonamides to furnish cyclic *N*-sulfonyl imidates is unknown.^{13e,16a}

We have been interested in identifying novel reactivity of olefins mediated by base metal–salen complexes.^{9a,c,10,30} In the presence of a Co^{II} catalyst, silane, and oxidant, olefins classically proceed through Mukaiyama hydration pathways (Figure 1B). When we subjected *N*-acyl sulfonamide **3a** to Mukaiyama hydration conditions, a mixture of products **I** and **II** was formed in 46% and 4% yield, respectively, in line with early observations by Mukaiyama.^{6a–c,f} Interestingly, when Co(acac)₂ was replaced by Co^{III}(salen)OTf complex **1** under otherwise identical conditions, the formation of *N*-sulfonyl imidate **4a** was observed (36%). Importantly, resubjecting **I** to Mukaiyama hydration conditions in the presence of catalyst **1** did not lead to **4a** (see the Supporting Information (SI)). When THF was replaced with toluene, **4a** was formed in 62% yield. Neither alcohol **I** nor ketone **II** was isolated from the reaction mixture. Collectively, these results compelled us to examine further the reaction in which Mukaiyama hydration of olefinic *N*-acyl sulfonamides is derailed to form cyclic imidates.

Although the cycloisomerization is isohypsic,³¹ it has been noted that for Co-mediated olefin functionalization reactions, the addition of hydroperoxide can be beneficial.^{6f,10a} Examination of the reaction conditions led to the identification of a procedure in toluene with 2 mol% **1**, *t*-BuOOH (2.2 equiv), and PhSiH₃ (2.2 equiv) that effects the cycloisomerization reaction **3a** → **4a** in 85% yield (Table 1, entry 1; see the SI). Control experiments revealed interesting details. No reactivity was observed in the absence of silane or catalyst **1** (entries 2 and 3). The use of a Co^{II}(salen-*t*Bu,*t*Bu) catalyst under otherwise identical conditions gave the product in merely 17% yield (entry 4; see the SI). Under an inert atmosphere, identical product yields were obtained (entry 5).

Table 1. Optimization of the Reaction Conditions

entry	deviation from the standard conditions	yield of 4a (%)
1	none	85
2	without PhSiH ₃	0
3	without 1	0
4	Co ^{II} (salen- <i>t</i> Bu, <i>t</i> Bu) instead of 1	17
5	under N ₂	84
6	air instead of <i>t</i> -BuOOH, 48 h	56
7	air, 10 mol% 1 , and 4 equiv of PhSiH ₃	82

Importantly, we found that the reaction could be conducted with air as the oxidant, leading to **4a** in 56% yield after 48 h (entry 6). Under conditions in which the catalyst loading was increased to 10 mol% with 4 equiv of silane and air, product **4a** was formed in 82% yield (entry 7). The fact that reaction conditions prescribe air and avoid the use of expensive or toxic oxidants makes this transformation attractive.³² Moreover, the use of air instead of oxygen renders the process safer.

With the optimized reaction conditions in hand, the scope and functional group tolerance of this transformation was investigated (Figure 2). Substrates incorporating a wide range of alkyl or aryl substituents, including rings at C_ω, were well-tolerated in the transformation, leading to products **4b–g** (62–83%). Ether-containing and *N*-Boc-amine-substituted alkenyl *N*-acyl sulfonamides readily underwent cycloisomerization and gave rise to products **4h–j** in 63–97% yield. β-Substituted olefins also underwent cyclization to give products **4k–m**. The reaction tolerated disubstituted olefins, allowing access to *N*-sulfonyl imidate **4n** in 55% yield.

For selected examples, namely, **4a–h**, **4j**, **4p**, and **4o**, the cycloisomerization reaction was repeated to showcase the broad applicability of air as the oxidant. The products were generally obtained in yields comparable to those under conditions employing *t*-BuOOH. The structures of the imidate products **4b**, **4h**, **4q**, and **4w** formed in this study were assigned by X-ray crystallography. For characterization of the remaining products, IR spectroscopy proved to be useful. *N*-Sulfonyl imidates display distinctive IR C=N absorbances at 1620 cm⁻¹, in stark contrast to ν_{C=O} = 1720 cm⁻¹ for *N*-tosyl lactams.

We next evaluated the formation of six-membered *N*-sulfonyl imidates. Gratifyingly, **3o–q** provided access to **4o–q** in 71–87% yield. Imidocarbonate **4r** was prepared in 52% yield from homoallylic *N*-sulfonyl carbamate **3r**. When **3s** was subjected to the reaction conditions, β-lactam **4s** was formed in 30% yield. The transformation was also amenable to 1,2-disubstituted olefins, with alkenyl *N*-acyl sulfonamide **3t** giving a mixture of five- and six-membered-ring products. In contrast, styrene derivative **4u** and diene **4v** were formed as single products. Finally, we established the feasibility of performing the reaction in a system with a trisubstituted olefin (**3w**), affording **4w** in 71% yield.

The effect of different sulfonamides on the transformation was investigated next (Figure 3). High yields were maintained for aryl sulfonamide substrates bearing arene groups




Figure 2. Substrate scope of the Co-catalyzed cycloisomerization of alkenyl *N*-acyl sulfonamides. ^aReaction yield obtained employing 10 mol% **1**, 4 equiv of PhSiH₃, and air. ^bThermal ellipsoids are shown at the 50% probability level.




Figure 3. Co-catalyzed cycloisomerization of a variety of sulfonamide and sulfonamide derivatives.

substituted with sterically demanding and electron-donating substituents (**6a** and **6b**). Methanesulfonamide derivative **5c** also produced the corresponding cyclic *N*-sulfonyl imidate **6c** in 88% yield. The use of olefinic *N*-acyl trifluoromethane sulfonamide **5d** yielded cycloisomerization products, albeit as a mixture of imidate **6d** and lactam **6e** in 30% and 25% yield, respectively. *N*-Sulfinyl imidate **6f** was formed in 68% yield, expanding the reaction to *N*-acyl sulfinamides.

The observation that the reaction conditions resemble those typically employed for Mukaiyama hydration^{6b} of olefins compelled us to conduct key control experiments (Figure 4A,B). In this respect, when **I** was subjected to either




Figure 4. Control experiments for the Co-catalyzed cycloisomerization reaction.

Mukaiyama hydration conditions or the reaction conditions, imidate **4a** was not formed, and the starting secondary alcohol was reisolated in 75 or 85% yield, respectively. This result precludes a mechanistic pathway for the overall transformation **3a** → **4a** involving Mukaiyama hydration of the olefin followed by ring closure. To rule out the possibility that the reaction is mediated by triflic acid formed from Co^{III}OTf catalyst **1**, starting material **3a** was exposed to a catalytic amount of

TfOH (10 mol%) (Figure 4C). No product formation was observed after 2 h, and **3a** was recovered in 75% yield.

Interestingly, there is paucity of data on the behavior of cyclic *N*-sulfonyl imidates. Accordingly, we conducted a brief study on **4a**. It displayed fast *E/Z* isomerization at room temperature as determined by HSQC and NOE NMR spectroscopy. The isomer (*E*)-**4a** prevails in toluene-*d*₈ (see the SI) on the basis of a cross-peak between the ring H₂C_α and *o*-CH_{aryl}. Cyclic imidates bearing C_α substituents crystallized solely as the *Z* isomer (see **4b**, **4h**, and **4q**). The barrier for *E/Z* isomerization of parent compound **4a** was determined to be $\Delta G^\ddagger = 16.0 \text{ kcal mol}^{-1}$ (Figure 5).³³




Figure 5. Determination of the barrier to *E/Z* isomerization in *N*-sulfonyl imidate **4a**. ¹H NMR spectra were recorded at 500 MHz in toluene-*d*₈ at -60 to 100 °C.

We then investigated derivatization of this underexplored product class (Figure 6). Compound **4p** was hydrolyzed to lactone **7a** in 88% yield (DMF–H₂O, 0.2 equiv of DBU, r.t., 12 h).^{29b,34} We wondered whether *N*-sulfonyl imidates could be employed as directing groups for arene C–H functionalization.




Figure 6. Selected functionalization of sulfonyl imidates.

The closest analogy we could find was in the work with Rh-catalyzed functionalization of cyclic *N*-sulfonyl ketimines.³⁵ Subjecting **4p** to $[(\text{Cp}^*)\text{RhCl}_2]_2$, AgSbF₆, Cu(OAc)₂, and methyl acrylate in 1,4-dioxane led to formation of *o*-C–H-alkenylated imidate **7b** (82% yield). Finally, reduction of **4p** with NaBH₄ gave amino alcohol **7c** in 77% yield.

The reactivity reported herein is intriguing and unexpected given previous studies involving cobalt complexes and simple olefins under similar conditions.^{9a,c,10,30b} A number of spectroscopic and computational investigations have suggested that canonical Co^{III}(salen) complexes such as **1** may be in equilibrium with species described as Co^{II}(salen^{•+}).³⁶ Various independent studies have separately indicated that equilibria involving Co species are sensitive to factors such as temperature, counterions, donor ligands, and solvent (CD₂Cl₂ vs DMSO-*d*₆ vs pyridine). Combined NMR and quantum-chemical studies revealed that Co(salen)Cl in THF-*d*₈ is present in diamagnetic and paramagnetic forms.³⁶

Our leading results outlined in Figure 1C hinted at a solvent effect (THF vs toluene), with the latter proving optimal for cyclization vis-à-vis the formation of Mukaiyama hydration/oxidation products I and II. Accordingly, we conducted a study of the cyclization reaction of **3a** with air as the oxidant over a range of solvents (Figure 7A). In proceeding from toluene to DMF, the relative amount of cyclization product decreases while the Mukaiyama products increase. A plot of the yield of cyclization product **4a** against solvent donor number³⁷ reveals a linear relationship (Figure 7; see the SI for details), with the yield of **4a** decreasing as a function of solvent coordination ability.³⁸ A recent report on olefin hydroamination reactions suggests a dependence of the product distribution *S* (cyclization vs Mukaiyama products) on solvent viscosity.^{16c} However, in the cyclization reactions of **3a** we did not observe such a dependence on the viscosity ($R^2 = 0.02$; see the SI); instead, once again a strong correlation ($R^2 = 0.94$) involving *S* and solvent donor ability was observed (Figure 7C).

For reactions involving Co complexes, silane, and an oxidant, it has been suggested in the literature that hydro-metalation of the starting olefin affords an organocobalt species.^{9a,c,10,13a–d,14,16a,30b} Our results are consistent with this putative intermediate partitioning itself between two possible pathways as a function of the solvent, namely, reaction with oxygen to form Mukaiyama hydration products I and II or, alternatively, a diverted course to give **4a**. It remains unclear whether the cyclization process we describe subsequently proceeds via cationic or radical intermediates.³⁹ In mechanistic inorganic studies of Co(salen) complexes,^{36a} Fuji has discussed the importance of the redox-active salen ligand in understanding the electronic structure of these complexes. This raises the question of whether the ability of canonical **1** to access its Co^{II}(salen^{•+}) form may be crucial to diverting the reaction from classical Mukaiyama reactivity to cycloisomerization. The design of complexes incorporating redox-non-innocent ligands for olefin functionalization reactions may provide new avenues for the identification of preparatively useful transformations.

In conclusion, we have reported the Co-catalyzed cycloisomerization of olefinic *N*-acyl sulfonamides employing *t*-BuOOH or air as the oxidant. The transformation was successful for a broad spectrum of olefin substrates and tolerated a variety of functional groups.⁴⁰ The barrier to *E/Z* isomerization of the imidates was determined by NMR coalescence experiments. We elaborated the imidate products




Figure 7. Solvent trends in the cycloisomerization of *N*-acyl sulfonamides. (A) Product distribution in various solvents. (B) Relationship between solvent donor number (Gutmann³⁷) and yield of **4a**. (C) Correlation between *S* ($=4a/(4a + I + II)$) and solvent donor number.

in further transformations, including cleavage of the sulfonamide, *N*-tosyl imidate reduction, and CH functionalization. Finally, we investigated the product distribution as a function of solvent, confirming that cycloisomerization is preferred over the traditional Mukaiyama hydration pathways in noncoordinating solvents. The access to cyclic *N*-sulfonyl imidate products provided by this method opens new possibilities for these structures as potentially useful building blocks in small-molecule discovery endeavors.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacsau.2c00186>.

Experimental and computational details and NMR data (PDF)

X-ray crystallographic data for **4w** (CIF)

X-ray crystallographic data for **4b** (CIF)

X-ray crystallographic data for **4h** (CIF)

X-ray crystallographic data for **4q** (CIF)

X-ray crystallographic data for **4s** (CIF)

■ AUTHOR INFORMATION

Corresponding Author

Erick M. Carreira – *ETH Zurich, 8093 Zurich, Switzerland*;

orcid.org/0000-0003-1472-490X;

Email: erickm.carreira@org.chem.ethz.ch

Authors

David M. Fischer – *ETH Zurich, 8093 Zurich, Switzerland*

Moritz Balkenhohl – *ETH Zurich, 8093 Zurich,*

Switzerland; orcid.org/0000-0002-3240-7708

Complete contact information is available at:

<https://pubs.acs.org/10.1021/jacsau.2c00186>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was funded by the European Research Council (OLECAT, Grant ID 833540). We are grateful to Dr. N. Trapp and M. Solar for X-ray crystallographic analysis, Dr. M.-O. Ebert and R. Falkenstein for low- and high-temperature NMR experiments, and T. Horak for starting material synthesis (all ETH Zurich). M.B. thanks the Deutsche Forschungsgemeinschaft (DFG) for a postdoctoral fellowship.

■ REFERENCES

- (1) (a) Crossley, S. W. M.; Obradors, C.; Martinez, R. M.; Shenvi, R. A. Mn-, Fe-, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev.* **2016**, *116*, 8912–9000. (b) Kyne, S. H.; Lefevre, G.; Ollivier, C.; Petit, M.; Ramis Cladera, V. A.; Fensterbank, L. Iron and cobalt catalysis: new perspectives in synthetic radical chemistry. *Chem. Soc. Rev.* **2020**, *49*, 8501–8542.
- (2) Ludwig, J. R.; Schindler, C. S. Catalyst: Sustainable Catalysis. *Chem.* **2017**, *2*, 313–316.
- (3) Lee, N.-H.; Byun, J.-C.; Baik, J.-S.; Han, C.-H.; Han, S.-b. Development of Mn(III)(Schiff Base) Complexes for the Catalyst of Olefin Oxygenation to Alcohols in the Presence of NaBH₄. *Bull. Korean Chem. Soc.* **2002**, *23*, 1365–1366.
- (4) (a) Matsushita, Y.-I.; Sugamoto, K.; Nakama, T.; Matsui, T. Synthesis of γ -hydroperoxy- α,β -unsaturated carbonyl compounds from $\alpha,\beta,\gamma,\delta$ -unsaturated carbonyl compounds by cobalt(II) porphyrin-catalyzed hydroperoxyoxygenation. *J. Chem. Soc., Chem. Commun.* **1995**, 567–568. (b) Sugamoto, K.; Matsushita, Y.-I.; Matsui, T. Direct hydroperoxyoxygenation of conjugated olefins catalyzed by cobalt(II) porphyrin. *J. Chem. Soc., Perkin Trans. 1* **1998**, 3989–3998. (c) Tokuyasu, T.; Kunikawa, S.; Masuyama, A.; Nojima, M. Co(III)–Alkyl Complex- and Co(III)–Alkylperoxy Complex-Catalyzed Triethylsilylperoxidation of Alkenes with Molecular Oxygen and Triethylsilane. *Org. Lett.* **2002**, *4*, 3595–3598. (d) O’Neill, P. M.; Hindley, S.; Pugh, M. D.; Davies, J.; Bray, P. G.; Park, B. K.; Kapu, D. S.; Ward, S. A.; Stocks, P. A. Co(thd)₂: a superior catalyst for aerobic epoxidation and hydroperoxysilylation of unactivated alkenes: application to the synthesis of spiro-1,2,4-trioxanes. *Tetrahedron Lett.* **2003**, *44*, 8135–8138. (e) Tokuyasu, T.; Kunikawa, S.; McCullough, K. J.; Masuyama, A.; Nojima, M. Synthesis of Cyclic Peroxides by Chemo- and Regioselective Peroxidation of Dienes with Co(II)/O₂/Et₃SiH. *J. Org. Chem.* **2005**, *70*, 251–260. (f) Hurlocker, B.; Miner, M. R.; Woerpel, K. A. Synthesis of silyl monoperoxyketals by regioselective cobalt-catalyzed peroxidation of silyl enol ethers: application to the synthesis of 1,2-dioxolanes. *Org. Lett.* **2014**, *16*, 4280–4283.

- (5) Zombeck, A.; Hamilton, D. E.; Drago, R. S. Novel catalytic oxidations of terminal olefins by cobalt(II)-Schiff base complexes. *J. Am. Chem. Soc.* **1982**, *104*, 6782–6784.
- (6) (a) Isayama, S.; Mukaiyama, T. Hydration of olefins with molecular oxygen and triethylsilane catalyzed by bis(trifluoroacetylacetonato)cobalt(II). *Chem. Lett.* **1989**, *18*, 569–572. (b) Inoki, S.; Kato, K.; Takai, T.; Isayama, S.; Yamada, T.; Mukaiyama, T. Bis(trifluoroacetylacetonato)cobalt(II) catalyzed oxidation-reduction hydration of olefins selective formation of alcohols from olefins. *Chem. Lett.* **1989**, *18*, 515–518. (c) Isayama, S.; Mukaiyama, T. A New Method for Preparation of Alcohols from Olefins with Molecular Oxygen and Phenylsilane by the Use of Bis(acetylacetonato)cobalt(II). *Chem. Lett.* **1989**, *18*, 1071–1074. (d) Inoki, S.; Kato, K.; Isayama, S.; Mukaiyama, T. A new and facile method for the direct preparation of α -hydroxycarboxylic acid esters from α,β -unsaturated carboxylic acid esters with molecular oxygen and phenylsilane catalyzed by Bis(dipivaloylmethanato)manganese(II) complex. *Chem. Lett.* **1990**, *19*, 1869–1872. (e) Mukaiyama, T.; Yamada, T. Recent advances in aerobic oxygenation. *Bull. Chem. Soc. Jpn.* **1995**, *68*, 17–35. (f) Isayama, S.; Mukaiyama, T. Novel method for the preparation of triethylsilyl peroxides from olefins by the reaction with molecular oxygen and triethylsilane catalyzed by bis(1,3-diketonato)cobalt(II). *Chem. Lett.* **1989**, *18*, 573–576.
- (7) (a) Okamoto, T.; Kobayashi, K.; Oka, S.; Tanimoto, S. Cobalt-catalyzed reaction of nitric oxide with aryl-substituted olefins in the presence of tetrahydroborate ion. *J. Org. Chem.* **1987**, *52*, S089–S092. (b) Okamoto, T.; Kobayashi, K.; Oka, S.; Tanimoto, S. A high-yield regioselective synthesis of keto oximes from aryl-conjugated ethylenes and ethyl nitrite in the presence of cobalt complex and BH_4^- ion. *J. Org. Chem.* **1988**, *53*, 4897–4901. (c) Kato, K.; Mukaiyama, T. A novel method for the preparation of 2-hydroxyiminocarboxylic acid esters. Cobalt (II) catalyzed α -oxidation of α, β -unsaturated ester with butyl nitrite and phenylsilane. *Chem. Lett.* **1990**, *19*, 1917–1920. (d) Kato, K.; Mukaiyama, T. Iron (III) complex catalyzed nitrosation of terminal and 1,2-disubstituted olefins with butyl nitrite and phenylsilane. *Chem. Lett.* **1992**, *21*, 1137–1140. (e) Sugamoto, K. A Synthesis of Oximes from Olefins by Cobalt(II) Porphyrin-Catalyzed Reduction-Nitrosation. *Synlett* **1998**, *1998*, 1270–1272. (f) Prateep-tongkum, S.; Jovel, I.; Jackstell, R.; Vogl, N.; Weckbecker, C.; Beller, M. First iron-catalyzed synthesis of oximes from styrenes. *Chem. Commun.* **2009**, 1990–1992.
- (8) (a) Gaspar, B.; Carreira, E. M. Mild Cobalt-Catalyzed Hydrocyanation of Olefins with Tosyl Cyanide. *Angew. Chem.* **2007**, *119*, 4603–4606. (b) Leggans, E. K.; Barker, T. J.; Duncan, K. K.; Boger, D. L. Iron(III)/ NaBH_4 -Mediated Additions to Unactivated Alkenes: Synthesis of Novel 20'-Vinblastine Analogues. *Org. Lett.* **2012**, *14*, 1428–1431.
- (9) (a) Waser, J.; Carreira, E. M. Convenient Synthesis of Alkylhydrazides by the Cobalt-Catalyzed Hydrohydrazination Reaction of Olefins and Azodicarboxylates. *J. Am. Chem. Soc.* **2004**, *126*, 5676–5677. (b) Sato, M.; Gunji, Y.; Ikeno, T.; Yamada, T. Stereoselective α -Hydrazination of α,β -Unsaturated Carboxylates Catalyzed by Manganese(III) Complex with Dialkylazodicarboxylate and Phenylsilane. *Chem. Lett.* **2005**, *34*, 316–317. (c) Waser, J.; González-Gómez, J. C.; Nambu, H.; Huber, P.; Carreira, E. M. Cobalt-Catalyzed Hydrohydrazination of Dienes and Enynes: Access to Allylic and Propargylic Hydrazides. *Org. Lett.* **2005**, *7*, 4249–4252. (d) Zheng, J.; Qi, J.; Cui, S. Fe-Catalyzed Olefin Hydroamination with Diazo Compounds for Hydrazone Synthesis. *Org. Lett.* **2016**, *18*, 128–131.
- (10) (a) Waser, J.; Nambu, H.; Carreira, E. M. Cobalt-Catalyzed Hydroazidation of Olefins: Convenient Access to Alkyl Azides. *J. Am. Chem. Soc.* **2005**, *127*, 8294–8295. (b) Carreira, E.; Gaspar, B.; Waser, J. Cobalt-Catalyzed Synthesis of Tertiary Azides from α,α -Disubstituted Olefins under Mild Conditions Using Commercially Available Reagents. *Synthesis* **2007**, *2007*, 3839–3845.
- (11) (a) Gui, J.; Pan, C.-M.; Jin, Y.; Qin, T.; Lo, J. C.; Lee, B. J.; Spergel, S. H.; Mertzman, M. E.; Pitts, W. J.; La Cruz, T. E.; et al. Practical olefin hydroamination with nitroarenes. *Science* **2015**, *348*, 886–891. (b) Obradors, C.; Martínez, R. M.; Shenvi, R. A. Ph(i-PrO)SiH₂: An Exceptional Reductant for Metal-Catalyzed Hydrogen Atom Transfers. *J. Am. Chem. Soc.* **2016**, *138*, 4962–71. (c) Zhu, K.; Shaver, M. P.; Thomas, S. P. Amine-bis(phenolate) Iron(III)-Catalyzed Formal Hydroamination of Olefins. *Chem. - Asian J.* **2016**, *11*, 977–80. (d) Ma, W.; Zhang, X.; Fan, J.; Liu, Y.; Tang, W.; Xue, D.; Li, C.; Xiao, J.; Wang, C. Iron-Catalyzed Anti-Markovnikov Hydroamination and Hydroamidation of Allylic Alcohols. *J. Am. Chem. Soc.* **2019**, *141*, 13506–13515.
- (12) (a) Gaspar, B.; Carreira, E. M. Catalytic hydrochlorination of unactivated olefins with para-toluenesulfonyl chloride. *Angew. Chem., Int. Ed.* **2008**, *47*, 5758–60. (b) Taniguchi, T.; Goto, N.; Nishibata, A.; Ishibashi, H. Iron-Catalyzed Redox Radical Cyclizations of 1,6-Dienes and Enynes. *Org. Lett.* **2010**, *12*, 112–115. (c) Ma, X.; Herzog, S. B. Non-classical selectivities in the reduction of alkenes by cobalt-mediated hydrogen atom transfer. *Chem. Sci.* **2015**, *6*, 6250–6255.
- (13) (a) Shigehisa, H.; Aoki, T.; Yamaguchi, S.; Shimizu, N.; Hiroya, K. Hydroalkoxylation of unactivated olefins with carbon radicals and carbocation species as key intermediates. *J. Am. Chem. Soc.* **2013**, *135*, 10306–9. (b) Shigehisa, H. Functional Group Tolerant Markovnikov-Selective Hydrofunctionalization of Unactivated Olefins Using a Cobalt Complex as Catalyst. *Synlett* **2015**, *26*, 2479–2484. (c) Shigehisa, H.; Hayashi, M.; Ohkawa, H.; Suzuki, T.; Okayasu, H.; Mukai, M.; Yamazaki, A.; Kawai, R.; Kikuchi, H.; Satoh, Y.; Fukuyama, A.; Hiroya, K. Catalytic Synthesis of Saturated Oxygen Heterocycles by Hydrofunctionalization of Unactivated Olefins: Unprotected and Protected Strategies. *J. Am. Chem. Soc.* **2016**, *138*, 10597–604. (d) Shigehisa, H.; Kikuchi, H.; Hiroya, K. Markovnikov-Selective Addition of Fluorous Solvents to Unactivated Olefins Using a Co Catalyst. *Chem. Pharm. Bull.* **2016**, *64*, 371–374. (e) For a singular example of an *N*-phenyl-*N*-tosylurea undergoing cyclization via oxygen, see: Nagai, T.; Mimata, N.; Terada, Y.; Sebe, C.; Shigehisa, H. Catalytic Dealkylative Synthesis of Cyclic Carbamates and Ureas via Hydrogen Atom Transfer and Radical-Polar Crossover. *Org. Lett.* **2020**, *22*, 5522–5527. (f) Ebisawa, K.; Izumi, K.; Ooka, Y.; Kato, H.; Kanazawa, S.; Komatsu, S.; Nishi, E.; Shigehisa, H. Catalyst- and Silane-Controlled Enantioselective Hydrofunctionalization of Alkenes by Cobalt-Catalyzed Hydrogen Atom Transfer and Radical-Polar Crossover. *J. Am. Chem. Soc.* **2020**, *142*, 13481–13490.
- (14) Shigehisa, H.; Ano, T.; Honma, H.; Ebisawa, K.; Hiroya, K. Co-Catalyzed Hydroarylation of Unactivated Olefins. *Org. Lett.* **2016**, *18*, 3622–3625.
- (15) Date, S.; Hamasaki, K.; Sunagawa, K.; Koyama, H.; Sebe, C.; Hiroya, K.; Shigehisa, H. Catalytic Direct Cyclization of Alkenyl Thioester. *ACS Catal.* **2020**, *10*, 2039–2045.
- (16) (a) Shigehisa, H.; Koseki, N.; Shimizu, N.; Fujisawa, M.; Niitsu, M.; Hiroya, K. Catalytic hydroamination of unactivated olefins using a Co catalyst for complex molecule synthesis. *J. Am. Chem. Soc.* **2014**, *136*, 13534–13537. The cyclization of two singular *N*-phenyl amides, 2-allyl-*N*-phenylbenzamide and 2-vinyl-*N*-phenylbenzamide, under HAT conditions to yield the corresponding cyclic *N*-phenyl imidates was reported herein. (b) Ohuchi, S.; Koyama, H.; Shigehisa, H. Catalytic Synthesis of Cyclic Guanidines via Hydrogen Atom Transfer and Radical-Polar Crossover. *ACS Catal.* **2021**, *11*, 900–906. (c) Ye, W.-T.; Zhu, R. Dioxxygen-promoted cobalt-catalyzed oxidative hydroamination using unactivated alkenes and free amines. *Chem. Catal.* **2022**, *2*, 345–357.
- (17) Thakur, R.; Jaiswal, Y.; Kumar, A. Imidates: an emerging synthon for *N*-heterocycles. *Org. Biomol. Chem.* **2019**, *17*, 9829–9843. Traditionally, acyclic *N*-sulfonyl imidates can be accessed through acid-catalyzed condensation of nitriles with alcohols (Pinner synthesis) followed by sulfonylation of the resultant imidate. Alternatively, they may be generated through dipolar cycloaddition of *N*-tosyl azide with alkynes in the presence of oxygen nucleophiles.
- (18) Volkova, Y. A.; Kozlov, A. S.; Kolokolova, M. K.; Uvarov, D. Y.; Gorbatov, S. A.; Andreeva, O. E.; Scherbakov, A. M.; Zavarzin, I. V. Steroidal *N*-Sulfonylimidates: Synthesis and biological evaluation in breast cancer cells. *Eur. J. Med. Chem.* **2019**, *179*, 694–706.

(19) Bundgaard, H.; Larsen, J. D. *N*-Sulfonyl imidates as a novel prodrug form for an ester function or a sulfonamide group. *J. Med. Chem.* **1988**, *31*, 2066–2069.

(20) (a) Iwata, M. Probenazole—a plant defence activator. *Pestic. Outlook* **2001**, *12*, 28–31. (b) Zhang, L.; Jiao, B.; Yang, X.; Liu, L.; Ma, S. Synthesis and antibacterial activity of new 4'-*O*-carbamates of 11,12-cyclic carbonate erythromycin A 6,9-imino ether. *J. Antibiot.* **2011**, *64*, 243–247.

(21) Liu, X.-Y.; Li, C.-H.; Che, C.-M. Phosphine Gold(I)-Catalyzed Hydroamination of Alkenes under Thermal and Microwave-Assisted Conditions. *Org. Lett.* **2006**, *8*, 2707–2710.

(22) (a) Nagamoto, M.; Yanagi, T.; Nishimura, T.; Yorimitsu, H. Asymmetric Cyclization of *N*-Sulfonyl Alkenyl Amides Catalyzed by Iridium/Chiral Diene Complexes. *Org. Lett.* **2016**, *18*, 4474–4477. (b) Nishimura, T.; Nagamoto, M.; Yorimitsu, H. Iridium-Catalyzed Intramolecular Oxidative Cyclization of Alkenyl Amides and Alkenoic Acids. *Synthesis* **2017**, *49*, 4272–4282.

(23) Ferrand, L.; Tang, Y.; Aubert, C.; Fensterbank, L.; Mouriès-Mansuy, V.; Petit, M.; Amatore, M. Niobium-Catalyzed Intramolecular Addition of O–H and N–H Bonds to Alkenes: A Tool for Hydrofunctionalization. *Org. Lett.* **2017**, *19*, 2062–2065.

(24) Chou, T.-H.; Yu, B.-H.; Chein, R.-J. $ZnI_2/Zn(OTf)_2$ -TsOH: a versatile combined-acid system for catalytic intramolecular hydrofunctionalization and polyene cyclization. *Chem. Commun.* **2019**, *55*, 13522–13525.

(25) Arai, T.; Watanabe, O.; Yabe, S.; Yamanaka, M. Catalytic Asymmetric Iodocyclization of *N*-Tosyl Alkenamides using Aminoiminophenoxy Copper Carboxylate: A Concise Synthesis of Chiral 8-Oxa-6-azabicyclo[3.2.1]octanes. *Angew. Chem., Int. Ed.* **2015**, *54*, 12767–12771.

(26) Takesue, T.; Fujita, M.; Sugimura, T.; Akutsu, H. A series of two oxidation reactions of ortho-alkenylbenzamide with hypervalent iodine(III): a concise entry into (3*R*,4*R*)-4-hydroxymellein and (3*R*,4*R*)-4-hydroxy-6-methoxymellein. *Org. Lett.* **2014**, *16*, 4634–4637.

(27) Cheng, X.; Hasimujiang, B.; Xu, Z.; Cai, H.; Chen, G.; Mo, G.; Ruan, Z. Direct Electrochemical Selenylation/Cyclization of Alkenes: Access to Functionalized Benzheterocycles. *J. Org. Chem.* **2021**, *86*, 16045–16058.

(28) Franz, J. E.; Dietrich, M. W.; Henshall, A.; Osuch, C. Reactions of Sulfonyl Azides and Sulfonamides with Vinyl Ethers. *J. Org. Chem.* **1966**, *31*, 2847–2853.

(29) (a) Neilson, D. G. Imidates including cyclic imidates. In *Amidines and Imidates*; John Wiley & Sons, 1991; pp 425–483. (b) Matsubara, R.; Berthiol, F.; Kobayashi, S. Sulfonylimidates as Nucleophiles in Catalytic Addition Reactions. *J. Am. Chem. Soc.* **2008**, *130*, 1804–1805.

(30) (a) Du Bois, J.; Hong, J.; Carreira, E. M.; Day, M. W. Nitrogen Transfer from a Nitridomanganese(V) Complex: Amination of Silyl Enol Ethers. *J. Am. Chem. Soc.* **1996**, *118*, 915–916. (b) Waser, J.; Carreira, E. M. Catalytic hydrohydrazination of a wide range of alkenes with a simple Mn complex. *Angew. Chem., Int. Ed.* **2004**, *43*, 4099–102. (c) Balkenhohl, M.; Kölbl, S.; Georgiev, T.; Carreira, E. M. Mn- and Co-Catalyzed Aminocyclizations of Unsaturated Hydrazones Providing a Broad Range of Functionalized Pyrazolines. *JACS Au* **2021**, *1*, 919–924.

(31) Hendrickson, J. B. Systematic characterization of structures and reactions for use in organic synthesis. *J. Am. Chem. Soc.* **1971**, *93*, 6847–6854.

(32) This is in contrast to previous methods employing *N*-fluorocollidinium salts as oxidants, which either require the reaction to be carried out under an inert atmosphere (hydrothioetherification, acetoxylation) or even prescribe degassing with argon to avoid the formation of Mukayama hydration (hydrofluorination, hydrohydroxylation, and hydroamination) products.

(33) (a) Gunther, H. *NMR Spectroscopy: Basic Principles, Concepts and Applications in Chemistry*, 3rd ed.; John Wiley & Sons, 2013. (b) Huggins, M. T.; Kesharwani, T.; Buttrick, J.; Nicholson, C.

Variable Temperature NMR Experiment Studying Restricted Bond Rotation. *J. Chem. Educ.* **2020**, *97*, 1425–1429.

(34) Gesmundo, N. J.; Grandjean, J. M.; Nicewicz, D. A. Amide and amine nucleophiles in polar radical crossover cycloadditions: synthesis of gamma-lactams and pyrrolidines. *Org. Lett.* **2015**, *17*, 1316–9.

(35) Wang, N. J.; Mei, S. T.; Shuai, L.; Yuan, Y.; Wei, Y. Aerobic oxidative C–H olefination of cyclic *N*-sulfonyl ketimines catalyzed by a rhodium catalyst. *Org. Lett.* **2014**, *16*, 3040–3043.

(36) (a) Kurahashi, T.; Fujii, H. Unique Ligand-Radical Character of an Activated Cobalt Salen Catalyst That Is Generated by Aerobic Oxidation of a Cobalt(II) Salen Complex. *Inorg. Chem.* **2013**, *52*, 3908–3919. (b) Kemper, S.; Hrobárik, P.; Kaupp, M.; Schlörer, N. E. Jacobsen's Catalyst for Hydrolytic Kinetic Resolution: Structure Elucidation of Paramagnetic Co(III) Salen Complexes in Solution via Combined NMR and Quantum Chemical Studies. *J. Am. Chem. Soc.* **2009**, *131*, 4172–4173. (c) Kochem, A.; Kanso, H.; Baptiste, B.; Arora, H.; Philouze, C.; Jarjays, O.; Vezin, H.; Luneau, D.; Orio, M.; Thomas, F. Ligand Contributions to the Electronic Structures of the Oxidized Cobalt(II) salen Complexes. *Inorg. Chem.* **2012**, *51*, 10557–10571.

(37) Gutmann, V. *The Donor-Acceptor Approach to Molecular Interactions*; Plenum Press: New York, 1978.

(38) A similar trend was observed for the transformation with *t*-BuOOH as the oxidant (see the SI).

(39) A more detailed discussion of feasible mechanistic pathways for the transformation described herein can be found in the SI.

(40) When the reaction was attempted in an intermolecular fashion employing phenylbut-4-ene and an excess of *N*-acetyl sulfonamide, no product was isolated.

Recommended by ACS

Directed, Nickel-Catalyzed Umpolung 1,2-Carboamination of Alkenyl Carbonyl Compounds

Vincent A. van der Puyl, Keary M. Engle, *et al.*

DECEMBER 06, 2018
ACS CATALYSIS

READ 

Non-Stabilized Vinyl Anion Equivalents from Styrenes by *N*-Heterocyclic Carbene Catalysis and Its Use in Catalytic Nucleophilic Aromatic Substitution

Sora Ito, Mamoru Tobisu, *et al.*

APRIL 11, 2022
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Chemoselective Primary Amination of Aryl Boronic Acids by PIII/PVO-Catalysis: Synthetic Capture of the Transient Nef Intermediate HNO

Seung Youn Hong and Alexander T. Radosevich

MAY 12, 2022
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Bioorthogonal Retro-Cope Elimination Reaction of *N,N*-Dialkylhydroxylamines and Strained Alkynes

Dahye Kang and Justin Kim

APRIL 08, 2021
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Get More Suggestions >