# Wastewater microorganisms impact microbial diversity and important ecological functions of stream periphyton

#### **Journal Article**

#### Author(s):

Carles, Louis; Wullschleger, Simon; Joss, Adriano; Eggen, Rik I.L.; Schirmer, Kristin; Schuwirth, Nele; Stamm, Christian (); Tlili, Ahmed

## Publication date: 2022-10-15

Permanent link: https://doi.org/10.3929/ethz-b-000578392

Rights / license: Creative Commons Attribution 4.0 International

Originally published in: Water Research 225, https://doi.org/10.1016/j.watres.2022.119119



Contents lists available at ScienceDirect

#### Water Research



journal homepage: www.elsevier.com/locate/watres

## Wastewater microorganisms impact microbial diversity and important ecological functions of stream periphyton

Louis Carles<sup>a</sup>, Simon Wullschleger<sup>a</sup>, Adriano Joss<sup>a</sup>, Rik I.L. Eggen<sup>a,b</sup>, Kristin Schirmer<sup>a,b,c</sup>, Nele Schuwirth<sup>a,b</sup>, Christian Stamm<sup>a</sup>, Ahmed Tlili<sup>a,b,\*</sup>

<sup>a</sup> Eawag: Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

<sup>b</sup> Department of Environmental Systems Science, ETH, Zürich, Switzerland

<sup>c</sup> School of Architecture, Civil and Environmental Engineering, EPFL Lausanne, Lausanne, Switzerland

#### ARTICLE INFO

#### ABSTRACT

Keywords: Stream biofilm Prokaryotes Eukaryotes DNA metabarcoding Micropollutants Pollution-induced community tolerance Effluents of wastewater treatment plants can impact microbial communities in the receiving streams. However, little is known about the role of microorganisms in wastewater as opposed to other wastewater constituents, such as nutrients and micropollutants. We aimed therefore at determining the impact of wastewater microorganisms on the microbial diversity and function of periphyton, key microbial communities in streams. We used a flow-through channel system to grow periphyton upon exposure to a mixture of stream water and unfiltered or ultra-filtered wastewater. Impacts were assessed on periphyton biomass, activities and tolerance to micro-pollutants, as well as on microbial diversity. Our results showed that wastewater microorganisms colonized periphyton and modified its community composition, resulting for instance in an increased abundance of Chloroflexi and a decreased abundance of diatoms and green algae. This led to shifts towards heterotrophy, as suggested by the changes in nutrient stoichiometry and the increased mineralization potential of carbon sub-strates. An increased tolerance towards micropollutants was only found for periphyton exposed to unfiltered wastewater but not to ultra-filtered wastewater, suggesting that wastewater microorganisms were responsible for this increased tolerance. Overall, our results highlight the need to consider the role of wastewater microorganisms when studying potential impacts of wastewater on the receiving water body.

#### 1. Introduction

Wastewater treatment plants (WWTPs) represent a major source for surface water pollution in urban areas, potentially leading to negative consequences for the structural and functional integrity of aquatic communities in the receiving streams (Gessner and Tlili, 2016; Stamm et al., 2016; Vörösmarty et al., 2010). Effluents from WWTPs typically contain various constituents such as dissolved organic matter, microorganisms and complex mixtures of micropollutants, which are not completely retained by the treatment processes. Due to such chemical and biological complexity, disentangling the specific effects of wastewater constituents on the receiving aquatic ecosystem from each other is a key challenge. This requires a study design that allows for controlled interventions, which cannot be easily achieved in natural environments. We therefore constructed a flow-through channel system that mimics the complexity of field conditions while allowing for targeted manipulations, and used stream periphyton as a biological model to explore the impact of WWTP effluents.

Stream periphyton, also referred to as aquatic biofilms, is a highly diverse and dynamic community of prokaryotic and micro-eukaryotic organisms that are embedded in an extracellular matrix, attached to the surface of submerged solid substrata. Periphyton plays a crucial role in streams as a basis of aquatic food webs and by contributing to important ecological processes such as primary production and nutrient cycling (Battin et al., 2016). With its high biological diversity and important ecological role, periphyton is widely used as a biological community model to assess effects of biotic and abiotic environmental factors, such as trophic interactions, chemical pollution or eutrophication (Montuelle et al., 2010; Sabater et al., 2007). Several studies have reported shifts in the structure of microbial communities in periphyton upon exposure to treated wastewater (Tlili et al., 2020, 2017), with an increase of the relative abundance of cyanobacteria and a decrease of

E-mail address: ahmed.tlili@eawag.ch (A. Tlili).

https://doi.org/10.1016/j.watres.2022.119119

Received 18 May 2022; Received in revised form 20 July 2022; Accepted 13 September 2022 Available online 14 September 2022

0043-1354/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

<sup>\*</sup> Corresponding author at: Department of Environmental Toxicology (Utox), Swiss Federal Institute of Aquatic Science and Technology (Eawag), Dübendorf, Switzerland. Überlandstrasse 133, 8600 Dübendorf, Switzerland.

diatoms (Carles et al., 2021; Romero et al., 2019). These shifts could also be linked to the effects of specific micropollutants from the wastewater effluent (Tamminen et al., 2022; Tardy et al., 2021). Overall, these studies have also shown that these shifts were accompanied by changes in important functional traits of the communities, such as tolerance to micropollutants, algal primary production and photosynthesis or bacterial secondary production. Notwithstanding these findings, most of these studies focused either on the overall effluent toxicity, or on specific wastewater constituents, such as nutrients and micropollutants, but not on the microorganisms from the WWTP.

It has been demonstrated that downstream of the WWTP, bacterial community profiles in the water column (i.e. planktonic) resembled a mixture of the upstream and the effluent communities (Mansfeldt et al., 2020; Price et al., 2018). Periphyton, however, has a different microbial lifestyle than planktonic cells: species interactions play a major role in determining microbial colonization dynamics and in shaping the final periphyton community structure. This means that changes in taxa composition in periphyton cannot be predicted based on hydraulic mixing alone. Systematic studies focusing on the invasion of stream periphyton by microorganisms from WWTPs and the resulting consequences for community structure and function are scarce. For instance, Chonova et al. (2019) have shown that 27% of diatom taxa found in the periphyton downstream of urban and hospital WWTPs potentially originated from the wastewater effluents. In a similar vein, Mußmann et al. (2013) showed that, despite the high diversity of bacterial nitrifiers in the wastewater effluents, only two taxa colonized the downstream periphyton. Nevertheless, monitoring the taxonomic profiles of these microorganisms alone is insufficient, as it does not inform about their contribution to changes in community functions.

Given this background, we aimed in this study to understand the specific contribution of the microorganisms that originate from WWTPs to the composition and functions of periphyton communities. To this end, we used a flow-through channel system where periphyton was grown in the presence of stream water mixed with various fractions of unfiltered or ultra-filtered treated urban wastewater. Ultrafiltration of the wastewater was intended to remove more than 99% of microorganisms while leaving dissolved nutrients and micropollutants unaffected. We measured a large set of structural and functional endpoints targeting the phototrophic and heterotrophic components in periphyton. We also carried out acute toxicity assays to determine the sensitivity of periphyton to a micropollutant mixture extracted from passive samplers deployed in the wastewater effluent. Moreover, we used next generation amplicon sequencing to describe the prokaryotic and eukaryotic community composition in periphyton, and we quantified micropollutants in periphyton and water samples.

#### 2. Material and methods

#### 2.1. Experimental system and design

The experiment was conducted in a flow-through channel system called "Maiandros" that mimics natural conditions in freshwater streams. Initially described by Burdon et al. (2020), this system was subsequently equipped with a light system to ensure a photoperiod of 12 h light: 12 h dark (Carles et al., 2021), as well as with an automated-online monitoring system for parameters such as flow rate, water conductivity and temperature (Desiante et al., 2022). The detailed description of the channel system and the monitoring system can be found in the Supplementary Materials in Desiante et al. (2022). The ultra-filtration (UF) unit added to the system consisted of a membrane module of 50  $\text{m}^2$  surface and a nominal pore size of 0.4  $\mu$ m, allowing for the removal of particulates (including microorganisms) from the effluent, with a microorganism removal efficiency of 99.3% (Desiante et al., 2022). The Maiandros system consists of 20 independent flow-through channels continuously filled with stream water from Chriesbach (47°24'16.7"N 8°36'41.4"E; Dübendorf, Switzerland), a

typical second-order stream of the Swiss Plateau, and wastewater treated for nitrification and denitrification (Desiante et al., 2022), thereafter called "wastewater". The detailed results of the 24 physico-chemical parameters that were monitored in wastewater are available in the Supplementary Materials presented in Desiante et al. (2022). The 20 channels were randomly assigned to five treatments (n = four independent replicate channels per treatment), corresponding to a nominal proportion of 0% (control), 30% and 80% of either unfiltered (WW) or ultra-filtered (UF) wastewater.

#### 2.2. Periphyton colonization and sampling

Periphyton grown for 28 days on glass slides installed in the channels was retrieved and transported to the laboratory for analyses as described by Carles et al. (2021). The duration of colonization was selected based on the assessment of periphyton growth (total biomass and algal biomass) in which we showed that the biofilms reached their maturity after 28 days in our experimental system (Fig. S1). Glass substrata are commonly used in periphyton studies since this inert material allows a better reproducibility and comparison among different conditions and studies. Briefly, five colonized glass slides per channel were scraped and periphyton was suspended in 200 mL of Evian natural water. Evian mineral water is a commonly used growth and exposure medium in microbiological and ecotoxicological studies because of its constant quality. Fresh suspensions were used for the biological analyses (see Section 2.5. below) and the remaining volume was lyophilized prior to micropollutant analysis and Carbon:Nitrogen:Phosphorus (C:N:P) ratio determination.

#### 2.3. Water sampling

Organic micropollutants were sampled in the channels by using passive samplers (AttractSPE®Disks SDB-RPS, 47 mm diameter, Affinisep, France) as described by Carles et al. (2021). Additional passive samplers were also deployed in the wastewater effluent and the accumulated micropollutants were extracted and used for the community tolerance bioassays described in Section 2.5.2 below (Tilil et al., 2017). In order to characterize the microbial community in all water sources by next generation sequencing, three composite water samples were taken weekly from the stream water as well as from the unfiltered and ultra-filtered wastewater during 24 h with an automated water sampler (Maxx, TP5 C Aktiv, Germany). The water samples were automatically taken (50 mL every 30 min), kept at 4 °C, pooled together and immediately filtered on Supor® polyethersulfone membrane disc filters with 0.2  $\mu$ m pore size (Pall Corporation, USA). The filters were stored at -80 °C prior to DNA extraction (see Section 2.5.3 below).

#### 2.4. Micropollutant analyses

Fifty-four substances (i.e. 22 pesticides, 25 pharmaceuticals, 4 artificial sweeteners, 2 corrosion inhibitors and caffeine) that we assumed to be potentially present in the wastewater were extracted and analysed from the passive samplers deployed in the channels according to Moschet et al. (2015) with few modifications (Carles et al., 2021). The selection of the 54 micropollutants was based on a list of priority substances established by Munz et al. (2018) from a large survey in 24 Swiss streams receiving effluents from WWTPs. Extraction of micropollutants used for the tolerance bioassays was carried out according to Tlili et al. (2017). Micropollutant were extracted from periphyton samples as described by Munz et al. (2018) with some modifications (Carles et al., 2021). Micropollutant analysis was performed by HPLC-MS/MS as described by Carles et al. (2021). The limit of quantification (LOQ with matrix factor correction) and relative recovery for each substance in each type of sample are reported in Table S1.

#### 2.5. Biological analyses

#### 2.5.1. Periphyton characterization

Total biomass was determined by measuring the ash free dry weight (AFDW) according to Tlili et al. (2008). Algal biomass was estimated by measuring chlorophyll-a content in periphyton (Sartory and Grobbelaar, 1984). Bacterial biomass was assessed with flow cytometry as described by Frossard et al. (2012) with few modifications (Carles et al., 2021). Total carbon and total nitrogen were determined in lyophilized periphyton samples by using an elemental analyser (HEKAtech Euro Elemental Analyzer; HEKAtech GmbH, Wegberg, Germany). Total phosphorus in freeze-dried periphyton was analysed by inductively coupled plasma mass spectrometry (8900 Triple Quadrupole ICP-MS, Agilent) after an additional digestion step consisting of mixing 5 mg of each periphyton sample with 3 mL of 65% HNO<sub>3</sub> and 1 mL of H<sub>2</sub>O<sub>2</sub> and heating at 250 °C (pressure 120 bars) for 5 min in an ultraCLAVE 4 (Milestone GmbH, Germany).

Photosynthetic efficiency was determined with an Imaging-PAM (pulse amplitude-modulated) fluorimeter (Heinz Walz GmbH, Germany). Primary and secondary productions were measured via the incorporation of <sup>14</sup>C-carbonate and <sup>14</sup>C-leucine according to Dorigo and Leboulanger (2001) and Buesing and Gessner (2003), respectively, with few modifications (Carles et al., 2021). Basal heterotrophic respiration was measured by using the MicroResp<sup>TM</sup> technique as described by Tlili et al. (2011a). The same method was also used to determine the community-level physiological profile (CLPP) of the periphyton suspensions from the various treatments. This was done by measuring the mineralization potential of 14 different carbon sources corresponding to 3 amino acids (glycine, L-cysteine and L-serine), 8 carbohydrates (D (+)-glucose, D(+)-xylose, L-arabinose, D(-)-fructose, D(+)-galactose, D (+)-mannose, D-sorbitol and sucrose), 2 carboxylic acids (citric acid and D(+)-galacturonic acid, monohydrate) and 1 organosulfonic acid (MOPS).

#### 2.5.2. Community tolerance bioassays

Periphyton tolerance to micropollutants was determined via shortterm assays as described in Carles et al. (2021) with few modifications. Briefly, a dilution series of the pure micropollutant extract was prepared in Evian mineral water (dilution factor of 3.16). The optical density of the biofilm suspensions was adjusted at 685 nm to 0.4. Then, 4.5 mL from each suspension were exposed to 0.5 mL of each of the six dilutions of the extract during four hours. Two additional controls were also prepared: Periphyton suspension exposed to 0.5 mL of mineral water (chemical-free control) or to 0.5 mL of 37% formaldehyde (i.e., formaldehyde control). Subsamples were then taken for photosynthetic efficiency, primary production and secondary production measurements, as described above. The same procedure was applied for substrate-induced respiration measurements by using the  $MicroResp^{TM}$ technique. An aliquot of 0.5 mL of each suspension was exposed in 96-deepwell microplates to 50 µL of each of the six dilutions of the extract for 14 h, in addition to the two controls describe above.

## 2.5.3. Next generation sequencing for prokaryotic and eukaryotic community compositions

Total genomic DNA extraction, library construction and sequencing of the 16S rRNA gene for prokaryotes and the 18S rRNA gene for eukaryotes were carried out as described in our previous work (Carles et al., 2021) with few modifications. Briefly, total genomic DNA was extracted from each biofilm suspension, stream water, and unfiltered or ultra-filtered wastewater (membrane disc filters) samples by using the DNeasy PowerBiofilm Kit (QIAGEN) following the manufacturer's instructions. The library construction consisted in two consecutive PCRs. The first PCR amplified the V3-V4 region of the 16S rRNA gene for prokaryotes and the V4-V5 region of the 18S rRNA gene for eukaryotes (see Table S2 for the detailed sequences of the primers) (Herlemann et al., 2011; Hugerth et al., 2014). Multiplexing indices and Illumina sequencing adapters were then added via a second PCR. The libraries were then normalized, pooled and sequenced (paired end  $2 \times 300$  nt, Illumina MiSeq) following the manufacture's run protocols (Illumina, Inc.). All raw sequences are available at the National Center for Biotechnology Information (NCBI) under the SRA accession ID PRJNA755072.

Sequencing data processing, Amplicon Sequence Variants (ASVs) binning and taxonomic assignment were done according to Carles et al. (2021) with few modifications. Briefly, the reads quality was checked with FastQC v0.11.2 (Andrews, 2010), the reads were end-trimmed by seqtk (https://github.com/lh3/seqtk) and then merged using FLASH v1.2.11 (Magoč and Salzberg, 2011). The primers were trimmed with cutadapt v1.12 (Martin, 2011). Quality filtering and subsequent size and GC selection step was carried out with PRINSEQ-lite v0.20.4 (Schmieder and Edwards, 2011). The reads were processed following the ASV analysis (Callahan et al., 2017). The sample reads were first denoised into ASVs by using UNOISE3 in the USEARCH software v.11.0.667. Final predicted taxonomic assignments were performed with SINTAX in the USEARCH software v.11.0.667 (Edgar, 2016) by using the SILVA v128 (16S rRNA) and the PR2 v4.14.0 (18S rRNA) sequence databases. The total number of reads obtained at each bioinformatics step is reported in Table S3.

#### 2.6. Data analyses

Significant differences among the treatments for the periphyton descriptors (i.e., AFDW, chlorophyll-a content, bacterial biomass, photosynthetic efficiency, primary production, secondary production, SIR, basal respiration, C:N:P molar ratios and taxonomic abundance) were assessed using one-way ANOVA followed by separate post hoc comparisons (Tukey's test,  $\alpha = 0.05$ ). The factor tested (i.e. treatment) consisted of five modalities: 0% WW, 30% WW, 30% UF, 80% WW and 80% UF. Normality and homogeneity of variance were checked with the Kolmogorov-Smirnov's and Levene's tests, respectively. Non-normally distributed data were transformed using logarithmic or Box-Cox functions. Statistical analyses were carried out in R 4.0.4 by using RStudio (version 1.4.1717).

The effective concentrations causing a 50% decrease of the measured activity ( $EC_{50}$ ) in the short-term assays were derived from concentration-activity relationships as described in Carles et al. (2021).

Sequencing data analyses for each periphyton and water sample were performed with the R package phyloseq version 1.34.0 (McMurdie and Holmes, 2013). A total of 499,968 and 72,072 reads was obtained after rarefaction for 16S and 18S rRNA datasets, respectively. Alpha diversity (i.e. Chao1 species richness and Shannon diversity index) was assessed with the R package phyloseq. The analysis of beta diversity was based on Bray-Curtis distances, which use the relative abundance of taxa. Permutational Multivariate Analysis of Variance (PERMANOVA) tests were carried out on the Bray-Curtis distances matrix using the R package vegan for prokaryotic and eukaryotic communities separately. Homogeneity in the dispersion of pro- and eukaryotic datasets was first assessed and the adonis function was then used to see if experimental treatments shared similar centroids. Additional pairwise comparisons were carried out by using the pairwise.adonis2 function (Martinez Arbizu, 2020). Graphical representations were generated by using the R package ggplot2 version 3.3.5.

The relative contributions of *source* communities (i.e. stream water, wastewater and UF wastewater) were determined for *sink* communities (i.e. periphyton communities) according to the mixture of stream water and wastewater in the channels by using the fast expectation-maximization for microbial source tracking (FEAST) package in R (Shenhav et al., 2019) for both prokaryotes and eukaryotes. The analysis was repeated five times (1000 iterations each) to reduce the effect of false predictions, with 12 replicates (sampling times) for each water *source* and four channel replicates for each *sink* periphyton community. The FEAST analysis also reports on the potential proportion in the *sink* 

community attributed to other origins (i.e. unknown source).

In order to identify taxa (ASVs) in periphyton responding positively or negatively to wastewater microorganisms, microbiome differential abundance testing was carried out for prokaryotes and eukaryotes by using the R package *DESeq2* version 1.30.1 (Love et al., 2014). This led to the selection of taxa that were assigned to three different groups for prokaryotes and eukaryotes (Figs. 1 and S2). *Group Positive direct* contains microorganisms from the wastewater that colonized periphyton. *Group Positive indirect* corresponds to taxa originating from the stream that were positively impacted by wastewater microorganisms in periphyton. *Group Negative* contains taxa that were negatively impacted by wastewater microorganisms. The differential abundance was tested using the nonrarefied ASV counts. Wald test and adjusted p-values were used to determine if each calculated log2 fold-change differed significantly from zero. In our study, we considered differentially abundant taxa with a log2 fold-change  $\geq |2|$  and Benjamini-Hochberg adjusted *p*-values < 0.05. The control (0 % WW), 30 and 80 % UF periphyton, as well as the UF wastewater were used as reference. Correlation among the prokaryotic and eukaryotic taxa selected by the microbial differential abundance testing was subsequently assessed based on the relative abundance (variance stabilizing transformation – vst-counts) of each taxon in periphyton from all treatments (N = 20). The correlation matrix was visualized with a heatmap displaying the Pearson (r) correlation coefficient (P < 0.05) with the R package *pheatmap* 1.0.12.

#### 3. Results and discussion

#### 3.1. Micropollutants in water and in periphyton

As we expected, the measured concentration of each quantifiable substance in the water in the channels increased with wastewater proportion, indicating that wastewater constituted the primary source of



Fig. 1. Workflow for the microbial differential abundance testing. The same procedure was applied to prokaryotes and eukaryotes datasets. The differential abundance analysis was done with the R package DESeq2. The detailed workflow is available in Fig. S2.

Water Research 225 (2022) 119119

micropollutants in the channels (Fig. 2A and Table S4). Our results also show that these concentrations were similar between the unfiltered and ultra-filtered wastewater for both 30 and 80 % wastewater proportions (Fig. S3). This finding confirms that the ultra-filtration, which was primarily intended to remove microorganisms from the effluent, did not modify the micropollutant composition and concentrations in the wastewater.

Because it is important to link effects to the actual exposure concentrations, we also quantified the micropollutants that accumulated in periphyton (Fig. 2B and Table S5). We found a positive correlation between micropollutant concentrations in periphyton and in the water (Fig. S4A), indicating that micropollutants accumulated in periphyton. However, accumulation varied depending on the measured substance as illustrated by bioconcentration factors (BCFs), which were derived by dividing the concentration in the periphyton by the concentration in the water.

The highest BCF values were found for the pesticides diuron, DEET and isoproturon, as well as for some pharmaceuticals such as clarithromycin and lidocaine (Fig. 2C and Table S6), a pattern that we also observed in a previous study using the same experimental system and treated wastewater (Carles et al., 2021). Differences in bioaccumulation could be explained by differences in the physicochemical properties of the compounds. This was reflected in the positive correlation that we found for the BCFs and log-transformed octanol/water partition coefficients (LogKow) (Fig. S4B). The comparatively high bioaccumulation of PSII inhibitors, such as the herbicides diuron and isoproturon, might be also explained by the presence of specific molecular binding sites, such as the protein D1 in the photosynthetic apparatus, within phototrophic organisms that dominate periphyton (Allen et al., 1983; Morin et al., 2018; Tlili et al., 2011b). On the contrary, the relatively low BCF values for other substances, such as the artificial sweetener acesulfame, may be explained by their high biotransformation rates within microbial cells in periphyton (Desiante et al., 2022, 2021).

#### 3.2. Periphyton characterization

We found no significant effect of wastewater, whether filtered or not, on most of the traditional periphyton descriptors, namely biomass, photosynthetic efficiency or primary and secondary production (Table 1), as also reported previously (Carles et al., 2021; Lebkuecher et al., 2018; Pereda et al., 2019; Tilil et al., 2017). On the other hand, basal respiration in the periphyton exposed to 80% of unfiltered and



**Fig. 2.** Micropollutant concentration in the channel water and in periphyton. Fifty-four substances were included in the targeted mass spectrometry analysis, including 4 artificial sweeteners, 2 corrosion inhibitors, 22 pesticides, 25 pharmaceuticals and caffeine. **A.** Micropollutant concentration in water samples ( $C_{water}$ ) in ng L<sup>-1</sup>. **B.** Micropollutant concentration in periphyton samples ( $C_{periphyton}$ ) in ng mg<sup>-1</sup> of periphyton ash-free dry weight (AFDW). **C.** Bioconcentration factor (BCF): ratio between the micropollutant concentration in periphyton and the average concentration in water for each substance and treatment. C<sub>periphyton</sub>, C<sub>water</sub> and BCF were reported for each substance and each treatment. The treatments correspond to 0% (control), 30% and 80% of unfiltered (WW) and ultra-filtered (UF) wastewater. Values are mean  $\pm$  SD from 4 channel replicates (C<sub>periphyton</sub>) and 4 passive samplers (C<sub>water</sub>).

#### Table 1

Periphyton descriptors (mean  $\pm$  SD, N = 4 replicate channels per treatment). Lower case letters denote significant differences among treatments (a < b < c, Tukey's test, P < 0.05): 0% (control), 30% and 80% wastewater (WW) and ultrafiltered wastewater (UF). Values in bold correspond to significant differences between each treatment and the control.

	0% WW	30% WW	30% UF	80% WW	80% UF
Biomass					
Ash-free dry weight	0.48 $\pm$	0.54 $\pm$	0.41 $\pm$	0.41 $\pm$	$0.37~\pm$
$(mg \ cm^{-2})$	0.06	0.05 (b)	0.05	0.06	0.03 (b)
	(ab)		(ab)	(ab)	
Chlorophyll-a (mg	14.4 $\pm$	$15.9 \pm$	$16.6 \pm$	$16 \pm 3.5$	$19.6 \pm$
g <sup>-1</sup> AFDW)	1.8 (a)	1.5 (a)	2.7 (a)	(a)	4.5 (a)
Bacterial biomass	$1.5 \pm$	$1.5 \pm$	$2.3 \pm$	$2\pm0.3$	$2.2 \pm$
(µg C g <sup>-1</sup> AFDW)	0.6 (a)	0.3 (a)	0.5 (a)	(a)	0.4 (a)
Nutrient ratio					
Carbon:Nitrogen	$\textbf{9.5}\pm\textbf{1.9}$	6.8 ±	6.3 <u>+</u>	5.9 ±	5.5 ±
molar ratio	(b)	0.4 (a)	0.4 (a)	0.4 (a)	0.4 (a)
Carbone:	113.3 $\pm$	61 ±	57.7 ±	60.5 ±	64.1 ±
Phosphorus	12.9 (b)	17.6 (a)	8.3 (a)	23.3 (a)	10.5 (a)
molar ratio					
Nitrogen:	12.5 $\pm$	$9\pm2.6$	$9.1 \pm$	10.1 $\pm$	11.6 $\pm$
Phosphorus	3.5 (a)	(a)	1.3 (a)	3.2 (a)	1.7 (a)
molar ratio					
Functional endpoints					
Photosynthetic	0.36 $\pm$	0.35 $\pm$	0.38 $\pm$	0.42 $\pm$	$0.29~\pm$
efficiency (Quantum	0.03 (a)	0.11 (a)	0.02 (a)	0.01 (a)	0.16 (a)
yield $\phi$ ')					
Primary production	$201.7~\pm$	$210.7~\pm$	$208.5 \ \pm$	320.8 $\pm$	144.8 $\pm$
( $\mu$ g C g <sup>-1</sup> AFDW	49.8 (a)	80.9 (a)	81.4 (a)	49.5 (a)	125.7
day <sup>-1</sup> )					(a)
Secondary	7.9 ±	$4.2 \pm$	$13.9 \pm$	$3.5 \pm$	$1.5 \pm$
production ( $\mu g C g^{-1}$ AFDW dav <sup>-1</sup> )	6.8 (ab)	5.3 (ab)	6.8 (b)	4.1 (ab)	0.7 (a)
Basal respiration (g	$3.8 \pm$	$3.9 \pm$	4.1 $\pm$	5.3 +	5.2 +
$CO2 g^{-1} AFDW$	0.3 (a)	0.5 (ab)	0.6	0.8 (c)	0.6 (bc)
$day^{-\tilde{1}}$ )			(abc)		

ultra-filtered wastewater significantly increased, indicating a potential shift towards heterotrophy. This conclusion is further supported by the clear effect of wastewater on nutrient stoichiometry. Regardless of the treatment, periphyton exposed to wastewater was characterized by lower C:N and C:P molar ratios than the control, with values closer to those described for bacteria than for algae (Table 1). For instance, heterotrophic bacterial cells were described to have a C:P ratio of about 45 (Goldman et al., 1987) while algal C:P ratio is more around 106 (Redfield et al., 1963). Moreover, bacteria have been shown to possess a higher ability to assimilate inorganic phosphorus than algae (Currie and Kalff, 1984). The shift towards heterotrophy could therefore also be explained by changes in the nutrient composition of the extracellular matrix of periphyton, which could be the result of the nutrient input from the wastewater. This was confirmed by the water analyses in the channels, which showed higher concentrations of ortho-phosphate, nitrate and dissolved organic carbon in both 80% unfiltered and ultra-filtered wastewater treatments (1.6  $\pm$  0.2, 7.3  $\pm$  1.7 and 5.7  $\pm$  0.6 mg  $L^{-1}$  , respectively) than in the control treatment (0.5  $\pm$  0.05, 4.9  $\pm$ 0.7 and 2.5  $\pm$  0.3 mg L<sup>-1</sup>, respectively) (Desiante et al., 2022).

Impacts of wastewater on the heterotrophic component of periphyton were also reflected by the changes in the community-level physiological profiles (CLPPs) that we established based on the capability of heterotrophs to mineralize various carbon sources for respiration (Fig. 3). Indeed, the respiration profiles of periphyton exposed to unfiltered and ultra-filtered wastewater differed from each other and from the control communities along both PCA 1 and PCA 2 axes. PCA 1 is clearly related to the wastewater proportion in the channels and shows a positive correlation between wastewater increase and a higher mineralisation potential of all tested carbon substrates. Such results underline again the fact that wastewater exposure favoured the increase of



**Fig. 3.** Principal component analysis (PCA) of the community-level physiological profiles (CLPP) of heterotrophs in periphyton. The profiles were established based on the capacity of heterotrophs to mineralize various carbon sources for respiration, including three amino acids, eight carbohydrates, two carboxylic acids and one organosulfonic acid, as well as a control without an additional carbon source (i.e. measurement of basal respiration). The treatments correspond to periphyton grown in the presence of 0% (control), 30% and 80% of unfiltered (WW) and ultra-filtered (UF) wastewater. The 95% confidence ellipse was added for each treatment.

heterotrophic activities in periphyton communities, likely due to the organic matter and nutrients in the wastewater effluent (Aristi et al., 2016, 2015; Proia et al., 2016). Interestingly, the dissimilarity along PCA 2 between the communities exposed to the ultra-filtered wastewater and the control was less pronounced than for periphyton exposed to the unfiltered wastewater. This suggests that microorganisms from the wastewater have contributed to some extent to the observed changes in the respiration profile of periphyton, which has never been described so far. We hypothesise here that microorganisms originating from wastewater have been exposed to a larger diversity and amounts of carbon sources than those found in the stream water, and may therefore have developed a higher potential capability of mineralization for a broader spectrum of substrates.

#### 3.3. Tolerance of periphyton to micropollutants

Increased microbial community tolerance in periphyton towards micropollutants from WWTPs has been suggested as an indicator to disentangle the specific effects of micropollutants from those of other stressors (Corcoll et al., 2014; Tlili et al., 2020, 2017). We assessed the tolerance of periphyton to a complex mixture of micropollutants that is representative of the micropollutants found in the wastewater effluent (Fig. S5 and Table S7). Irrespective of the treatment, no effect of wastewater was observed on periphyton tolerance based on primary production and respiration, and inconclusive results were obtained when measuring secondary production due to the absence of inhibition and the high variability among replicates (Table 2 and Fig. S6). In sharp contrast, calculated EC50 values based on photosynthetic PSII yield measurements were significantly higher for periphyton exposed to 30% and 80% unfiltered wastewater than for the controls. This reflects an increased tolerance of the phototrophic communities to the micropollutant mixture, as also shown in our previous study (Carles et al., 2021). A potential explanation could be related to the mode of action of PSII inhibitor herbicides, which bind to the QB-binding niche on the D1 protein of the PSII complex, thus blocking electron transport from QA to QB (Jansen et al., 1993). It has been suggested that upon exposure to such herbicides, tolerant phototrophic species can upregulate the

#### Table 2

Tolerance measurements (EC<sub>50</sub> values, N = 4 replicate channels per treatment) of periphyton from the five experimental treatments: 0% (control), as well as 30% and 80% of unfiltered (WW) or ultra-filtered wastewater (UF). Values in parentheses correspond to the 95% confidence interval. R-values are the ratios between the mean of EC<sub>50</sub> of 30% WW, 30% UF, 80% WW and 80% UF divided by the EC<sub>50</sub> of the control for each endpoint. n.d. means not determined due to the absence of inhibition. Values in bold correspond to significant difference between each treatment and the control.

Endpoint	0% WW EC <sub>50</sub>	R	30% WW EC <sub>50</sub>	R	30% UF EC <sub>50</sub>	R	80% WW EC <sub>50</sub>	R	80% UF EC <sub>50</sub>	R
Photosynthetic efficiency	354 (231–545)	-	<b>800 (541–1186)</b>	<b>2.3</b>	348 (213–571)	0.9	<b>653 (492–869)</b>	<b>1.8</b>	299 (167–536)	0.8
Primary production	133 (119–149)	-	139 (108–179)	1	151 (116–198)	1.1	154 (121–196)	1.2	136 (106–175)	1
Secondary production	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Substrate-induced respiration	54 (32–90)	-	58 (35–97)	1.1	63 (37–106)	1.2	40 (21–76)	0.7	89 (42–191)	1.7

expression of the D1 protein, leading to an increased abundance of the QB-PSII inhibitor complex. A community dominated by such tolerant species can therefore cope with higher exposure concentrations than a community dominated by sensitive species, leading to higher  $EC_{50}$  values (Tlili et al., 2011b). This is in agreement with the higher BCFs we found for the herbicide PSII inhibitors, such as diuron and isoproturon. Nevertheless, as epitomized by the concept of stress-induced community sensitivity (Vinebrooke et al., 2004), an increased community tolerance to micropollutants might lead to an enhanced sensitivity towards other stressors and altered ecological functions with negative consequences for the ecosystem functioning.

Regardless of the underlying mechanisms, an important insight from our study is that following the ultrafiltration of the wastewater, which removed more than 99% of the microorganisms, no increased tolerance was observed in periphyton (Table 2 and Fig. S6). Similarly, Desiante et al. (2022) have reported on a loss of micropollutant biotransformation potential by periphyton, one of the mechanisms potentially explaining tolerance, following the ultrafiltration of the wastewater. Collectively, such findings point for the first time towards a major role that microorganisms originating from the WWTPs play in community tolerance to micropollutants. It is conceivable that these microorganisms might have developed a tolerance to micropollutants in the WWTP before being released into the streams. Understanding whether their contribution occurs directly via the colonization of periphyton by micropollutant-tolerant taxa, or indirectly by modifying species interactions within the community, requires a comprehensive characterization of the microbial diversity not only in periphyton but as well in wastewater and stream water.

## 3.4. Microbial diversity and taxonomic abundance in periphyton and water

### *3.4.1.* Relative contribution of stream water and wastewater communities to the periphyton community

Little is known about the relative contribution of wastewater- and stream-microorganisms (i.e. source communities) to periphyton microbial composition (i.e. sink community). To address this knowledge gap, we used the microbial source tracking tool FEAST (Shenhav et al., 2019) and analysed the NGS data for prokaryotes and eukaryotes. Our results clearly indicated that wastewater contributed largely to periphyton communities, with a higher proportion for prokaryotes than for eukaryotes (Fig. 4). For instance, the relative proportion of wastewater in periphyton exposed to 80% wastewater reached up to 79% and 38% for prokaryotes and eukaryotes, respectively. This is in line with the shift towards heterotrophy that we observed in periphyton exposed to wastewater, since prokaryotes in periphyton correspond mainly to heterotrophic bacteria. Another line of evidence comes from the comparison of periphyton exposed to unfiltered and ultra-filtered wastewater: we observed a strong reduction of the relative proportion of wastewater community for periphyton exposed to ultra-filtered wastewater (Fig. 4). This led, for instance for eukaryotes, to a proportion of wastewater community  $\leq$  5% in periphyton exposed to 30% and 80% ultra-filtered



**Fig. 4.** Proportion of water sources in periphyton samples using fast expectation-maximization for microbial source tracking (FEAST) for prokaryotes and eukaryotes. Periphyton was grown in the presence of 0% (control), 30% and 80% of unfiltered (WW) and ultra-filtered (UF) wastewater. The relative contribution of each *source* community (i.e. stream water, unfiltered wastewater and ultra-filtered wastewater) was determined for each *sink* community (i.e. periphyton community) according to the mixture of stream water and wastewater in the channels. FEAST also reports on the potential proportion of the sink attributed to other origins (i.e. *unknown source*). The analysis was repeated five times (1000 iterations each) with 12 replicates (sampling times) for each water source and four channel replicates for each periphyton community. Data are mean of each source proportion (five independent repetitions).

wastewater, confirming in turn the capacity of wastewater microorganisms to colonize periphyton.

The proportion of each sink community that did not match the signature of the sources included in our analysis was assigned to unknown sources. Such analysis is used to identify potential contamination of the sinks by other unidentified microbial sources (Liang et al., 2021; Shenhav et al., 2019). In our study, ultra-filtration did not only lead to an increase of the relative proportion of stream communities in periphyton, but also to an increased proportion of these unknown sources. The relatively high proportion of the unknown sources may be explained by the presence of microorganisms in the Maiandros channel system, and more specifically for the UF treatment, by the colonization of the backside of the membranes by microorganisms forming distinct communities. The colonization dynamics per se, as well as species interactions, in periphyton communities could also lead to different community assemblages compared to the surrounding water column (Peng et al., 2018), and thus potentially contribute to the relatively high proportion of the *unknown* sources. This makes periphyton different from free-living microorganisms in the water column in streams, for which several field surveys have shown that downstream bacterial community profiles were a mixture between the upstream and the effluent (Mansfeldt et al., 2020; Pascual-Benito et al., 2020; Price et al., 2018).

## 3.4.2. Impact of wastewater and wastewater microorganisms on periphyton community structure

We further evaluated the impacts of wastewater on periphyton community structure based on the commonly used descriptors alpha- (i. e. taxonomic richness and Shannon diversity index of a given community) and beta-diversity (i.e. structural differences among several microbial communities). While alpha-diversity for eukaryotes was not affected by wastewater, both taxonomic richness (Chao1) and Shannon diversity of prokaryotes decreased in periphyton exposed to 80% wastewater compared to the control, with no significant difference between unfiltered and ultra-filtered wastewater treatments (Fig. 5A). In contrast, wastewater led to a clear separation in the beta-diversity of both prokaryotic and eukaryotic communities among all treatments (PERMANOVA, P = 0.001, Fig. 5B). Indeed, periphyton communities exposed to unfiltered and ultra-filtered wastewater were distinct from each other and from the control communities (pairwise PERMANOVA, P < 0.05). Several field and mesocosm studies have also reported on the effects of wastewater effluents on the structure of periphyton communities (Carles et al., 2021; Chonova et al., 2019; Lebkuecher et al., 2018; Romero et al., 2019; Tardy et al., 2021), with contrasting results between alpha- and beta-diversity (Carles et al., 2021; Lebkuecher et al., 2018). This may be explained by the various wastewater constituents, such as nutrients, micropollutants and microorganisms. While micropollutants can negatively affect the abundance of certain taxa, nutrients can favor the growth of others (Aristi et al., 2015). Collectively, this may result in distinct communities with regard to beta-diversity but with similar alpha-diversity indices (Hugerth and Andersson, 2017).

## 3.4.3. Identification of periphyton taxa impacted by wastewater microorganisms

By comparing periphyton communities exposed to the unfiltered and ultra-filtered wastewater, as well as with the control communities, we were able to examine how and which microorganisms originating from the wastewater potentially affected the final composition of the communities. For this, we applied the microbial differential abundance testing to identify prokaryotic and eukaryotic taxa in periphyton that are positively or negatively impacted by wastewater microorganisms, in terms of relative abundance. This analysis led to the selection of 129 prokaryotic and 20 eukaryotic taxa (Table S8) that were all correlated to each other (Fig. S7) and subsequently assigned to three groups.

The first group, named *Group Positive direct*, was composed of 98 prokaryotic and 8 eukaryotic taxa that were removed by the ultra-filtration from the wastewater and directly colonized periphyton

exposed to unfiltered wastewater (Fig. 6). All these taxa had a higher abundance in periphyton exposed to unfiltered wastewater than in periphyton exposed to the ultra-filtered wastewater and in the control. Taxa composing this group are therefore potentially major players in periphyton metabolic alterations and increased tolerance that we observed in our study. For instance, two prokaryotic phyla (Chlorobi and Firmicutes) that were only found in this group (Fig. 6A), have been frequently detected in the outlet and downstream-periphyton of WWTPs (Aubertheau et al., 2017; Ross et al., 2012; Thiel et al., 2019; Ziganshina et al., 2016) as well as in biofilters used to treat urban wastewater (Aguirre-Sierra et al., 2016). Several taxa that belong to the prokaryotic phylum Chloroflexi were also assigned to this group. Chloroflexi is a phylum of filamentous bacteria possessing a wide diversity of metabolisms and are also known as photoheterotrophs (Overmann, 2008). This phylum was found to be highly abundant in unfiltered wastewater and periphyton exposed to this wastewater, while almost no Chloroflexi were detected in ultra-filtered wastewater and, accordingly, in periphyton exposed to ultra-filtered wastewater (Fig. S8A). This suggests that these phototrophic bacteria could have contributed directly or indirectly to the observed increased tolerance of phototrophs in periphyton to micropollutants.

The second group, named Group Positive indirect, contained 13 prokaryotic and 8 eukaryotic taxa that did not originate from the WWTP but were positively impacted, in terms of relative abundance in periphyton, by the microorganisms from the wastewater (Fig. 6). Among the prokaryotic taxa, two families (Mycobacteriaceae and Rhodospirillaceae) were specific to this group. Rhodospirillaceae have been shown to be more abundant in lower and medium order streams (Chen et al., 2018), as it is the case for the stream Chriesbach used in our study. It has also been shown that Mycobacteriaceae can thrive in environments that are influenced by anthropogenic activities, such as WWTPs (Amha et al., 2017; Falkinham, 2015; Makovcova et al., 2014). Group Positive indirect also contained members of the phylum Chloroflexi, mainly the family Caldilineaceae that was also found in the Group Positive direct (Fig. 6A). Caldilineaceae may have contributed to the changes in the respiration profiles of periphyton since it has been shown that this family has different metabolic potentials for substrate utilization compared to other microorganisms involved in the enhanced biological phosphorus removal process of WWTPs (Kindaichi et al., 2013).

Microorganisms from the wastewater also had negative impacts on the relative abundance of several taxa in periphyton, which were assigned to the third group, named Group Negative. This group was composed by 18 prokaryotic and 4 eukaryotic taxa, among which several were specific to Group Negative (Fig. 6). For instance, this was the case for several prokaryotic taxa belonging to Cyanobacteria, Deinococcaceae, Methylobacteriaceae and Microbacteriaceae (Fig. 6A). Cyanobacteria are known to respond differently (i.e. an increase or a decrease of abundance) to wastewater effluents (Carles et al., 2021; Carles and Artigas, 2020; Corcoll et al., 2014; Mansfeldt et al., 2020; Romero et al., 2019). However, we are not aware of any study that investigated the specific effect of wastewater microorganisms on this phylum. Deinococcaceae have already been associated to periphyton growing in reference (unpolluted) sites in a previous study comparing community composition along an urban pollution gradient in a stream (Pineda-Mora et al., 2020), indicating their potential sensitivity to wastewater constituents. The presence of Methylobacteriaceae and Microbacteriaceae in the Group Negative may be explained by their association with microalgae (Levy et al., 2009; Paddock et al., 2020), which seemed to be negatively impacted by wastewater microorganisms in terms of abundance (i.e. Class Chlorophyceae in Fig. S7B). Finally, Group Negative also contained one eukaryotic taxon affiliated to Bacillariophyta (i.e. diatoms, Fig. 6B), which is consistent with the observed negative impact of wastewater microorganisms on diatom abundance (Fig. S8B).

These results highlight the need to consider wastewater microbial communities as a stressor *per se*, which can influence the final



**Fig. 5.** Diversity of prokaryotes and eukaryotes in water and in periphyton. The analysis is based on the next generation sequencing of the 16S rRNA (prokaryotes) and 18S rRNA (eukaryotes) genes from water samples (stream water, unfiltered wastewater (WW) and ultra-filtered-wastewater (UF)) and from periphyton grown in the presence of 0% (control), 30% and 80% unfiltered (WW) and ultra-filtered (UF) wastewater. **A:** Alpha diversity; the values of taxonomic richness Chao1 and Shannon's diversity index H' are reported as a boxplot for water (N = 12 sampling times) and periphyton samples (N = 4 channel replicates). Significant differences are indicated by lowercase letters, a > b > c > d > e (Tukey's test, P < 0.05). **B:** Beta diversity; Non-metric Multi-Dimensional Scaling (NMDS) of prokaryotic and eukaryotic communities based on Bray-Curtis distances. The 95% confidence ellipse was added for each sample type.



В



Fig. 6. Repartition of the prokaryotic (A) and eukaryotic (B) taxa selected by the microbial differential abundance testing using DESeq2. The total number of taxa is indicated for each group. The repartition of taxa in each group is described at the phylum, class and family levels for prokaryotes, and at the phylum and class levels for eukaryotes.

composition of periphyton communities, when examining the potential impacts of wastewater on stream ecosystems for water quality assessment. This aspect was poorly described so far, as most studies looked either at the overall effluent toxicity (Liao et al., 2019; Nega et al., 2019; Peng et al., 2018; Romero et al., 2019), or at other wastewater constituents, such as nutrients (Lebkuecher et al., 2018) and micropollutants (Carles et al., 2021; Chonova et al., 2019; Tamminen et al., 2022; Tardy et al., 2021; Tlili et al., 2020). Our study also points towards the importance of species interactions within periphyton communities, as well as between prokaryotes and eukaryotes, which occur among co-existing species in periphyton (Gubelit and Grossart, 2020). For instance, studies on diatom-bacterial interactions revealed a high species-specific interdependence of the algal host and bacteria, as each diatom species developed a bacterial community that differed in its composition (Grossart et al., 2005; Koedooder et al., 2019; Stock et al., 2019).

#### 4. Conclusions

Overall, our study provides compelling evidence that microorganisms originating from wastewater strongly affected periphyton communities. Specifically, we show that these microorganisms are able to colonize periphyton and modify its community composition, either directly or indirectly via species interactions, contributing to changes in respiration profiles. Being at the basis of the food web in streams, such changes in periphyton communities, downstream of WWTPs, bear potential significant environmental costs for higher trophic levels with probable impacts on the overall flow of energy through the food chain in fresh waters. Furthermore, our results also showed that tolerance of periphyton communities to micropollutants was governed by microorganisms released from the WWTP and not by in-stream exposure to the micropollutants. This finding underlines the fact that the Pollution-Induced Community Tolerance (PICT) concept, which stipulates that increased tolerance is directly caused by the exerted selection pressure of micropollutants, should be reconsidered in the context of WWTPs. Instead, selection pressures that occur in highly contaminated compartments such as in the WWTP itself have to be taken into account. Collectively, our findings also have important implications for WWTP management. Microbial communities that are released in the wastewater should be considered as a potential stressor for the receiving streams, similarly to other stressors, such as nutrients, micropollutants or increased temperature. This in turn implies that the measures currently implemented in WWTPs to reduce the load of released microorganisms are not sufficient to completely reduce the ecological hazard they might represent.

#### CRediT authorship contribution statement

Louis Carles: Conceptualization, Data curation, Formal analysis, Writing – original draft, Writing – review & editing. Simon Wullschleger: Data curation, Formal analysis, Writing – review & editing. Adriano Joss: Funding acquisition, Conceptualization, Data curation, Writing – review & editing. Rik I.L. Eggen: Funding acquisition, Conceptualization, Writing – review & editing. Kristin Schirmer: Funding acquisition, Conceptualization, Writing – review & editing. Nele Schuwirth: Funding acquisition, Conceptualization, Writing – review & editing. Christian Stamm: Funding acquisition, Conceptualization, Writing – review & editing, Project administration. Ahmed Tlili: Funding acquisition, Conceptualization, Writing – original draft, Writing – review & editing.

#### **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Data availability

Data will be made available on request.

#### Acknowledgments

This work was supported by the project EcoImpact 2.0 funded by Eawag. The authors thank Eawag colleagues Marco Kipf and Richard Fankhauser for their help in setting up the artificial channels, Werner Desiante for his help in maintaining the artificial channels and Bettina Wagner for her help in sampling and periphyton characterization. As well, we would like to thank Aria Minder, Silvia Kobel and Jean-Claude Walser from the Genetic Diversity Centre (GDC, ETH Zürich) for their support with genetic diversity analyses. Data on molecular diversity produced and analysed in this paper were generated in collaboration with the GDC, ETH Zurich.

#### Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.watres.2022.119119.

#### References

- Aguirre-Sierra, A., Bacchetti-De Gregoris, T., Berná, A., Salas, J.J., Aragón, C., Esteve-Núñez, A., 2016. Microbial electrochemical systems outperform fixed-bed biofilters in cleaning up urban wastewater. Environ. Sci. Water Res. Technol. 2, 984–993. https://doi.org/10.1039/C6EW00172F.
- Allen, M.M., Turnburke, A.C., Lagace, E.A., Steinback, K.E., 1983. Effects of photosystem II herbicides on the photosynthetic membranes of the cyanobacterium *Aphanocapsa* 6308. Plant Physiol. 71, 388–392. https://doi.org/10.1104/pp.71.2.388.
- Amha, Y.M., Anwar, M.Z., Kumaraswamy, R., Henschel, A., Ahmad, F., 2017. Mycobacteria in municipal wastewater treatment and reuse: microbial diversity for screening the occurrence of clinically and environmentally relevant species in arid regions. Environ. Sci. Technol. 51, 3048–3056. https://doi.org/10.1021/acs. est.6b05580.
- Andrews, S., 2010. FastQC: a quality control tool for high throughput sequence data [WWW Document]. URL http://www.bioinformatics.babraham.ac.uk/projects/f astac.
- Aristi, I., Casellas, M., Elosegi, A., Insa, S., Petrovic, M., Sabater, S., Acuña, V., 2016. Nutrients versus emerging contaminants–Or a dynamic match between subsidy and stress effects on stream biofilms. Environ. Pollut. 212, 208–215. https://doi.org/ 10.1016/j.envpol.2016.01.067.
- Aristi, I., von Schiller, D., Arroita, M., Barceló, D., Ponsatí, L., García-Galán, M.J., Sabater, S., Elosegi, A., Acuña, V., 2015. Mixed effects of effluents from a wastewater treatment plant on river ecosystem metabolism: subsidy or stress? Freshw. Biol. 60, 1398–1410. https://doi.org/10.1111/fwb.12576.
- Aubertheau, E., Stalder, T., Mondamert, L., Ploy, M.-C., Dagot, C., Labanowski, J., 2017. Impact of wastewater treatment plant discharge on the contamination of river biofilms by pharmaceuticals and antibiotic resistance. Sci. Total Environ. 579, 1387–1398. https://doi.org/10.1016/j.scitotenv.2016.11.136.
- Battin, T.J., Besemer, K., Bengtsson, M.M., Romani, A.M., Packmann, A.I., 2016. The ecology and biogeochemistry of stream biofilms. Nat. Rev. Microbiol. 14, 251–263. https://doi.org/10.1038/nrmicro.2016.15.
- Buesing, N., Gessner, M.O., 2003. Incorporation of radiolabeled leucine into protein to estimate bacterial production in plant litter, sediment, epiphytic biofilms, and water samples. Microb. Ecol. 45, 291–301. https://doi.org/10.1007/s00248-002-2036-6.
- Burdon, F.J., Bai, Y., Reyes, M., Tamminen, M., Staudacher, P., Mangold, S., Singer, H., Räsänen, K., Joss, A., Tiegs, S.D., Jokela, J., Eggen, R.I.L., Stamm, C., 2020. Stream microbial communities and ecosystem functioning show complex responses to multiple stressors in wastewater. Glob. Change Biol. https://doi.org/10.1111/ gcb.15302.
- Callahan, B.J., McMurdie, P.J., Holmes, S.P., 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. ISME J. 11, 2639–2643. https://doi.org/10.1038/ismej.2017.119.
- Carles, L., Artigas, J., 2020. Interaction between glyphosate and dissolved phosphorus on bacterial and eukaryotic communities from river biofilms. Sci. Total Environ. 719, 137463 https://doi.org/10.1016/j.scitotenv.2020.137463.
- Carles, L., Wullschleger, S., Joss, A., Eggen, R.I.L., Schirmer, K., Schuwirth, N., Stamm, C., Tlili, A., 2021. Impact of wastewater on the microbial diversity of periphyton and its tolerance to micropollutants in an engineered flow-through channel system. Water Res. 117486 https://doi.org/10.1016/j.watres.2021.117486.
- Chen, W., Wilkes, G., Khan, I.U.H., Pintar, K.D.M., Thomas, J.L., Lévesque, C.A., Chapados, J.T., Topp, E., Lapen, D.R., 2018. Aquatic bacterial communities associated with land use and environmental factors in agricultural landscapes using a metabarcoding approach. Front. Microbiol. 9, 2301. https://doi.org/10.3389/ fmicb.2018.02301.

Chonova, T., Kurmayer, R., Rimet, F., Labanowski, J., Vasselon, V., Keck, F., Illmer, P., Bouchez, A., 2019. Benthic diatom communities in an alpine river impacted by waste water treatment effluents as revealed using DNA metabarcoding. Front. Microbiol. 10 https://doi.org/10.3389/fmicb.2019.00653.

Corcoll, N., Acuña, V., Barceló, D., Casellas, M., Guasch, H., Huerta, B., Petrovic, M., Ponsatí, L., Rodríguez-Mozaz, S., Sabater, S., 2014. Pollution-induced community tolerance to non-steroidal anti-inflammatory drugs (NSAIDs) in fluvial biofilm communities affected by WWTP effluents. Chemosphere 112, 185–193. https://doi. org/10.1016/j.chemosphere.2014.03.128.

Currie, D.J., Kalff, J., 1984. A comparison of the abilities of freshwater algae and bacteria to acquire and retain phosphorus. Limnol. Oceanogr. 29, 298–310. https://doi.org/ 10.4319/lo.1984.29.2.0298.

Desiante, W.L., Carles, L., Wullschleger, S., Joss, A., Stamm, C., Fenner, K., 2022. Wastewater microorganisms impact the micropollutant biotransformation potential of natural stream biofilms. Water Res. 118413 https://doi.org/10.1016/j. watres.2022.118413.

Desiante, W.L., Minas, N.S., Fenner, K., 2021. Micropollutant biotransformation and bioaccumulation in natural stream biofilms. Water Res. 116846 https://doi.org/ 10.1016/j.watres.2021.116846.

Dorigo, U., Leboulanger, C., 2001. A pulse-amplitude modulated fluorescence-based method for assessing the effects of photosystem II herbicides on freshwater periphyton. J. Appl. Phycol. 13, 509–515.

Edgar, R.C., 2016. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing (preprint). Bioinformatics. https://doi.org/10.1101/081257.

Falkinham, J.O., 2015. Environmental sources of nontuberculous mycobacteria. Clin. Chest Med. 36, 35–41. https://doi.org/10.1016/j.ccm.2014.10.003.

Frossard, A., Gerull, L., Mutz, M., Gessner, M.O., 2012. Disconnect of microbial structure and function: enzyme activities and bacterial communities in nascent stream corridors. ISME J. 6, 680–691. https://doi.org/10.1038/ismej.2011.134.

Gessner, M.O., Tlili, A., 2016. Fostering integration of freshwater ecology with ecotoxicology. Freshw. Biol. 61, 1991–2001. https://doi.org/10.1111/fwb.12852.

Goldman, J.C., Caron, D.A., Dennett, M.R., 1987. Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C: N ratio 1. Limnol. Oceanogr. 32, 1239–1252.

Grossart, H.-P., Levold, F., Allgaier, M., Simon, M., Brinkhoff, T., 2005. Marine diatom species harbour distinct bacterial communities: marine diatom species harbour distinct bacterial communities. Environ. Microbiol. 7, 860–873. https://doi.org/ 10.1111/j.1462-2920.2005.00759.x.

Gubelit, Y.I., Grossart, H.-P., 2020. New methods, new concepts: what can be applied to freshwater periphyton? Front. Microbiol. 11, 1275. https://doi.org/10.3389/ fmicb.2020.01275.

- Herlemann, D.P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J., Andersson, A.F., 2011. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. ISME J. 5, 1571–1579. https://doi.org/10.1038/ismej.2011.41.
- Hugerth, L.W., Andersson, A.F., 2017. Analysing microbial community composition through amplicon sequencing: from sampling to hypothesis testing. Front. Microbiol. 8, 1561. https://doi.org/10.3389/fmicb.2017.01561.
- Hugerth, L.W., Muller, E.E.L., Hu, Y.O.O., Lebrun, L.A.M., Roume, H., Lundin, D., Wilmes, P., Andersson, A.F., 2014. Systematic design of 18s rRNA gene primers for determining eukaryotic diversity in microbial consortia. PLoS One 9, e95567. https://doi.org/10.1371/journal.pone.0095567.

Jansen, M.A.K., Mattoo, A.K., Malkin, S., Edelman, M., 1993. Direct demonstration of binding-site competition between photosystem II inhibitors at the QB niche of the D1 protein. Pesticide Biochem. Physiol. 46, 78–83.

Kindaichi, T., Nierychlo, M., Kragelund, C., Nielsen, J.L., Nielsen, P.H., 2013. High and stable substrate specificities of microorganisms in enhanced biological phosphorus removal plants: substrate specificities of microorganisms in EBPR plants. Environ. Microbiol. 15, 1821–1831. https://doi.org/10.1111/1462-2920.12074.

Koedooder, C., Stock, W., Willems, A., Mangelinckx, S., De Troch, M., Vyverman, W., Sabbe, K., 2019. Diatom-bacteria interactions modulate the composition and productivity of benthic diatom biofilms. Front. Microbiol. 10 https://doi.org/ 10.3389/fmicb.2019.01255.

Lebkuecher, J.G., Bojic, S., Breeden, C.A., Childs, S.L., Evans, M.C., Hauskins, B.S., Irick, Z.A., Kraft, J.C., Krausfeldt, J.M., Santoyo, N.I., 2018. Photoautotrophicperiphyton composition in reaches with differing nutrient concentrations in the harpeth river of middle Tennessee. Castanea 83, 288. https://doi.org/10.2179/18-163.

Levy, J.L., Stauber, J.L., Wakelin, S.A., Jolley, D.F., 2009. The effect of bacteria on the sensitivity of microalgae to copper in laboratory bioassays. Chemosphere 74, 1266–1274. https://doi.org/10.1016/j.chemosphere.2008.10.049.

Liang, H., Yu, Z., Wang, B., Ndayisenga, F., Liu, R., Zhang, H., Wu, G., 2021. Synergistic application of molecular markers and community-based microbial source tracking methods for identification of fecal pollution in river water during dry and wet seasons. Front. Microbiol. 12. 660368 https://doi.org/10.3389/fmicb.2021.660368.

seasons. Front. Microbiol. 12, 660368 https://doi.org/10.3389/fmicb.2021.660368. Liao, K., Bai, Y., Huo, Y., Jian, Z., Hu, W., Zhao, C., Qu, J., 2019. Use of convertible flow cells to simulate the impacts of anthropogenic activities on river biofilm bacterial communities. Sci. Total Environ. 653, 148–156. https://doi.org/10.1016/j. scitotenv.2018.10.363.

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15, 550. https://doi.org/ 10.1186/s13059-014-0550-8.

Magoč, T., Salzberg, S.L., 2011. FLASH: fast length adjustment of short reads to improve genome assemblies. Bioinformatics 27, 2957–2963. https://doi.org/10.1093/ bioinformatics/btr507. Makovcova, J., Slany, M., Babak, V., Slana, I., Kralik, P., 2014. The water environment as a source of potentially pathogenic mycobacteria. J. Water Health 12, 254–263. https://doi.org/10.2166/wh.2013.102.

Mansfeldt, C., Deiner, K., Mächler, E., Fenner, K., Eggen, R.I.L., Stamm, C., Schönenberger, U., Walser, J.-C., Altermatt, F., 2020. Microbial community shifts in streams receiving treated wastewater effluent. Sci. Total Environ. 709, 135727 https://doi.org/10.1016/j.scitotenv.2019.135727.

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet. J. 17 (10) https://doi.org/10.14806/ej.17.1.200.

Martinez Arbizu, P., 2020. pairwiseAdonis: Pairwise multilevel comparison using adonis. R package version 0.4.

McMurdie, P.J., Holmes, S., 2013. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. PLoS One 8, e61217. https://doi. org/10.1371/journal.pone.0061217.

Montuelle, B., Dorigo, U., Bérard, A., Volat, B., Bouchez, A., Tlili, A., Gouy, V., Pesce, S., 2010. The periphyton as a multimetric bioindicator for assessing the impact of land use on rivers: an overview of the Ardières-Morcille experimental watershed (France). Hydrobiologia 657, 123–141. https://doi.org/10.1007/s10750-010-0105-2.

Morin, S., Chaumet, B., Mazzella, N., 2018. A time-dose response model to assess diuroninduced photosynthesis inhibition in freshwater biofilms. Front. Environ. Sci. 6 https://doi.org/10.3389/fenvs.2018.00131.

Moschet, C., Vermeirssen, E.L.M., Singer, H., Stamm, C., Hollender, J., 2015. Evaluation of *in-situ* calibration of Chemcatcher passive samplers for 322 micropollutants in agricultural and urban affected rivers. Water Res. 71, 306–317. https://doi.org/ 10.1016/j.watres.2014.12.043.

Munz, N.A., Fu, Q., Stamm, C., Hollender, J., 2018. Internal concentrations in gammarids reveal increased risk of organic micropollutants in wastewater-impacted streams. Environ. Sci. Technol. 52, 10347–10358. https://doi.org/10.1021/acs.est.8b03632.

Mußmann, M., Ribot, M., von Schiller, D., Merbt, S.N., Augspurger, C., Karwautz, C., Winkel, M., Battin, T.J., Martí, E., Daims, H., 2013. Colonization of freshwater biofilms by nitrifying bacteria from activated sludge. FEMS Microbiol. Ecol. 85, 104–115. https://doi.org/10.1111/1574-6941.12103.

Nega, M., Braun, B., Künzel, S., Szewzyk, U., 2019. Evaluating the impact of wastewater effluent on microbial communities in the Panke, an Urban River. Water 11, 888. https://doi.org/10.3390/w11050888.

Overmann, J., 2008. Green nonsulfur bacteria. In: Pettis, G. (Ed.), Encyclopedia of Life Sciences. John Wiley & Sons, Ltd, Chichester, pp. 1–10.

Paddock, M.B., Fernández-Bayo, J.D., VanderGheynst, J.S., 2020. The effect of the microalgae-bacteria microbiome on wastewater treatment and biomass production. Appl. Microbiol. Biotechnol. 104, 893–905. https://doi.org/10.1007/s00253-019-10246-x.

Pascual-Benito, M., Ballesté, E., Monleón-Getino, T., Urmeneta, J., Blanch, A.R., García-Aljaro, C., Lucena, F., 2020. Impact of treated sewage effluent on the bacterial community composition in an intermittent mediterranean stream. Environ. Pollut. 266, 115254 https://doi.org/10.1016/j.envpol.2020.115254.

Peng, Y., Li, J., Lu, J., Xiao, L., Yang, L., 2018. Characteristics of microbial community involved in early biofilms formation under the influence of wastewater treatment plant effluent. J. Environ. Sci. 66, 113–124. https://doi.org/10.1016/j. jes.2017.05.015.

Pereda, O., Acuña, V., von Schiller, D., Sabater, S., Elosegi, A., 2019. Immediate and legacy effects of urban pollution on river ecosystem functioning: a mesocosm experiment. Ecotoxicol. Environ. Saf. 169, 960–970. https://doi.org/10.1016/j. ecoenv.2018.11.103.

Pineda-Mora, D., Juárez-López, A.L., Toribio-Jiménez, J., Leal-Ascencio, M.T., Ruvalcaba-Ledezma, J.C., Castelán-Sánchez, H.G., Aguirre-Noyola, J.L., Arp, P.A., 2020. Diversity and functions of epilithic riverine biofilms. Water Air Soil Pollut. 231, 391. https://doi.org/10.1007/s11270-020-04692-x.

Price, J.R., Ledford, S.H., Ryan, M.O., Toran, L., Sales, C.M., 2018. Wastewater treatment plant effluent introduces recoverable shifts in microbial community composition in receiving streams. Sci. Total Environ. 613 (614), 1104–1116. https://doi.org/ 10.1016/j.scitotenv.2017.09.162.

Proia, L., von Schiller, D., Gutierrez, C., Casas-Ruiz, J.P., Gómez-Gener, L., Marcé, R., Obrador, B., Acuña, V., Sabater, S., 2016. Microbial carbon processing along a river discontinuum. Freshw. Sci. 35, 1133–1147. https://doi.org/10.1086/689181.

Redfield, A.C., Ketchum, B.H., Richards, F.A., 1963. The Influence of Organisms on the Composition of Seawater. The Sea. Hill M.N., New York, pp. 26–77.

Romero, F., Sabater, S., Font, C., Balcázar, J.L., Acuña, V., 2019. Desiccation events change the microbial response to gradients of wastewater effluent pollution. Water Res. 151, 371–380. https://doi.org/10.1016/j.watres.2018.12.028.

Res. 151, 371–380. https://doi.org/10.1016/j.watres.2018.12.028.
Ross, K.A., Feazel, L.M., Robertson, C.E., Fathepure, B.Z., Wright, K.E., Turk-MacLeod, R. M., Chan, M.M., Held, N.L., Spear, J.R., Pace, N.R., 2012. Phototrophic phylotypes dominate mesothermal microbial mats associated with hot springs in yellowstone national park. Microb. Ecol. 64, 162–170. https://doi.org/10.1007/s00248-012-0012-3.

Sabater, S., Guasch, H., Ricart, M., Romaní, A., Vidal, G., Klünder, C., Schmitt-Jansen, M., 2007. Monitoring the effect of chemicals on biological communities. The biofilm as an interface. Analyt. Bioanalyt. Chem. 387, 1425–1434. https://doi.org/ 10.1007/s00216-006-1051-8.

Sartory, D.P., Grobbelaar, J.U., 1984. Extraction of chlorophyll a from freshwater phytoplankton for spectrophotometric analysis. Hydrobiologia 114, 177–187. https://doi.org/10.1007/BF00031869.

Schmieder, R., Edwards, R., 2011. Quality control and preprocessing of metagenomic datasets. Bioinformatics 27, 863–864. https://doi.org/10.1093/bioinformatics/ btr026.

Shenhav, L., Thompson, M., Joseph, T.A., Briscoe, L., Furman, O., Bogumil, D., Mizrahi, I., Pe'er, I., Halperin, E., 2019. FEAST: fast expectation-maximization for

#### L. Carles et al.

microbial source tracking. Nat. Methods 16, 627–632. https://doi.org/10.1038/ s41592-019-0431-x.

- Stamm, C., Räsänen, K., Burdon, F.J., Altermatt, F., Jokela, J., Joss, A., Ackermann, M., Eggen, R.I.L., 2016. Unravelling the impacts of micropollutants in aquatic ecosystems. Adv. Ecol. Res. 55, 183–223. https://doi.org/10.1016/bs. aecr.2016.07.002. Elsevier.
- Stock, W., Blommaert, L., De Troch, M., Mangelinckx, S., Willems, A., Vyverman, W., Sabbe, K., 2019. Host specificity in diatom–bacteria interactions alleviates antagonistic effects. FEMS Microbiol. Ecol. 95 https://doi.org/10.1093/femsec/ fiz171 fiz171.
- Tamminen, M., Spaak, J., Tlili, A., Eggen, R., Stamm, C., Räsänen, K., 2022. Wastewater constituents impact biofilm microbial community in receiving streams. Sci. Total Environ. 807, 151080 https://doi.org/10.1016/j.scitotenv.2021.151080.
- Tardy, V., Bonnineau, C., Bouchez, A., Miège, C., Masson, M., Jeannin, P., Pesce, S., 2021. A pilot experiment to assess the efficiency of pharmaceutical plant wastewater treatment and the decreasing effluent toxicity to periphytic biofilms. J. Hazard. Mater. 411, 125121 https://doi.org/10.1016/j.jhazmat.2021.125121.
- Thiel, V., Garcia Costas, A.M., Fortney, N.W., Martinez, J.N., Tank, M., Roden, E.E., Boyd, E.S., Ward, D.M., Hanada, S., Bryant, D.A., 2019. Candidatus thermonerobacter thiotrophicus," A non-phototrophic member of the bacteroidetes/ chlorobi with dissimilatory sulfur metabolism in hot spring mat communities. Front. Microbiol. 9, 3159. https://doi.org/10.3389/fmicb.2018.03159.
- Tlili, A., Corcoll, N., Arrhenius, Å., Backhaus, T., Hollender, J., Creusot, N., Wagner, B., Behra, R., 2020. Tolerance patterns in stream biofilms link complex chemical pollution to ecological impacts. Environ. Sci. Technol. 54, 10745–10753. https:// doi.org/10.1021/acs.est.0c02975.

- Tlili, A., Dorigo, U., Montuelle, B., Margoum, C., Carluer, N., Gouy, V., Bouchez, A., Berard, A., 2008. Responses of chronically contaminated biofilms to short pulses of diuronAn experimental study simulating flooding events in a small river. Aquat. Toxicol. 87, 252–263. https://doi.org/10.1016/j.aquatox.2008.02.004.
- Tlili, A., Hollender, J., Kienle, C., Behra, R., 2017. Micropollutant-induced tolerance of *in situ* periphyton: establishing causality in wastewater-impacted streams. Water Res. 111, 185–194. https://doi.org/10.1016/j.watres.2017.01.016.
- Tlili, A., Marechal, M., Montuelle, B., Volat, B., Dorigo, U., Bérard, A., 2011a. Use of the MicroResp<sup>TM</sup> method to assess pollution-induced community tolerance to metals for lotic biofilms. Environ. Pollut. 159, 18–24. https://doi.org/10.1016/j. envpol.2010.09.033.
- Tlili, A., Montuelle, B., Bérard, A., Bouchez, A., 2011b. Impact of chronic and acute pesticide exposures on periphyton communities. Sci. Total Environ. 409, 2102–2113. https://doi.org/10.1016/j.scitotenv.2011.01.056.
- Vinebrooke, R.D., Cottingham, K.L., Norberg, J., Dodson, S.I., Maberly, S.C., Sommer, U., 2004. Impacts of multiple stressors on biodiversity and ecosystem functioning: the role of species co-tolerance. Oikos 104, 451–457. https://doi.org/10.1111/j.0030-1299.2004.13255.x.
- Vörösmarty, C.J., McIntyre, P.B., Gessner, M.O., Dudgeon, D., Prusevich, A., Green, P., Glidden, S., Bunn, S.E., Sullivan, C.A., Liermann, C.R., Davies, P.M., 2010. Global threats to human water security and river biodiversity. Nature 467, 555–561. https://doi.org/10.1038/nature09440.
- Ziganshina, E.E., Ibragimov, E.M., Ilinskaya, O.N., Ziganshin, A.M., 2016. Bacterial communities inhabiting toxic industrial wastewater generated during nitrocellulose production. Biologia 71, 70–78. https://doi.org/10.1515/biolog-2016-0014.