

Minimal mirror twin Higgs

Journal Article**Author(s):**

Barbieri, Riccardo; Hall, Lawrence J.; Harigaya, Keisuke

Publication date:

2016-11

Permanent link:

<https://doi.org/10.3929/ethz-b-000123676>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

Journal of High Energy Physics 2016(11), [https://doi.org/10.1007/JHEP11\(2016\)172](https://doi.org/10.1007/JHEP11(2016)172)

Minimal mirror twin Higgs

Riccardo Barbieri,^{a,b} Lawrence J. Hall^{c,d} and Keisuke Harigaya^{c,d}

^a*Institute of Theoretical Studies, ETH Zurich,
CH-8092 Zurich, Switzerland*

^b*Scuola Normale Superiore,
Piazza dei Cavalieri 7, 56126 Pisa, Italy*

^c*Department of Physics, University of California,
Berkeley, California 94720, U.S.A.*

^d*Theoretical Physics Group, Lawrence Berkeley National Laboratory,
Berkeley, California 94720, U.S.A.*

E-mail: riccardo.barbieri@sns.it, ljhall@lbl.gov,
keisukeharigaya@berkeley.edu

ABSTRACT: In a Mirror Twin World with a maximally symmetric Higgs sector the little hierarchy of the Standard Model can be significantly mitigated, perhaps displacing the cutoff scale above the LHC reach. We show that consistency with observations requires that the Z_2 parity exchanging the Standard Model with its mirror be broken in the Yukawa couplings. A minimal such effective field theory, with this sole Z_2 breaking, can generate the Z_2 breaking in the Higgs sector necessary for the Twin Higgs mechanism. The theory has constrained and correlated signals in Higgs decays, direct Dark Matter Detection and Dark Radiation, all within reach of foreseen experiments, over a region of parameter space where the fine-tuning for the electroweak scale is 10-50%. For dark matter, both mirror neutrons and a variety of self-interacting mirror atoms are considered. Neutrino mass signals and the effects of a possible additional Z_2 breaking from the vacuum expectation values of $B - L$ breaking fields are also discussed.

KEYWORDS: Beyond Standard Model, Cosmology of Theories beyond the SM

ARXIV EPRINT: [1609.05589](https://arxiv.org/abs/1609.05589)

Contents

1	Introduction	1
2	Review of the twin Higgs mechanism	3
3	Necessity of Z_2 symmetry breaking in Yukawa couplings	6
4	Minimal mirror twin Higgs	8
4.1	Constraint on Yukawa couplings from Δm_H^2	9
4.2	Standard Model like Higgs decays	10
4.3	Mirror QCD phase transition temperature	10
5	Thermal history with Higgs exchange	12
5.1	QCD' and constraints on the q' spectrum	12
5.2	Decoupling temperature	13
5.3	Dark radiation	16
5.4	Cosmological signals of mirror neutrino masses	17
6	Thermal history including kinetic mixing	17
6.1	Decoupling temperature	18
6.2	Dark radiation	19
6.3	Milli-charged particle	20
7	Mirror baryon dark matter	20
7.1	Dark matter candidates	20
7.2	Direct detection of dark matter	23
8	Non-minimal Z_2 breaking from $B - L$ vevs	25
8.1	Dark radiation	25
8.2	Additional dark matter candidates	25
9	Conclusions	27
A	Heavy axion and Z_2 symmetry breaking in Froggatt Nielsen sector	29

1 Introduction

An intriguing idea, that has persisted over several decades, is that of the Mirror World: the Standard Model, with quarks and leptons (q, l) and gauge interactions $SU(3) \times SU(2) \times U(1)$, is supplemented by an identical sector where mirror quarks and leptons (q', l') interact via

mirror gauge interactions $SU(3)' \times SU(2)' \times U(1)'$. There are two prime motivations for this idea. The discrete symmetry that interchanges the ordinary and mirror worlds can be interpreted as spacetime parity, P , allowing a neat restoration of parity [1, 2]. Secondly, mirror baryons are expected to be produced in the early universe and to be sufficiently stable to yield dark matter, and this may lead to an understanding of why the cosmological energy densities of baryons and dark matter are comparable.

A third, more recent motivation for the Mirror World arises from the absence so far of new physics at colliders to explain the origin of the weak scale. If the Higgs doublets of the two sectors (H, H') possess a potential with maximal $SO(8)$ symmetry at leading order, the observed Higgs boson becomes a pseudo-Nambu-Goldstone boson with a mass that is insensitive to the usual Standard Model (SM) quadratic divergences; this is the Twin Higgs idea [3]. Furthermore, if the symmetric quartic coupling of this potential, λ , is large relative to the SM quartic coupling, λ_{SM} , the Mirror World reduces the amount of fine-tuning by a factor of $2\lambda/\lambda_{SM}$ to reach any particular UV cutoff of the effective theory. Since we now know that $\lambda_{SM} = 0.13$ is small, this improvement can be very significant, allowing a Little Hierarchy between the weak scale and the UV cutoff, which may be beyond the LHC reach.

In this paper we formulate a minimal, experimentally viable, low energy effective theory for this idea, Minimal Mirror Twin Higgs, and study its signals. This is a pressing issue: mirror baryon dark matter, the Twin Higgs mechanism and consistency with cosmological limits on the amount of dark radiation all require a breaking of parity, P [4]. How is this to be accomplished? We do not attempt a UV completion, whether supersymmetric [5, 6] or with composite Higgs [7–10], but note that both account for the approximate $SO(8)$ symmetry of the Higgs potential.

The $SO(8)$ invariant quartic interaction contains an interaction $H^\dagger H H'^\dagger H'$ thermally coupling the two sectors at cosmological temperatures above a few GeV, so that the bound on dark radiation provides a very severe constraint on Mirror Twin Higgs. The Twin Higgs mechanism requires a parity breaking contribution to the Higgs mass terms in the potential, Δm_H^2 . We find this term by itself to be insufficient to solve the dark radiation problem, nomatter what other interactions connect the two sectors, at least for fine-tunings above the percent level. This then implies that the Yukawa couplings of the two sectors differ, $y' \neq y$.

Hence we introduce an effective theory, Minimal Mirror Twin Higgs, where all P violation arises from a breaking of flavor symmetry, yielding different Yukawa couplings in the two sectors. This single source of P violation leads simultaneously to three key results

- The Δm_H^2 term necessary for the Twin Higgs mechanism is generated via q' loops.
- The mirror QCD phase transition temperature is raised above the decoupling temperature of the two sectors, solving the problem of excessive dark radiation.
- The mirror baryon mass is raised, allowing viable dark matter.

A striking signature would be the discovery at the LHC, or a future collider, of the mirror Higgs itself, decaying to WW or ZZ ; see [4] and figure 10 of [11]. As the mirror Higgs mass depends on the $SO(8)$ invariant quartic, λ , it could be beyond the LHC range, and here we focus on other signals. The size of P breaking in the Yukawa couplings to obtain

the above three results leads to a preferred range of the lightest q' mass of $(2 - 20)$ GeV, leading us to compute signals for the following quantities

- The signal strength, μ , and the invisible width, $\Gamma(h \rightarrow \text{inv})$, of the Higgs boson.
- The amount of dark radiation, ΔN_{eff} .
- The direct detection rate for mirror baryon dark matter from Higgs exchange.
- The effective sum of neutrino masses affecting large scale structure and the CMB.

These signals are tightly correlated as they all depend on the Higgs portal between the two sectors, the ratio of the weak scales, and on the masses of the light q' . For dark matter, both mirror neutrons and a variety of self-interacting mirror atoms are considered.

After a brief review of the Twin Higgs mechanism in section 2, we demonstrate that the breaking of P in the Yukawa couplings is necessary in section 3. We define the Minimal Mirror Twin Higgs theory in section 4, and discuss the consequences of breaking P in the Yukawa couplings. We constrain the q' Yukawa couplings from $\Gamma(h \rightarrow \text{inv})$ and Δm_H^2 , and study how large the mirror QCD phase transition temperature T'_c can be. In section 5 we study the cosmological history of the two sectors when the only communication between them arises from the Higgs interaction $H^\dagger H H'^\dagger H'$ and find that the decoupling temperature can be lower than T'_c , allowing ΔN_{eff} to lie inside the observational limit. We predict the amount of dark radiation and the effective sum of neutrino masses. In section 6 we examine an alternative cosmology when communication between the sectors is dominated by kinetic mixing of the hypercharge gauge bosons. We study a variety of candidates for mirror dark matter in section 7, and find that the $H^\dagger H H'^\dagger H'$ interaction, together with the enhanced q' Yukawa couplings, will allow direct detection at planned experiments over a large part of the mass range. In section 8 we briefly study ΔN_{eff} and dark matter candidates when additional P breaking arises from the absence of Majorana masses for right-handed mirror neutrinos. In the appendix we show that a PQ symmetry common to both sectors allows a solution to the strong CP problem, with the axion mass enhanced by the mirror sector by a factor of order 10^3 , leading to the possibility that f_a is of order 10 TeV.

2 Review of the twin Higgs mechanism

In this section, we review the Twin Higgs mechanism using a linear sigma model. The Standard Model Higgs doublet H , together with a scalar field H' , is embedded into a representation of some approximate global symmetry. The global symmetry is broken down to a subgroup containing $SU(2)_L \times U(1)_Y$ by the vacuum expectation value (VEV) of H' . Four of the pseudo-Nambu-Goldstone bosons form a doublet of $SU(2)_L$ and are identified with the Standard Model Higgs doublet. The lightness of the Higgs mass in comparison with the scale of the Higgs potential can be understood in this way.

Let us take a closer look at the Higgs potential. A global symmetry preserving potential is given by

$$V_{\text{sym}} = \lambda (|H|^2 + |H'|^2)^2 + m_H^2 (|H|^2 + |H'|^2). \quad (2.1)$$

We have neglected possible higher order terms which are expected in composite Twin Higgs models, as they are irrelevant for the following discussion. Since the global symmetry is explicitly broken by Yukawa couplings and the electroweak gauge interaction, we expect a breaking of the global symmetry in the Higgs potential, at least by quantum corrections. The quantum correction to the mass term is the most dangerous, and to suppress it we assume a Z_2 symmetry $H \leftrightarrow H'$ and call H' the mirror Higgs. We also introduce appropriate mirrors of other SM particles. In the following, we use “ ’ ” to denote mirror objects. The global symmetry of the Higgs potential is now $SO(8)$. A Z_2 symmetric mass term is accidentally $SO(8)$ symmetric, and an $SO(8)$ breaking potential is given by¹

$$V_{SO(8)} = \Delta\lambda (|H|^4 + |H'|^4). \tag{2.2}$$

As we will see later, we need small Z_2 breaking terms to obtain a correct electroweak symmetry breaking scale. We assume Z_2 breaking in the mass term,

$$V_{Z_2} = \Delta m_H^2 |H|^2. \tag{2.3}$$

The origin of the Z_2 breaking mass term is explained in the next section.

Let us derive the VEVs of the Higgs fields in the small $SO(8)$ breaking limit. Assuming $\Delta m_H^2 > 0$, we expect $\langle H' \rangle^2 > \langle H \rangle^2$. Setting $\langle H \rangle = 0$ initially, the VEV of $\langle H' \rangle$ is given by

$$v'^2 \equiv \langle H' \rangle^2 = \frac{-m_H^2}{2\lambda}. \tag{2.4}$$

In the unitary gauge of $SU(2)_L \times U(1)_Y$, the pseudo-Nambu Goldstone boson, namely the Standard Model like Higgs h , is given by

$$\begin{pmatrix} H \\ H' \end{pmatrix} \rightarrow v' \begin{pmatrix} \sin \frac{h}{\sqrt{2}v'} \\ \cos \frac{h}{\sqrt{2}v'} \end{pmatrix}. \tag{2.5}$$

Minimizing the potential of h given by eqs. (2.2) and (2.3), we obtain

$$v^2 \equiv \langle H \rangle^2 = v'^2 \sin^2 \frac{\langle h \rangle}{\sqrt{2}v'} \simeq \frac{v'^2}{2} \left(1 - \frac{\Delta m_H^2}{2\Delta\lambda v'^2} \right). \tag{2.6}$$

The mass of h is given by

$$m_h^2 \simeq 8\Delta\lambda v^2, \tag{2.7}$$

whereas the mass of the mirror Higgs boson, h' , is

$$m_{h'}^2 \simeq 4\lambda v'^2. \tag{2.8}$$

From eqs. (2.6) and (2.7), the required Z_2 breaking is given by

$$\Delta m_H^2 \simeq \frac{1}{4} \frac{v'^2}{v^2} m_h^2 \simeq (200 \text{ GeV})^2 \times \left(\frac{v'/v}{3} \right)^2. \tag{2.9}$$

¹Quantum corrections from top quarks gives $\Delta\lambda = 3y_t^4/64\pi^2(\log(\Lambda^2/m_t^2) + A)$, with Λ the cut off scale of the Higgs sector and A a finite UV-dependent term. To obtain the observed Higgs mass requires $\log(\Lambda/m_t) + A/2 \approx 7$. Alternatively a tree level $\Delta\lambda$ may exist, as, e.g., from a supersymmetric D-term. See [9] for the case of a composite Twin Higgs model.

Let us comment on the fine-tuning in the Twin Higgs model. In order to obtain the hierarchy between v' and v , Δm_H^2 must be tuned against $\Delta \lambda v'^2$. The standard fine-tuning measure is given by

$$\Delta \equiv \left| \frac{\partial \ln v^2}{\partial \ln \Delta m_H^2} \right| = \frac{1}{2} \frac{v'^2}{v^2}. \quad (2.10)$$

The mixing in the physical higgs bosons, h and h' , imply an overall reduction of the couplings of the Standard Model higgs to any Standard Model particles by the relative amount $(1 - 1/2(v/v')^2)$. The precision measurements of these couplings [13] require that $v'/v > 2$ at 95% C.L. (see section 4.2).² Hence we need a tuning of at least 50%: this is the unavoidable, minimal fine-tuning in Twin Higgs models.

In general we expect a fine tuning also in the mass of the mirror Higgs boson. Assuming that the dominant contribution to this fine tuning comes from the top loop suitably cutoff at a scale Λ_{TH} ,³ it is

$$\Delta_{m_{h'}} \approx \frac{3/8\pi^2 y_t^2 \Lambda_{\text{TH}}^2}{2\lambda v'^2}. \quad (2.11)$$

If $\Delta_{m_{h'}} > 1$, the overall fine tuning in Twin Higgs is given by

$$\Delta_{\text{TH}} = \frac{1}{2} \frac{v'^2}{v^2} \times \Delta_{m_{h'}}, \quad (2.12)$$

that can be compared with the fine tuning in the SM

$$\Delta_{\text{SM}} = \frac{3/8\pi^2 y_t^2 \Lambda_{\text{SM}}^2}{2\lambda_{\text{SM}} v^2}, \quad (2.13)$$

where λ_{SM} is the SM quartic coupling and Λ_{SM} is the cut off scale of the SM.

Thus the fine tuning in TH relative to the SM is

$$\Delta_{\text{SM}}/\Delta_{\text{TH}} = \frac{2\lambda}{\lambda_{\text{SM}}} \frac{\Lambda_{\text{SM}}^2}{\Lambda_{\text{TH}}^2}. \quad (2.14)$$

As $\lambda_{\text{SM}} \simeq 0.13$, this improvement is significant: for a moderate tuning Λ_{TH} can be above the scales directly explorable at the LHC.

Let us comment on the required quality of the Z_2 symmetry [9, 12]. The top Yukawa coupling y_t gives a one-loop quantum correction,

$$\Delta m_H^2|_{y_t} \simeq \frac{3}{8\pi^2} (y_{t'}^2 - y_t^2) \Lambda^2, \quad (2.15)$$

where Λ is the cut off of the Higgs sector. In composite Twin Higgs models, Λ is the scale of higher resonances, which is expected be as large as $\Lambda \sim g_* v'$, where g_* is the coupling strength of hadrons. The naive-dimensional analysis [14, 15] and the large N counting [16] suggest $g_* \sim 4\pi/\sqrt{N}$, where N is the size of the confining gauge group of the composite

²The electroweak precision measurement as well sets an indirect bound on v'/v , which depends on the mass of the mirror Higgs. For $m_{h'} = 1$ TeV, considering the IR contribution only, one obtains $v'/v > 3$ at 90% C.L.

³For $\lambda > 1$ this requires a suppression of the Higgs loop contribution in the UV completion of the Twin Higgs model considered here.

Twin Higgs model. In the supersymmetric Twin Higgs model, Λ is the stop mass scale multiplied by a log enhancement factor. Requiring that this correction does not exceed the required one in eq. (2.9), we obtain

$$\left| \frac{y_{t'} - y_t}{y_t} \right| \lesssim 0.03 \left(\frac{v'/v}{3} \right)^2 \left(\frac{5 \text{ TeV}}{\Lambda} \right)^2. \quad (2.16)$$

The strong interaction gives a two loop quantum correction,

$$\Delta m_H^2|_{\text{strong}} \simeq \frac{3y_t^2 (g_3^2 - g_3'^2)}{8\pi^4} \Lambda^2, \quad (2.17)$$

leading to the requirement

$$\left| \frac{g_3'^2 - g_3^2}{g_3^2} \right| \lesssim 0.5 \left(\frac{v'/v}{3} \right)^2 \left(\frac{5 \text{ TeV}}{\Lambda} \right)^2. \quad (2.18)$$

Finally, the $SU(2)_L$ interaction gives a one-loop quantum correction,

$$\Delta m_H^2|_{\text{weak}} \simeq \frac{9 (g_2^2 - g_2'^2)}{64\pi^2} \Lambda^2, \quad (2.19)$$

requiring

$$\left| \frac{g_2'^2 - g_2^2}{g_2^2} \right| \lesssim 0.2 \left(\frac{v'/v}{3} \right)^2 \left(\frac{5 \text{ TeV}}{\Lambda} \right)^2. \quad (2.20)$$

A natural explanation for the quality of the Z_2 symmetry shown in eqs. (2.16), (2.18) and (2.20) is that the Lagrangian is precisely Z_2 symmetric, with a complete copy of all Standard Model particles — this is nothing but the Mirror World. A key question then becomes the form and origin of the Z_2 breaking necessary to construct a fully realistic theory.

3 Necessity of Z_2 symmetry breaking in Yukawa couplings

As reviewed in the previous section, the Twin Higgs mechanism requires a Z_2 symmetry under which the Standard Model and mirror particles are exchanged. The Z_2 symmetry must be broken eventually to obtain the correct electroweak symmetry breaking scale. In this section, we show that it is mandatory to break the Z_2 symmetry in Yukawa couplings to suppress the abundance of dark radiation, no matter what the origin of Z_2 breaking is and no matter what interactions might be added to the theory to couple the two sectors. Thus the minimal phenomenologically viable way to break the Z_2 symmetry is via $y_{f'} \neq y_f$.

The Standard Model and mirror particles interact with each other by the interaction in eq. (2.1) so that in the early universe at temperatures of order the weak scale the two sectors are kept in equilibrium via Higgs exchange. At lower temperatures the mirror particles eventually annihilate/decay into mirror photons (and mirror neutrinos), giving an extra component of relativistic particles, which is often referred to as dark radiation. The amount of dark radiation depends on the decoupling temperature between the two sectors, T_d , below which the interaction rate between Standard Model and mirror particles is smaller than the Hubble expansion rate. Without introducing extra interactions, T_d is determined by Higgs




Figure 1. Prediction for the amount of extra dark radiation arising from only e' , μ' , τ' and γ' as a function of the decoupling temperature T_d between the Standard Model and mirror sectors.

exchange which depends on the Yukawa couplings [4, 5]. In this section, to be general, we allow additional interactions coupling the sectors and treat T_d as a free parameter.

Assuming the Yukawa couplings are Z_2 symmetric, the mirror charged lepton masses are determined solely by the ratio v'/v . (The masses of mirror hadrons could be also affected by the Z_2 symmetry breaking in the $SU(3)_c$ gauge coupling.) As their masses are relatively small, they remain in the thermal bath until low temperature, and contribute too much dark radiation. In figure 1, we show the prediction for the abundance of dark radiation, which by convention is expressed as an effective extra number of neutrinos, ΔN_{eff} , which is given by

$$\Delta N_{\text{eff}} = \frac{4}{7} g'_r \times \left(\frac{10.75}{g(T_d)} \right)^{4/3} \times \left(\frac{g'(T_d)}{g'_r} \right)^{4/3}, \quad (3.1)$$

where $g(T)$ and $g'(T)$ are the effective entropy degrees of freedom (d.o.f) of the Standard Model particles and the mirror particles at temperature T , respectively, and g'_r is the d.o.f. of the radiation component of the mirror sector. The second factor expresses the heating of the Standard Model neutrinos, and the third factor expresses the heating of the dark radiation. The d.o.f of the Standard Model particles $g(T)$ is extracted from [17]. We assume that only the mirror electron, muon, tau and photon are light and contribute to the dark radiation. Light mirror neutrinos, as considered in sections 4 to 7, make the claim in this section even stronger. Then $g'_r = 2$ and

$$\begin{aligned} g'(T_d) &= 2 + \frac{45}{2\pi^2} \frac{\rho'(T_d) + p'(T_d)}{T_d^4}, \\ \rho'(T_d) &= \frac{2}{\pi^2} \sum_{f=e,\mu,\tau} m_f^4 \left(\frac{v'}{v} \right)^4 \int_1^\infty dx \frac{(x^2 - 1)^{1/2} x^2}{\exp(xm_f v'/T_d v) + 1}, \\ p'(T_d) &= \frac{2}{3\pi^2} \sum_{f=e,\mu,\tau} m_f^4 \left(\frac{v'}{v} \right)^4 \int_1^\infty dx \frac{(x^2 - 1)^{3/2}}{\exp(xm_f v'/T_d v) + 1}. \end{aligned} \quad (3.2)$$

The Planck collaboration puts a bound on the effective number of neutrinos, $N_{\text{eff}} = 3.2 \pm 0.5$ [21], which leads to the upper bound of $\Delta N_{\text{eff}} < 0.65$ (2σ) and 0.40 (1σ), indicated

by dashed lines in figure 1. The 2σ bound can be only marginally satisfied. The 1σ bound can be satisfied when $v'/v \gtrsim 40$, which requires a fine-tuning of more than 0.1% to obtain the electroweak symmetry breaking scale. Note the importance to reach this conclusion of the deviation of the actual $g'(T_d)$ from a naive stepwise function. We conclude that it is necessary to break the Z_2 symmetry in the Yukawa couplings to further raise the mirror charged lepton masses.

The Yukawa couplings can naturally acquire Z_2 symmetry breaking if they arise from VEVs of fields, as in the Froggatt-Nielsen (FN) mechanism [22]. Once these fields take asymmetric VEVs, the Z_2 symmetry of the Yukawa couplings is spontaneously broken. As long as the top Yukawa coupling does not depend on these field values, the Twin Higgs mechanism is maintained. In this paper we do not specify the model of the FN mechanism, but study the physical consequences of a low energy effective field theory for Twin Higgs with $y_{f'} \neq y_f$.

4 Minimal mirror twin Higgs

The effective field theory below Λ_{TH} for Minimal Mirror Twin Higgs is

$$\begin{aligned} \mathcal{L}_{\text{EFT}} = & \mathcal{L}_{321}(g_i, y_f, \lambda, m_H) + \mathcal{L}'_{3'2'1'}(g_{i'}, y_{f'}, \lambda', m_{H'}) + 2\lambda'' H^\dagger H H'^\dagger H' + \frac{1}{2} \frac{\epsilon}{\cos\theta_W^2} B^{\mu\nu} B'_{\mu\nu} \\ & + (LH)^2/M_1 + (L'H')^2/M_1 + (LH)(L'H')/M_2 + \dots \end{aligned} \quad (4.1)$$

where $\mathcal{L}_{321}(g_i, y_f, \lambda, m_H)$ describes the Standard Model including all operators up to dimension 4 and $\mathcal{L}'_{3'2'1'}$ similarly describes the mirror sector. The lepton-Higgs interactions of the second line are the most general Z_2 symmetric set of dimension 5 operators and are suppressed by large masses $M_{1,2}$. Including all operators consistent with gauge invariance allows kinetic mixing via ϵ .

The Twin Higgs mechanism is imposed by boundary conditions at Λ_{TH} :

$$\lambda'' = \lambda', \quad m_{H'} = m_H, \quad g_{i'} = g_i, \quad y_{t'} = y_t. \quad (4.2)$$

As discussed in footnote 1, there could also be a boundary condition giving non-zero $\Delta\lambda$. Z_2 breaking is the minimal consistent with the requirement of the previous section, namely

$$y_{f'} \neq y_f \quad f \neq t. \quad (4.3)$$

This breaking of Z_2 is hard, meaning that the boundary conditions of (4.2) are not exact and are broken by loop corrections at Λ_{TH} . We restrict the size of $y_{f'}$, for $f \neq t$, so that these corrections maintain the Twin Higgs mechanism and satisfy eqs. (2.16), (2.18), (2.20). The dimension 5 operators have flavor structure suppressed and would arise from the see-saw mechanism with Z_2 symmetric neutrino Yukawa couplings and right-handed neutrino masses. Comparable $M_{1,2}$ are excluded from limits on oscillation to sterile neutrinos, suggesting that lepton number symmetries (or $B - L$ symmetries) play an important role in neutrino masses.⁴ We assume these symmetries lead to neutrino couplings $LNH + L'N'H'$

⁴If the hierarchy between M_1 and M_2 is not too large, the theory may have neutrino oscillation signals.

and masses for right-handed neutrinos, N and N' , so that $M_2 \gg M_1$ or $M_1 \gg M_2$; the light neutrinos are then Majorana or Dirac, respectively.

For $M_2 \gg M_1$, arising if NN' mass mixing is absent, neutrinos are Majorana with light mirror neutrino masses given by

$$m_{\nu'} = (v'/v)^2 m_\nu. \quad (4.4)$$

For $M_1 \gg M_2$, arising if the only right-handed neutrino mass is NN' , Standard Model and mirror left-handed neutrinos obtain Dirac masses, so that

$$m_{\nu'} = m_\nu. \quad (4.5)$$

Such Dirac masses from a seesaw are an interesting possible consequence of parity restoration in the mirror scheme.

Next we constrain the Z_2 breaking in $y_{f'}$ by requiring that quantum corrections to the Higgs masses yield non-zero Δm_H^2 as required for the Twin Higgs mechanism in eq. (2.9). We then place experimental bounds on $y_{f'}$ from the invisible decay width of the SM-like Higgs. Finally, we study mirror fermion spectra, consistent with these constraints, that maximize the QCD' scale.

4.1 Constraint on Yukawa couplings from Δm_H^2

We derive a constraint on Z_2 symmetry breaking in the Yukawa couplings by requiring that quantum correction to the soft Higgs masses yields the Twin Higgs mechanism. We assume the top Yukawa couplings of the Standard Model and mirror sectors are identical but allow other Yukawa couplings to differ, inducing a Z_2 breaking Higgs mass term

$$\Delta m_H^2|_{\text{Yuk}} \simeq \sum_{f \neq t} \frac{N_{f'}}{8\pi^2} y_{f'}^2(\Lambda) \Lambda^2, \quad (4.6)$$

where $N_{f'}$ is the multiplicity of the mirror fermion f' ; 3 for mirror quarks and one for mirror leptons. It should be remarked that the sign of the mass term is the required one. For this correction to explain Δm_H^2 of eq. (2.9), the sum of the square of the mirror Yukawa couplings, and hence the mirror fermions masses, are determined

$$\begin{aligned} \sum_{f \neq t} \left(\frac{N_{f'}}{3} \right) y_{f'}^2(\Lambda) &\simeq 0.04 \left(\frac{v'/v}{3} \right)^2 \left(\frac{5 \text{ TeV}}{\Lambda} \right)^2, \\ \sum_{f=u,d,c,s,b} \left(\frac{N_{f'}}{3 \delta_{f',\Lambda}^2} \right) m_{f'}^2 &\simeq (100 \text{ GeV})^2 \left(\frac{v'/v}{3} \right)^4 \left(\frac{5 \text{ TeV}}{\Lambda} \right)^2. \end{aligned} \quad (4.7)$$

Here, $\delta_{f',\mu} \equiv y_{f'}(m_{f'})/y_{f'}(\mu)$ encodes the effect of the renormalization between a scale μ and $m_{f'}$. $\delta_{q',\Lambda}$ is about 1.3–1.5 for $m_{q'} = (100-10) \text{ GeV}$, and $\delta_{l',\Lambda} \simeq 1$. In the following we take $\delta_{q',\Lambda} = 1.4$. Note that the estimation of $\Delta m_H^2|_{\text{Yuk}}$ is UV sensitive and hence involves $O(1)$ uncertainty, which we formally treat by varying the value of Λ in eq. (4.7). The Z_2 breaking correction to the Higgs quartic couplings is proportional to $y_{f'}^4$ and is negligible.

4.2 Standard Model like Higgs decays

The Standard Model like Higgs h is an admixture of the two doublets H and H' , $h = c_\gamma H + s_\gamma H'$, with $s_\gamma^2 \equiv \sin^2 \gamma = (v/v')^2$ up to corrections of order $(v/v')^4$ and $(m_h/m_{h'})^2$. In turn this leads to a universal reduction by a factor c_γ of the Higgs couplings to all pairs of SM particles as well as to a coupling of the same Higgs to the mirror fermions

$$\mathcal{L} \supset -y_{f'} H' f'_L \bar{f}'_R \rightarrow -\frac{v}{\sqrt{2}v'} y_{f'} h f'_L \bar{f}'_R = -\frac{vm_{f'}}{\sqrt{2}v'^2 \delta_{f',m_h}} h f'_L \bar{f}'_R, \quad (4.8)$$

where the QCD running factor for a mirror quark $f' = q'$, δ_{q',m_h} , is about 1.3 for $m_{q'}$ around 10 GeV, and the precise value depends on the details of the mirror quark spectrum.⁵

Higgs decays to mirror fermions leads to an invisible branching ratio

$$\text{Br}_{\text{inv}} = \text{Br}(h \rightarrow f' \bar{f}') \simeq 0.1 \times \left(\frac{3}{v'/v}\right)^4 \sum_{f', 2m_{f'} < m_h} \frac{N_{f'}}{3} \left(\frac{m_{f'}}{10\text{GeV}}\right)^2 \delta_{f',m_h}^{-2}. \quad (4.9)$$

where phase space has been neglected. Figure 2 shows the correlations between v/v' , Br_{inv} , and the universal deviation from unity of the Higgs signal-strengths at the LHC into any SM final state,

$$1 - \mu = 1 - c_\gamma^2(1 - \text{Br}_{\text{inv}}) \simeq s_\gamma^2 + \text{Br}_{\text{inv}}, \quad (4.10)$$

versus the relevant combination of mirror fermion masses

$$\sum_{f', 2m_{f'} < m_h} m_{f'}^2 \frac{N_{f'}}{3} \left(\frac{1.3}{\delta_{f',m_h}}\right)^2 \equiv m_{\text{sum}}^2. \quad (4.11)$$

We adopt the bound $\mu > 0.75$ (95% C.L.) for the gluon fusion channel [13], as it has the smallest uncertainty. The bound is so strong that mirror fermions with $m_{f'} < m_h/2$ give sub-dominant contributions to Δm_H^2 (see eq. (4.7)): there must be at least one mirror fermion heavier than $m_h/2$ other than t' . The high-luminosity LHC can probe $\mu < 0.93$ [23], which is shown by a dashed line in figure 2. Irrespective of the mirror fermions masses, $v'/v < 4$ can be probed. The ILC is expected to measure the Higgs signal-strength with an accuracy of 1% [24], which probes $v'/v < 10$.

4.3 Mirror QCD phase transition temperature

The larger Yukawa couplings of the mirror quarks leads to a larger mass of the mirror quarks, and hence to a larger mirror QCD phase transition temperature, T'_c . Since the number of degrees of freedom of the mirror sector changes rapidly near the phase transition, increasing T'_c above the decoupling temperature of the two sectors is critical to obtaining ΔN_{eff} below the current bound, as specifically illustrated in section 5.3. However, there is a limit to how much T'_c can be increased, and here we derive an upper bound on T'_c .

First, we take the masses of N_+ mirror quarks (m_+) above $m_h/2$ and degenerate, and determine their mass according to eq. (4.7). Next, we take the masses of the remaining

⁵In the following, we take $\delta_{q',m_h} = 1.3$ for simplicity. This simplification essentially does not change the following discussion.




Figure 2. Prediction for the universal Higgs signal-strength as a function of a sum of mirror fermion masses defined in (4.11), for given v'/v . The dotted lines show contours of the invisible branching ratio, and the dashed line shows the sensitivity of the high-luminosity running of the LHC.




Figure 3. The prediction of the mirror QCD phase transition temperature as a function of the light mirror quark mass.

$5 - N_+$ mirror quarks to be the same (m_q), and constrained by the bound $\mu > 0.75$. With these masses, we solve the two-loop renormalization group equation of the mirror QCD gauge coupling constant. We find that, for a wide range of parameter space, m_q is well above the scale at which the gauge coupling diverges. Adopting the lattice calculation for the pure SU(3) Yang-Mills gauge theory in ref. [25], matching the renormalization scale and the inverse of the lattice spacing, we obtain T'_c , as shown in figure 3. The figure shows that T'_c can be well above 1 GeV, and reach a few GeV. Here the right end of each line shows the point where the bound from the invisible decay of the Higgs, as inferred from the limit on μ , is saturated. We also show values of $\text{Br}(h \rightarrow \text{inv}) = 0.02, 0.05, 0.1$ by dots on each line, from left to right.

Note that the sum of the square of Yukawa couplings is bounded as eq. (4.7) and $\mu > 0.75$, while the dynamical scale of mirror QCD depends on the product of the mirror quark masses. Thus, for a given N_+ , the universal m_+ and m_- saturating these bounds maximize T'_c : the maximal T'_c can be read off from the right hand end of each line.

5 Thermal history with Higgs exchange

In this section, we discuss the thermal history of the Minimal Mirror Twin Higgs theory of (4.1) in the limit of $\epsilon < 10^{-5}$, so that the sectors are coupled only via Higgs exchange, and focus on the amount of dark radiation. Note that radiative corrections to ϵ in the effective theory below Λ_{TH} vanish at 3 loops,⁶ and any 4-loop contribution would be of order 10^{-10} . In the early universe, at temperatures larger than several GeV, the Standard Model and mirror particles interact with each other and have the same temperature. Below some temperature, T_d , the interaction between the sectors becomes inefficient and they evolve independently. Heavy mirror particles eventually decay or annihilate into mirror photons and neutrinos, which are observed as dark radiation estimated in eq. (3.1). To satisfy observational constraints requires $g'(T_d) \ll g(T_d)$ so that at T_d the colored states u, d, s, g contribute to $g(T_d)$ but QCD' states contribute very little to $g'(T_d)$; roughly speaking, decoupling must occur between QCD' and QCD phase transitions, and we explore this further below.

5.1 QCD' and constraints on the q' spectrum

If some q' are light compared to the QCD' scale Λ' , then T'_c is lower than computed in the previous section resulting in large QCD' contributions to $g'(T_d)$ that are excluded by bounds on ΔN_{eff} . Hence the QCD' phase transition is purely gluonic with zero q' flavors. The temperature dependence of this zero flavor QCD, $g'_{\text{QCD}}(T)$ has been accurately computed on the lattice [26]. As the temperature decreases from high values, $g'_{\text{QCD}}(T)$ drops from its large perturbative value of 16 only very gradually, and then sharply drops very close to the critical temperature T'_c ; immediately after the phase transition at T'_c the glueball contribution to $g'(T'_c)$ is 0.6. If $T_d > 1.1 T'_c$, much of the entropy of the mirror gluon plasma is distributed solely to the mirror particles, which leads to too large ΔN_{eff} . Hence, in the next sub-section we seek regions of parameter space where $T_d < T'_c$, which is at most (1–2.8) GeV, as shown in figure 3.

Below T'_c , mirror glueballs S' decay to $\gamma'\gamma'$ via

$$\mathcal{L} \sim \frac{\alpha_s \alpha}{m_{q'}^4} G' G' F' F', \quad (5.1)$$

generated by integrating out the lightest mirror quark, of mass $m_{q'}$, giving a decay rate

$$\Gamma(S' \rightarrow \gamma'\gamma') = \frac{\alpha_s^2 \alpha^2 \Lambda'^9}{64\pi^3 m_{q'}^8} \sim 10^{-17} \text{ GeV} \left(\frac{\Lambda'}{2 \text{ GeV}} \right)^9 \left(\frac{20 \text{ GeV}}{m_{q'}} \right)^8. \quad (5.2)$$

Λ' is the scale at which QCD' becomes strongly coupled, and is comparable to T'_c . In the case with all q' heavier than $m_h/2$, labelled $N_+ = 5$ in figure 3, the mirror glueball is not in thermal equilibrium just above T'_c where its contribution to g' is large. In this case, the

⁶Candidates for such three-loop corrections are diagrams where a SM photon is connected with two Higgs with a loop of SM particles, and a mirror photon is connected with two mirror Higgs in a similar manner, and the two Higgs connect to two mirror Higgs via Higgs mixing. We find that the sub-diagram involving a photon and two Higgs vanishes due to the Bose statistic of the Higgs. Intuitively speaking, the symmetrized two Higgs have vanishing angular momentum, and vanishing correlators with a photon.

mirror glueballs decay late, well after T_d computed below, and hence is excluded by the limit on ΔN_{eff} . Figure 3 shows that, in all the remaining cases of $N_+ = 1-4$, there must be one q' lighter than 22 GeV. Over much of the parameter space of the q' spectrum, the decay rate of (5.2) is fast enough to ensure that the glueballs at T'_c are indeed in thermal equilibrium with γ' , and contribute 0.6 to $g'(T'_c)$. Cases with $\Gamma(\mathcal{S}' \rightarrow \gamma'\gamma')$ less than the Hubble expansion rate at T'_c must be discarded as they give too much dark radiation.

Mirror glueballs also decay to $\nu'\bar{\nu}'$ via

$$\mathcal{L} \sim \frac{\alpha_s \alpha_2}{m_q^2 m_{Z'}^2} \nu^\dagger \sigma \nu D G' G', \quad (5.3)$$

giving a decay rate

$$\Gamma(\mathcal{S}' \rightarrow \nu'\bar{\nu}') = \frac{\alpha_s^2 \alpha_2^2}{64\pi^3} \frac{\Lambda'^9}{m_q^4 m_{Z'}^4} \sim 10^{-20} \text{ GeV} \left(\frac{\Lambda'}{2 \text{ GeV}} \right)^9 \left(\frac{20 \text{ GeV}}{m_{q'}} \right)^4 \left(\frac{3}{v'/v} \right)^4, \quad (5.4)$$

which is negligible in comparison with $\Gamma(\mathcal{S}' \rightarrow \gamma'\gamma')$.

5.2 Decoupling temperature

Now, let us estimate the decoupling temperature T_d . Mirror fermions f' and standard fermions f interact with each other by the exchange of the standard model Higgs. Since f' interact with γ' , thermal equilibrium between γ' and Standard Model particles can be maintained, even if f' is heavy and its number density is much smaller than that of relativistic particles. We discuss the thermal equilibrium of mirror neutrinos later.

Let us first estimate the interaction rate assuming that the dynamics of f' in the thermal bath is described by that of free fermions. This is certainly correct for the mirror leptons. We comment on the case with mirror quarks later. The scattering cross section between f and f' by Higgs exchange is given by

$$\sigma v(ff' \rightarrow ff') = \frac{1}{8\pi} \left(\frac{m_f}{v} \right)^2 \left(\frac{v m_{f'}}{v'^2} \right)^2 \frac{m_f m_{f'}}{m_f + m_{f'}} \frac{p_{\text{cm}}}{m_h^4} \quad (5.5)$$

where we assume a non-relativistic limit. Here p_{cm} is the momentum of the fermion in the center of mass frame. In the thermal bath, it has a typical size

$$p_{\text{cm}}^2 = \frac{4T(m_f + m_{f'} + \sqrt{m_f m_{f'}})}{3(2 + m_f/m_{f'} + m_{f'}/m_f)}. \quad (5.6)$$

The annihilation cross section of a pair of f' into a pair of f is given by

$$\sigma(f' \bar{f}' \rightarrow f \bar{f}) v = \frac{N_f}{4\pi} \left(\frac{m_f}{v} \right)^2 \left(\frac{v m_{f'}}{v'^2} \right)^2 \frac{(m_{f'}^2 - m_f^2)^{3/2}}{m_{f'}^3 m_h^4} p_{f'}^2. \quad (5.7)$$

Here $p_{f'}$ is the momentum of f' in the center of mass frame. In the thermal bath, it is as large as $p_{f'}^2 \simeq 3m_{f'}T/2$. N_f is the multiplicity of the Dirac fermion f ; for one lepton (quark), $N_f = 1(3)$.

The energy density of mirror particles, ρ' , is transferred into Standard Model particles at a rate

$$\begin{aligned} \frac{d}{dt}\rho' = & \sum_f 4(N_f n_F(m_f, T)) (N_{f'} n_F(m_{f'}, T)) \sigma v(ff' \rightarrow ff') \times \Delta E \\ & + \sum_f N_{f'} n_F(m_{f'}, T)^2 \sigma v(f' \bar{f}' \rightarrow f \bar{f}) \times 2m_{f'}. \end{aligned} \quad (5.8)$$

Here $n_F(m, T)$ is the number density of a Weyl fermion of mass m in the thermal bath at temperature T , and $\Delta E \simeq T$ is a typical energy transfer by the scattering $ff' \rightarrow ff'$. We find that the annihilation $f' \bar{f}' \rightarrow f \bar{f}$ is more important than the scattering $ff' \rightarrow ff'$ in determining T_d by Higgs exchange. This is because the energy transfer per annihilation is $2m'_{f'}$ and is larger than that per scattering, $\Delta E \sim T$, for $T < m_{f'}$.

In figure 4, solid curves show the temperature T_d , below which $(d\rho'/dt)/\rho'$ is smaller than the expansion rate of the universe, as a function of $m_{f'}$ with various colors for different $N_{f'} = 1, 2, 3, 6, 9, 12, 15$. The black and red dashed lines show the maximal T'_c as estimated in section 4.3. Other dashed lines shows the maximal possible T'_c when $N_{f'}/3$ mirror quarks have a common mass $m_{f'}$, with their color chosen to match those of the corresponding solid lines. The maximal T'_c are determined in the following way. We first put as many quarks as possible (N_+) above $m_h/2$ with the same mass and determine their mass so that the bound in eq. (4.7) is saturated. We then determine the masses of remaining quarks ($5 - N_+ - N_{f'}/3$) so that the bound $\mu > 0.75$ is saturated. (For $v'/v = 3, 5$ and $\Lambda = 3$ TeV, we find that $N_+ = 5 - N_{f'}/3$, and hence the second step is not necessary.) On the right end of each dashed line, the invisible decay width of the Standard Model Higgs into mirror fermions f' saturate the current bound. For a mirror fermion mass in the range between about 5 and 20-30 GeV for $v'/v = 3-5$, T_d is smaller than T'_c , so that the energy of the mirror gluon plasma is efficiently transferred into the Standard Model particles.

In figure 5, we show the predictions for the invisible branching ratio of the Standard Model Higgs. The ranges of $m_{f'}$ which yield $T_d > T'_c$ are depicted by dotted lines. The shaded region is excluded by the measurement of the Higgs-signal strength. The high-luminosity LHC can probe $\mu < 0.93$, whose corresponding invisible branching ratio is shown by a dashed line in the right panel. It is also sensitive to the invisible branching ratio of 10% [23]. For $v'/v \sim 3$, the deviation of the Higgs signal-strength from unity as well as non-zero invisible branching ration may be detected in the high-luminosity running of the LHC. The ILC is sensitive to an invisible branching fraction of sub-percent level [24]. The region with $T_d < T'_c$ can be probed by the Higgs-signal strength as well as the invisible decay of the Higgs at the high-luminosity LHC and the ILC.

In the above analysis we calculated T_d assuming that the dynamics of mirror fermions is described by that of free fermions. This is not correct for mirror quarks when the temperature is smaller than the binding energy B_D of the mirror QCD interaction. As the temperature drops below the binding energy, some mirror quarks form bound states, namely mirror quarkonia. This effect is expected to enhance the energy transfer rate by the




Figure 4. The decoupling temperature, T_d , between the two sectors as a function of the mirror fermion mass, $m_{f'}$, for various $N_{f'}$ degenerate states labelled by color. The dashed lines show the maximal possible mirror QCD phase transition temperature.




Figure 5. Predictions for the invisible branching ratio of the Standard Model Higgs. The ranges of $m_{f'}$ that yield $T_d < (>)T'_c$ are depicted by solid (dotted) lines.

annihilation of mirror fermions. The annihilation rate of mirror fermions inside quarkonia is

$$\Gamma(q'\bar{q}' \rightarrow f\bar{f}) \sim \sigma(q'\bar{q}' \rightarrow f\bar{f})v|_{p_{f'} \simeq m_{q'}\alpha'_3} \times \frac{1}{4\pi} (m_{f'}\alpha'_3)^3, \quad (5.9)$$

where the second factor is the inverse volume of a quarkonium. The energy transfer rate by annihilation is given by

$$\frac{d}{dt}\rho'|_{\text{annihilation}} \sim N_{q'} \sum_f n_B(2m_{q'}, T) \Gamma(\eta_{q'} \rightarrow f\bar{f}) \times 2m_{q'}, \quad (5.10)$$

where $N_{q'}$ is the number of mirror quarks with a mass $m_{q'}$, and $n_B(m, T)$ is the number density of a real scalar with a mass m in the thermal bath with a temperature T . The ratio of the energy transfer rate by the annihilation of free fermions to that by the annihilation inside quarkonia is

$$\frac{d\rho'/dt|_{\text{free}}}{d\rho'/dt|_{\text{quarkonia}}} \sim \left(\frac{T}{m_{q'}\alpha'_3}\right)^{5/2} \sim \left(\frac{T}{E_B}\right)^{5/2}. \quad (5.11)$$

Here we have used the non-relativistic approximation for $n_{F,B}$. In the parameter space we have discussed, the binding energy E_B is comparable to the temperature, and hence the

formation of the bound state does not change the result in figure 4 significantly. But we note that it is possible that $T_d < T'_c$ is achieved for a wider parameter region.

Here we show that mirror neutrinos can be in thermal equilibrium down to T_d . Chemical equilibrium of mirror neutrinos is maintained by the annihilation process $f' \bar{f}' \leftrightarrow \nu' \bar{\nu}'$, with cross section

$$\sigma(f' \bar{f}' \rightarrow \nu' \bar{\nu}')v = \frac{g_2^4/c_w^4}{64\pi} (I_{3f'} - 2s_w^2 Q_{f'})^2 \frac{m_{f'}^2}{m_{Z'}^4}. \quad (5.12)$$

The number of mirror neutrinos produced/annihilated per unit volume and time is given by

$$\sigma(f' \bar{f}' \rightarrow \nu' \bar{\nu}')v \times n_F(m_{f'}, T)^2 \times N_{f'}. \quad (5.13)$$

Comparing this rate with $H \times n(\nu')$, we find that chemical equilibrium as well as kinetic equilibrium are maintained down to a temperature of about $m_{f'}/10$. Kinetic equilibrium alone is maintained by the scattering $f' \nu' \rightarrow f' \nu'$, with a cross section

$$\sigma(f' \nu' \rightarrow f' \nu')v = \frac{g_2^4/c_w^4}{32\pi} \left((I_{3f'} - 2s_w^2 Q_{f'})^2 + \frac{1}{4} \right) \frac{T^2}{m_{Z'}^4}, \quad (5.14)$$

and is effective down to a temperature of about $m_{f'}/20$. Comparing $m_{f'}/10$ and T_d in figure 4, thermal equilibrium of mirror neutrinos is also maintained down to temperature T_d .

5.3 Dark radiation

As we have shown, mirror photons and mirror neutrinos can be in thermal equilibrium with Standard Model particles down to temperature $T_d < T'_c$. Taking account mirror photons and neutrinos, the prediction for ΔN_{eff} is

$$\Delta N_{\text{eff}, \gamma' \nu'} = \frac{4}{7} \times \left(2 + \frac{7}{4} \times 3 \right) \times \left(\frac{10.75}{g(T_d)} \right)^{4/3} = 0.29 \times \left(\frac{80}{g(T_d)} \right)^{4/3}, \quad (5.15)$$

which is consistent with the upper bound, $\Delta N_{\text{eff}} < 0.65$.

To keep T_d smaller than T'_c , some mirror fermions must have masses not far above T_d and so they also contribute to dark radiation, giving

$$\Delta N_{\text{eff}, \gamma' \nu' f'} = 0.29 \times \left(\frac{80}{g(T_d)} \right)^{4/3} \left(\frac{7.25 + g'_f(T_d)}{7.25} \right)^{4/3}, \quad (5.16)$$

where g'_f is the effective d.o.f. of mirror fermions f' . In figure 6, we show the prediction for ΔN_{eff} as a function of $m_{f'}$. Here we have neglected the contribution from the mirror gluon plasma, which is correct for $m_{f'}$ that give $T_d < T'_c$. Regions that give $T_d > T'_c$ are depicted by dotted lines. We also assume that $g_{s, f'}$ is well approximated by that of the ideal gas of f' . This is correct for mirror leptons; for mirror quarks, the actual $g_{s, f'}$ is smaller. Lines in the left panel are terminated if the Higgs coupling-strength falls below 0.75. The predicted amount of the dark radiation can be consistent with the experimental bound.

To summarize: for the amount of dark radiation to be below the experimental bound, there must be a mirror fermion with mass in the range (4 – 28) GeV.




Figure 6. Prediction for ΔN_{eff} as a function of the mirror fermion mass $m_{f'}$, for various multiplicities, $N_{f'}$. Thermal equilibrium between the two sectors is maintained by the exchange of the SM-like Higgs. The ranges of $m_{f'}$ that yield $T_d < (>)T'_c$ are depicted by solid (dotted) lines.

5.4 Cosmological signals of mirror neutrino masses

As discussed in section 4, the dimension 5 operators of the Minimal Mirror Twin Higgs theory of (4.1) result in neutrino masses that are either Majorana, with $m_{\nu'} = (v'/v)^2 m_\nu$, or Dirac with $m_{\nu'} = m_\nu$. Such masses can have significant effects on structure formation in the universe as well as the CMB spectrum. Taking into account dilution by Standard Model particles, the ν' number density is

$$n_{\nu'} = \frac{10.75}{g(T_d)} \times \frac{g'(T_d)}{g'_r} \times n_\nu = \left(\frac{7}{29} \Delta N_{\text{eff}} \right)^{3/4} n_\nu. \quad (5.17)$$

For the case of Majorana neutrinos, the effective total mass of light neutrinos, constrained by data on structure formation, is

$$\begin{aligned} \left(\sum m_\nu \right)_{\text{eff}} &\equiv \sum m_\nu + \sum m_{\nu'} \frac{n_{\nu'}}{n_\nu} \\ &= \left(\sum m_\nu \right) \times \left(1 + \left(\frac{7}{29} \Delta N_{\text{eff}} \right)^{3/4} \left(\frac{v'}{v} \right)^2 \right) > 2.3 \sum m_\nu. \end{aligned} \quad (5.18)$$

Here we have used the prediction of figure 6, $\Delta N_{\text{eff}} > 0.3$, and the experimental constraint $v'/v > 3$, to obtain the last inequality. Although the current cosmological data are more constraining on ΔN_{eff} than on $(\sum m_\nu)_{\text{eff}}$, both parameters may play a comparably important role in observations in the near future. We note that $(\sum m_\nu)_{\text{eff}}$ can be larger or smaller if the Z_2 symmetry is also broken in the Dirac mass term of neutrinos.

For the case of Dirac neutrinos, the effective total mass of neutrinos is

$$\left(\sum m_\nu \right)_{\text{eff}} = \left(\sum m_\nu \right) \times \left(1 + \left(\frac{7}{29} \Delta N_{\text{eff}} \right)^{3/4} \right) \simeq \sum m_\nu \quad (5.19)$$

so that $m_{\nu'}$ have a small effect on structure formation and the CMB spectrum.

6 Thermal history including kinetic mixing

In the last section, we discussed the thermal history of the Minimal Mirror Twin Higgs theory with $\epsilon < 10^{-5}$, so that Higgs exchange is the unique interaction coupling the two

sectors, and found that light mirror fermion must be in the range of about (4-20) GeV. In this section we allow larger values of ϵ , from the UV completion above Λ_{TH} , so that the Standard Model and mirror sectors also interact by kinetic mixing between $U(1)_Y$ and $U(1)'_Y$ gauge bosons, described by the $\epsilon B^{\mu\nu} B'_{\mu\nu}$ term of eq. (4.1). As we will see, the allowed range of mirror fermions masses are wider than the case without the kinetic mixing.

6.1 Decoupling temperature

Here we list the decoupling temperatures of various processes that maintain thermal equilibrium of mirror photons and/or neutrinos. We take a field basis such that the mirror photon is shifted to eliminate kinetic mixing, $A' \rightarrow A' + \epsilon A$. In this basis, Standard Model charged particles interact only with photons, while mirror particles interact with both photons and mirror photons.

Mirror photons are in thermal equilibrium with mirror charged fermions f' , which also interact with photons and through mixing, maintaining thermal equilibrium between Standard Model particles and mirror photons. The cross section for $f'\gamma' \leftrightarrow f'\gamma$ is given by,

$$\sigma(f'\gamma' \leftrightarrow f'\gamma)v = \frac{8\pi}{3}\epsilon^2\alpha^2q_{f'}^4\frac{1}{m_{f'}^2}, \tag{6.1}$$

where we assume $T \ll m_{f'}$, and $q_{f'}$ is the electromagnetic charge of f' . The scattering rate is smaller than the expansion rate of the universe below a temperature $T_{d,\gamma'}$,

$$T_{d,\gamma'} \simeq \frac{m_{f'}}{20 + 2\ln\frac{\epsilon}{10^{-2}}}. \tag{6.2}$$

$U(1)$ kinetic mixing also mixes the mirror Z' boson and the Standard Model photon, allowing mirror neutrinos to interact with Standard Model fermions with a cross section,

$$\sigma(f\nu' \rightarrow f\nu')v \simeq \sigma(f\bar{f} \leftrightarrow \nu'\bar{\nu}')v \simeq \frac{16\pi}{3}q_f^2\frac{\alpha^2}{c_w^4}\epsilon^2\frac{T^2}{m_{Z'}^4}. \tag{6.3}$$

This scattering becomes ineffective below the temperature $T_{d,\nu'}$,

$$T_{d,\nu'} \simeq 0.3 \text{ GeV} \left(\frac{\epsilon}{10^{-2}}\right)^{-2/3} \left(\frac{\nu'/v}{3}\right)^{4/3}. \tag{6.4}$$

$T_{d,\nu'}$ must be larger than the QCD phase transition temperature, otherwise the mirror neutrino abundance exceeds the upper bound on dark radiation. This gives an upper bound on kinetic mixing, $\epsilon \lesssim 10^{-2}$, so that $T_{d,\gamma'} \gtrsim m_{f'}/20$.

Mirror photons are in the thermal equilibrium with mirror charged fermions f' . If the scattering rate of the process $f'\bar{f}' \leftrightarrow \nu'\bar{\nu}'$ is large enough, kinetic as well as chemical equilibrium of mirror neutrinos is maintained. As we have seen, this interaction is effective down to the temperature of $m_{f'}/10 \equiv T_{d,\nu'\gamma',\text{che}}$. Kinetic equilibrium of mirror neutrinos is maintained by $f'\nu' \rightarrow f'\nu'$, which is efficient down to a temperature of $m_{f'}/20 \equiv T_{d,\nu'\gamma',\text{kin}}$.




Figure 7. Decoupling temperatures of various process when the lightest mirror charged fermion is a quark of mass 20 GeV.

6.2 Dark radiation

From the above consideration on decoupling temperatures, we obtain a bound on mirror fermion masses. In order for the energy of mirror gluons to be transferred into Standard Model particles, $T_{\gamma'}$ or $\max(T_{\nu'}, T_{d,\nu'\gamma',\text{kin}})$ must be smaller than T'_c . This requires a charged mirror fermion lighter than $20T'_c = 20\text{--}50$ GeV. Charged mirror fermion masses are also bounded from below. Even if mirror and Standard Model sectors decouple just before the QCD phase transition, ΔN_{eff} is too large unless charged mirror leptons have masses larger than 2 GeV. Since mirror quarks are bound to form mesons below T'_c , they can be lighter than 2 GeV as long as the meson masses are above 2 GeV. The allowed range of mirror fermion masses is wider than the case without the kinetic mixing.

Here we estimate ΔN_{eff} for a representative point of parameter space, illustrating the importance of a variety of reactions between mirror and QCD phase transitions. Suppose that the lightest mirror charged fermion is a quark of mass 20 GeV, so that mirror glueballs decay into mirror photons just below T'_c . In figure 7, we show the decoupling temperatures of various processes as a function of ϵ . We also show the mirror QCD phase transition temperature, which we assume to be the maximal one we estimated in section 4.3 for $v'/v = 3$ and $\Lambda = 3$ TeV. In region A, kinetic mixing is insufficient to transfer the energy of the mirror gluon plasma into Standard Model particles, and ΔN_{eff} is determined by Higgs exchange. In region B ΔN_{eff} is given by that in eq. (5.15) with $T_d = T_{d,\gamma'}$. In region C, the number density of mirror neutrinos per comoving volume is conserved below $\min(T_{d,\nu'\gamma',\text{che}}, T_{d,\nu'}) \equiv T_{d,\nu',\text{che}}$, giving

$$\Delta N_{\text{eff}} \simeq \frac{4}{7} \times 2 \left(\frac{10.75}{g(T_{d,\gamma'})} \right)^{4/3} + 3 \left(\frac{10.75}{g(T_{d,\gamma'})} \right)^{4/3} \left(1 - \frac{4860\zeta(3)^2}{7\pi^6} \frac{g(T_{d,\nu',\text{che}}) - g(T_{d,\gamma'})}{g(T_{d,\nu',\text{che}}) + 2} \right). \quad (6.5)$$

The last factor in the second term accounts for the conservation of the comoving number density of mirror neutrinos. However, we find this factor is ≥ 0.94 , so that ΔN_{eff} is well approximated by eq. (5.15) with $T_d = T_{d,\gamma'}$. In region D, ΔN_{eff} is given by eq. (5.15) with

$T_d = T_{d,\nu'}$. In region E, mirror photons decouple from the thermal bath at $T_{d,\nu'\gamma',\text{che}}$, while mirror neutrinos decouple at $T_{d,\nu'}$, giving

$$\Delta N_{\text{eff}} = \frac{4}{7} \times 2 \left(\frac{10.75}{g(T_{d,\nu'})} \right)^{4/3} \left(\frac{g(T_{d,\nu'}) + 5.25}{g(T_{d,\nu'\gamma',\text{kin}}) + 5.25} \right)^{4/3} + 3 \left(\frac{10.75}{g(T_{d,\nu'})} \right)^{4/3}. \quad (6.6)$$

We find that in region B, C, and D, ΔN_{eff} is about 0.3. In region E, ΔN_{eff} is larger than 0.3, and can saturate the bound on ΔN_{eff} for $\epsilon \simeq 10^{-2}$.

If the lightest mirror quark is heavier than 20 GeV, mirror gluons do not decay immediately below T'_c , but decay later. This is excluded if kinetic mixing is absent, because T'_c and T_d are close to each other. With kinetic mixing, the energy of mirror photons can be transferred into SM particles well below T'_c , if a charged mirror lepton is light enough. As long as mirror glueballs decay into mirror photons before the QCD phase transition, sufficiently light mirror charged leptons can suppress ΔN_{eff} to be within the allowed range.

6.3 Milli-charged particle

With U(1) kinetic mixing, mirror fermions of mirror charge $q_{f'}$ can be understood to carry SM electric charge $\epsilon q_{f'}$. In the range $m_{f'} = O(1-10)$ GeV, the most stringent constraint comes from collider experiments [28, 29], which is much weaker than the bound from ΔN_{eff} , $\epsilon \lesssim 10^{-2}$. A proposed search at the LHC can search down to mixings of $\epsilon = 10^{-2} - 10^{-3}$ [30].

7 Mirror baryon dark matter

The mirror sector, like the SM, possesses accidental baryon and lepton symmetries. Mirror baryons and leptons may account for dark matter in the universe [31] (see [32–34] for earlier work on astrophysical considerations of mirror baryons and leptons.). We assume a non-zero mirror matter-antimatter asymmetry with the asymmetric component comprising dark matter. The proximity of the energy densities of baryons and dark matter could be understood if the sectors which generate the Standard Model and mirror asymmetry are close to Z_2 symmetric, and the dark matter mass is $O(1-30)$ GeV.

We consider the mirror up quark u' , down quark d' and electron e' as possible components of dark matter. To simplify the discussion, and motivated by section IVC, we take the masses of u' and d' larger than the dynamical scale of mirror QCD, so that the masses of mirror baryons mainly originate from mirror quark masses.

7.1 Dark matter candidates

We consider dark matter candidates in Minimal Mirror Twin Higgs, where the Z_2 symmetry is broken solely by Yukawa couplings. A dark matter candidate depends on the mass spectrum of u' , d' and e' . Below we consider several interesting possibilities.

7.1.1 Light u' , d' : mirror neutron

Taking e' sufficiently heavy, as the temperature drops below $m_{e'}$, it decays into u' , d' and ν' and its abundance becomes negligible. For the remaining u' , d' , number changing processes from W' exchanges are absent, so that u' and d' are both stable and have separately

conserved comoving numbers. Mirror charge neutrality implies $n_{d'} = 2n_{u'}$. Below the mirror QCD phase transition temperature, u' and d' are combined into mirror baryons B' and mesons M' . The stable hadrons are

$$B'_{uuu}, B'_{uud}, B'_{udd}, B'_{ddd}, M'_{u\bar{d}},$$

with an obvious notation. The meson $M'_{u\bar{d}}$ is captured by mirror baryons, e.g.

$$M'_{u\bar{d}} + B'_{ddd} \rightarrow B'_{udd} (+\gamma').$$

As the sum of the masses of $M'_{u\bar{d}}$ and B'_{ddd} is larger than the mass of B'_{udd} by $2m_{d'}$, the meson $M'_{u\bar{d}}$ disappear from the thermal bath by this capture process. Finally, B' scatter with each other and almost all become the mirror neutron B'_{udd} . For example, the scattering process

$$B'_{uud} + B'_{ddd} \rightarrow 2B'_{udd} (+\gamma')$$

eliminates B'_{uud} and B'_{ddd} from the thermal bath. Note that the sum of the masses of B'_{uud} and B'_{ddd} is larger than twice of the mass of B'_{udd} as the mirror baryon B'_{ddd} is spin-3/2 and has a contribution to its mass from a spin-spin interaction.

Without any IR effects, the scattering cross section between mirror neutrons is $O(m_{B'_{udd}}^{-2})$, and does not affect structure formation. The cross section can be enhanced up to the unitarity limit by some IR effects, e.g. the Sommerfeld effect [35, 36] or the existence of resonance states [37]. In our case, mirror pions are also heavy due to large mirror quark masses, and those effects are expected to be suppressed.

7.1.2 Light d' , e' : mirror atom

If u' is sufficiently heavy, it decays to d' , e' and ν' and disappears from the thermal bath. The mirror electron e' is now stable, as its decay is kinematically forbidden. After the QCD' phase transition, d' combines into B'_{ddd} . To preserve charge neutrality, there are the same number of \bar{e}' . The mirror baryons B'_{ddd} and positrons \bar{e}' eventually “recombine” into mirror atoms.

The self scattering cross section of mirror atoms is given by [38]

$$\sigma/m_D \simeq \frac{100}{\alpha^2} \frac{(R+1)^4}{R^2} \frac{1}{m_D^3} = 6.6 \text{ cm}^2/\text{g} \times \left(\frac{10 \text{ GeV}}{m_D} \right)^3 \frac{(R+1)^4}{16R^2}, \quad (7.1)$$

where $R = \max(m_{B'_{ddd}}/m_{e'}, m_{e'}/m_{B'_{ddd}})$ and m_D is the mass of the dark atom. Here we assume that $R \sim 1$ and the kinetic energy of the dark atom is much smaller than the binding energy. The cross section is minimized for $R = 1$, which we assume in the following. The cored dark matter halo profiles could be explained by $m_D \sim 10 \text{ GeV}$. The upper bound on the cross section in dwarf galaxies, $\sigma/m_D < 10 \text{ cm}^2/\text{g}$ [39] requires that $m_D > 9 \text{ GeV}$. The constraint from galaxy clusters is weak, since the velocity dispersion of dark matter is so large that its kinetic energy is comparable to the binding energy, and the self interaction cross section is suppressed.

The recombination of mirror baryons and electrons is incomplete. From the numerical estimation in ref. [40] we obtain the ionized fraction,

$$x_{e'} \simeq 0.02 \times \left(\frac{m_D}{10 \text{ GeV}} \right)^2 \frac{4R}{(R+1)^2}. \quad (7.2)$$

The ionized components interact with each other and with atomic dark matter strongly, and may affect halo shape [40] and the scattering of clusters. The estimation of these and other constraints on the ionized fraction is beyond the scope of this paper.

With the sizable kinetic mixing that we considered in section 6, the ionized components participate in acoustic oscillations and affect the CMB spectrum [41, 42].⁷ The upper bound on the ionized fraction is $\Omega_{\text{ionized}}/\Omega_{\text{atom}} < 0.01$ [43] which, together with the limit from self scattering in dwarfs, disfavors atomic dark matter for such kinetic mixing.⁸

7.1.3 Light u' , e' : mirror atom

If d' is sufficiently heavy, B'_{uuu} and e' are stable and eventually form a mirror atom, $(B'_{uuu} + 2e')$ with a binding energy larger than that of $(B'_{ddd} + \bar{e}')$. The constraints from self interactions and the ionized components of the atomic dark matter are weakened. The precise determination of the constraints is beyond the scope of this paper.

7.1.4 Light u' , d' , e' : mirror neutron

Let us assume that u' , d' , e' are all light, with a spectrum such that e' , u' , d' are stable, $m_{u'} + m_{d'} > m_{e'}$, $m_{e'} + m_{d'} > m_{u'}$, $m_{e'} + m_{u'} > m_{d'}$. (Other mass spectra belong to one of the fore-mentioned three cases.) Then the following mirror particles are stable,

$$B'_{uuu}, B'_{uud}, B'_{udd}, B'_{ddd}, e'.$$

With u' , d' , and e' all light, reactions mediated by W' result in the removal of the e' , for example by $e'u' \rightarrow \nu'd'$. Scattering among the various B' remove B'_{uuu} and B'_{ddd} and mirror charge neutrality implies that only the mirror neutron remains.

In the expanding universe, however, the W' mediated interaction may freeze out before mirror protons and mirror electrons are removed from the thermal bath. At $T \lesssim m'$, where m' is the mass scale of mirror electrons and mirror protons, the freeze-out temperature is given by solving

$$\frac{g_2^4}{8\pi} \frac{m'^2}{m_{W'}^4} \times \frac{\rho_{\text{DM}}/s}{m'} s(T) = H(T), \quad (7.3)$$

where $s(T)$ is the entropy density, and $\rho_{\text{DM}}/s \simeq 3.6 \times 10^{-9} \text{ GeV}$ is the energy density of dark matter divided by the entropy density. The first factor in the left-hand side is the cross

⁷Mirror baryons and electrons interact with SM particles via kinetic mixing and are heated, which may change the ionized fraction.

⁸The galactic magnetic field prevents the ionized component from entering the disk, and ionized components initially in the disk are likely to escape the disk owing to Fermi acceleration by Super-Nova remnants. Furthermore, the magnetic field of the Earth prevents any ionized component from reaching the Earth. Hence, the ionized component may evade direct detection experiments performed on the Earth [44].

section of the W' mediated process, and the second factor is a rough estimate on the number density of mirror protons and electrons. The freeze-out temperature is then given by

$$T_{\text{fo},W'} \simeq 1 \text{ GeV} \times \frac{5 \text{ GeV}}{m'} \left(\frac{v'/v}{3} \right)^4. \quad (7.4)$$

If the freeze-out temperature is much smaller than $m'_p + m'_e - m'_n$, mirror protons and mirror electrons disappear from the thermal bath. This is the case for $m' \gtrsim 10 \text{ GeV}$. For $m' \lesssim 10 \text{ GeV}$, a non-negligible amount of mirror electrons and protons remain. Some of them later recombine into mirror atomic states. Unlike the cases of sections 7.1.2 and 7.1.3, these states decay into mirror neutrons and mirror neutrinos through mirror electron capture. Thus there is no constraint from the self-interaction of atomic dark matter. However, there is a constraint on the ionized component.

7.2 Direct detection of dark matter

The above dark matter candidates, N' , interact with Standard Model nucleons, N , through the exchange of the Standard Model Higgs, and may be observed in direct detection experiments [45–47]. The interaction of the Standard Model Higgs relevant for the direct detection is given by

$$\begin{aligned} \mathcal{L} &= \frac{h}{\sqrt{2}v} \left[- \left(\frac{v}{v'} \right)^2 \sum_{f' \in N'} m_{f'} f' \bar{f}' - \sum_{q=u,d,s} m_q q \bar{q} + \sum_{q=c,b,t} \frac{\alpha_3}{12\pi} \left(1 + \frac{11}{4} \frac{\alpha_3(m_q)}{\pi} \right) G_{\mu\nu}^a G^{a\mu\nu} \right] \\ &= \frac{h}{\sqrt{2}v} \left[- \left(\frac{v}{v'} \right)^2 \sum_{f' \in N'} m_{f'} f' \bar{f}' - \sum_{q=u,d,s} m_q q \bar{q} + 3.5 \times \frac{\alpha_3}{12\pi} G_{\mu\nu}^a G^{a\mu\nu} \right], \end{aligned} \quad (7.5)$$

where we have taken into account the one-loop QCD correction to the coupling with gluons [48]. The relevant matrix elements of N' is given by

$$\sum_{f' \in N'} \langle N' | m_{f'} f' \bar{f}' | N' \rangle = 2m_{N'}^2. \quad (7.6)$$

Here we assume that the mass of N' is mainly given by the masses of mirror fermions. Using the trace anomaly formula for the Standard Model nucleon [49],

$$2m_N^2 = \langle N | T_\mu^\mu | N \rangle = -\frac{9\alpha_3}{8\pi} \langle N | G_{\mu\nu}^a G^{a\mu\nu} | N \rangle + \sum_{q=u,d,s} \langle N | m_q q \bar{q} | N \rangle, \quad (7.7)$$

and matrix elements derived by a lattice calculation [50],⁹

$$\sum_{q=u,d,s} \langle N | m_q q \bar{q} | N \rangle \simeq 0.1 \times 2m_N^2, \quad (7.8)$$

the scattering cross section between N' and N through Higgs exchange is given by

$$\sigma_{NN'} = \frac{0.028}{\pi} \frac{m_{N'}^2 m_N^2}{v'^4 m_h^4} \left(\frac{m_N m_{N'}}{m_N + m_{N'}} \right)^2. \quad (7.9)$$

⁹This matrix element, obtained from a lattice calculation, is consistent with the one extracted from hadron scattering data with the aid of chiral perturbation theory [51, 52].




Figure 8. The scattering cross section between dark matter N' and a Standard Model nucleon N as a function of the mass of dark matter. Here we assume that the mass of dark matter is given by mirror fermion masses, and the matrix element of the trace anomaly is saturated by the mirror fermion mass terms.

If kinetic mixing is absent, $m_{N'}$ is bounded from below. Let us consider the mirror neutron, which is free of constraints from self interactions as well as from the efficiency of mirror recombination. It is made of one u' and two d' and the lines in figure 6 for $N_{f'} = 6$ show that the constraint on ΔN_{eff} requires the mirror neutron to be heavier than 19 GeV (26 GeV) at the 2σ level and 36 GeV (51 GeV) at the 1σ level for $v'/v = 3$ (5). The mirror neutron mass of $m_{N'} = O(1-30)$ GeV can explain the proximity of the energy densities of baryons and dark matter. Combined with the constraint from ΔN_{eff} , $m_{N'} = O(10)$ GeV is an interesting region. Note that, however, ΔN_{eff} is estimated by the ideal gas approximation of mirror quarks. The actual lower bound on $m_{N'}$ can be weaker.

In figure 8, we show the prediction on $\sigma_{NN'}$ as a function of $m'_{N'}$. The regions depicted by thin (dotted) lines are disfavored by ΔN_{eff} at the $1(2)\sigma$ level, if the mirror neutron is dark matter and kinetic mixing is absent. We show the constraints from the LUX experiment [53] and the Panda-II experiment [54] by solid lines. The higher mass region, $m_{N'} > 40$ (110) GeV is excluded for $v'/v = 3$ (5). We also show the sensitivity of the XENON1T experiment [55], the LZ experiment [56], and the neutrino floor [57] by dashed lines. We conclude that dark matter can be detected by near future experiments in the region consistent with the upper bound on ΔN_{eff} without kinetic mixing.

Once kinetic mixing between hypercharge gauge bosons is introduced, $m'_{N'}$ may be smaller and hence direct detection via Higgs exchange may be difficult to observe. In this case, however, direct detection via the kinetic mixing is possible. The mirror neutron dark matter interacts through its magnetic dipole moment. The magnetic moment of mirror neutron is expected to be of order

$$\mu_{N'} \sim \epsilon e \frac{1}{m_{N'}}. \tag{7.10}$$

Translating the bound derived in [58], we obtain the bound $\epsilon \lesssim 10^{-3}-10^{-4}$ for $m_{N'} = O(1-10)$ GeV. Projected low-threshold experiments such as CRESST III [59] and SuperCDMS SNOLAB [60] will probe smaller kinetic mixing.

Mirror atoms have a large radius and interact through screened charges of mirror baryons and electrons. The scattering cross section between atomic dark matter and a nucleus (per nucleon) is given by [61]

$$\sigma \simeq \frac{4\pi\alpha^2\epsilon^2}{\alpha_{\text{bind}}^4 m_{N'}^2} \simeq 10^{-34} \text{ cm}^2 \left(\frac{\alpha}{\alpha_{\text{bind}}}\right)^4 \left(\frac{\epsilon}{10^{-5}}\right)^2 \left(\frac{10 \text{ GeV}}{m_{N'}}\right)^2, \quad (7.11)$$

where α_{bind} is the fine-structure constant of the binding force that forms the atom. Here we assume $R \simeq 1$. A kinetic mixing that allows $T_d < T'_c$ ($\epsilon > 10^{-5}$) is excluded by various direct detection experiments. Especially, the light mass region ($m_{N'} = O(1) \text{ GeV}$) is excluded by CRESST II [62]).

8 Non-minimal Z_2 breaking from $B - L$ vevs

In this section we consider Z_2 symmetry breaking beyond Minimal Mirror Twin Higgs. While the SM neutrinos become very light via the seesaw mechanism, mirror neutrinos can be much heavier if the mirror right-handed neutrinos do not obtain large masses. This can be achieved if the Z_2 symmetry is also spontaneously broken by a large VEV breaking $B - L$ symmetry and a small/vanishing VEV breaking $B' - L'$.

8.1 Dark radiation

Let us first consider the case where a $B' - L'$ gauge field is absent, or the gauged $B' - L'$ symmetry is broken at an intermediate mass scale and hence the $B' - L'$ gauge field is heavy.¹⁰ The prediction on ΔN_{eff} in eq. (5.16) becomes

$$\Delta N_{\text{eff},\gamma'f'} = \frac{4}{7} \times 2 \left(\frac{10.75}{g(T_d)}\right)^{4/3} \left(\frac{2 + g'_f(T_d)}{2}\right)^{4/3} = 0.08 \left(\frac{80}{g(T_d)}\right)^{4/3} \left(\frac{2 + g'_f(T_d)}{2}\right)^{4/3}. \quad (8.1)$$

ΔN_{eff} can be as small as 0.08.

Next we consider the case with a massless $B' - L'$ gauge field. As we will see in the next sub-section, the massless $B' - L'$ gauge field is beneficial in identifying mirror baryons with dark matter. The prediction for the amount of the dark radiation are given by

$$\Delta N_{\text{eff},\gamma'f'} = \frac{4}{7} \times 4 \left(\frac{10.75}{g_{*s}(T_d)}\right)^{4/3} \left(\frac{4 + g'_f(T_d)}{4}\right)^{4/3} = 0.17 \left(\frac{80}{g(T_d)}\right)^{4/3} \left(\frac{4 + g'_f(T_d)}{4}\right)^{4/3}. \quad (8.2)$$

ΔN_{eff} can be as small as 0.17.

8.2 Additional dark matter candidates

The mirror neutrino masses now arise from Dirac Yukawa couplings and the lightest, ν' , could have a mass of order the lighter states of u' , d' and e' , or could be heavier than these light states. Also, a massless $B' - L'$ gauge field may alter dark matter phenomenology.

¹⁰In the supersymmetric Twin Higgs model, this is naturally achieved by a condensation of a mirror right-handed sneutrino at the TeV scale.

When $B' - L'$ is exact, so that $B' - L'$ charge is conserved, the B' and L' asymmetries must be produced after the mirror sphaleron process freezes out to avoid washout of B' and L' asymmetries. On the other hand, if the $B' - L'$ symmetry is only approximate, a $B' - L'$ asymmetry may be produced before the mirror sphaleron process freezes out.

8.2.1 Light u' , d' , ν' with unbroken $B' - L'$ gauge symmetry: mirror neutron and neutrino

Let us assume an unbroken $B' - L'$ gauge symmetry in the setup of section 7.1.1. Then the mirror neutron itself is no longer a dark matter candidate, as it feels a long-range force by the unbroken $B' - L'$ gauge field. However, a mirror neutron and a mirror neutrino, whose number densities are identical due to $B' - L'$ charge neutrality, can form an atom. The bound from self interactions as well as the ionized fraction is evaded for $\alpha'_{B-L} \gtrsim 10^{-2}$.

8.2.2 Light d' , e' , ν' with unbroken $B' - L'$ gauge interaction: mirror atom

With unbroken $B' - L'$ gauge symmetry in the setup in section 7.1.2, charge and $B' - L'$ neutrality remove ν' states, leaving $(B'_{ddd} + \bar{e}')$ atoms. The constraints from self interactions and the ionized fraction of atomic dark matter are evaded if $\alpha'_{B-L} \gtrsim 10^{-2}$.

8.2.3 Light u' , d' , e' with no light $B' - L'$ gauge boson: mirror atom and/or neutron

If ν' is sufficiently heavy, it decays to u' , d' and e' and disappears from the thermal bath. Reactions mediated by W' are absent, so that u' , d' and e' numbers are separately conserved. The following mirror particles are stable,

$$B'_{uuu}, B'_{uud}, B'_{udd}, B'_{ddd}, M'_{u\bar{d}}, e'.$$

Among them, B'_{uuu} , B'_{ddd} and $M'_{u\bar{d}}$ disappear as discussed in section 7.1.1. The asymmetry is stored in mirror protons B'_{uud} , neutrons B'_{udd} , and electrons e' , with abundances that depend on the B' and L' asymmetries. If $L' = 0$, only the mirror neutron remains, whereas if $B' - L' = 0$, as would occur if $B' - L'$ is unbroken during asymmetry genesis, only the mirror proton and the mirror electron remain.¹¹

If $B' - L'$ symmetry is broken at a sufficiently high scale, a non-zero $B' - L'$ asymmetry may be generated above the mirror electroweak phase transition temperature. After the mirror sphaleron process freezes out, the B' and L' asymmetries are given by

$$B' = \frac{1}{4}(B - L)', \quad L' = -\frac{3}{4}(B - L)'. \tag{8.3}$$

Here we assume that the matter content of the mirror sector is identical to the standard model plus three right-handed neutrinos just before the mirror sphaleron process freezes out. The resultant number densities of mirror protons, neutrons and electrons are given by

$$n_{e'} = n_{L'} = -\frac{3}{4}n_{B'-L'} = -\frac{3}{4}n_{n'}, \quad n_{p'} = n_{e'} = -\frac{3}{4}n_{n'}, \quad n_{n'} = n_{B'-L'} \tag{8.4}$$

¹¹These cases require asymmetry generation after the mirror sphaleron freeze out, as can be achieved in the supersymmetric Twin Higgs model, with Affleck-Dine baryogenesis [63, 64] from the $\bar{u}'\bar{d}'\bar{s}'$ flat direction.

Dark matter is composed of comparable numbers of mirror atoms and mirror neutrons. The constraints from self interactions and ionized fraction of atomic dark matter are relaxed.

9 Conclusions

The Twin Higgs mechanism significantly relaxes fine-tuning of the electroweak scale, and allows for a larger cut off scale. The cut off of the Standard Model is

$$\Lambda_{\text{SM}} \simeq 1.4 \text{ TeV} \times \left(\frac{\Delta_{\text{SM}}}{10} \right)^{1/2}, \quad (9.1)$$

while that of the Twin Higgs theory is

$$\Lambda_{\text{TH}} = 5.7 \text{ TeV} \times \left(\frac{\Delta_{\text{TH}}}{10} \right)^{1/2} \lambda^{1/2}. \quad (9.2)$$

The minimal theory has no new colored states to be produced at the LHC. It does offer the possibility of discovery modes at the LHC, such as production of the mirror Higgs via Higgs mixing; but the larger cut off may raise the masses of new particles above the LHC reach.

Mirror Twin Higgs models, however, predict the existence of extremely light particles, mirror photons and mirror neutrinos, that contribute to the dark radiation of the universe, leading to constraints on a realistic theory. We have found that, independent of the interactions that couple the two sectors, it is *necessary* to break the mirror Z_2 symmetry in the Yukawa couplings.

Minimal mirror twin Higgs. We have constructed a completely realistic effective field theory of Twin Higgs below the cut off Λ_{TH} . It contains a complete mirror sector, so that a UV completion, which we did not study, can restore spacetime parity symmetry. In Minimal Mirror Twin Higgs, the only Z_2 breaking arises from Yukawa couplings and the only communication between the sectors is from Higgs mixing, required by Twin Higgs, and kinetic mixing of hypercharge fields, allowed by gauge invariance. Furthermore, the Z_2 breaking Yukawa couplings not only suppress dark radiation to acceptable levels, but generate the Z_2 breaking Higgs mass term necessary for the Twin Higgs mechanism and raise the mirror baryon mass as required for realistic dark matter.

The Z_2 breaking Yukawa couplings induce a sizable invisible branching ratio of the SM-like Higgs boson through its mixing with the mirror Higgs. This, together with the reduction of the Higgs coupling to Standard Model particles, leads to a universal deviation from unity of the Higgs signal-strengths correlated with the masses of mirror fermions, as shown in figure 2. Irrespective of the mirror fermions masses, the high-luminosity running of the LHC and the ILC can probe $v'/v < 4$ and 10 respectively.

As illustrated in figure 4, it is non-trivial that Higgs mixing can lead to a decoupling temperature less than the QCD' phase transition temperature, necessary for a solution of the dark radiation problem. Figure 4 shows that, with Higgs mixing alone, there must be light mirror fermions with mass $m_{f'}$ in the range of about (4 – 28) GeV. The upper bound on $m_{f'}$ gives a lower bound of $\Delta N_{\text{eff}} \gtrsim 0.3$. The allowed range for $m_{f'}$ depends on $N_{f'}$, the

number of light mirror fermion states, and narrows considerably for larger values of $N_{f'}$. As the upper bound on $m_{f'}$ becomes tighter, ΔN_{eff} increases, which is shown in figure 6.

The lower bounds on $m_{f'}$ from figures 4 and 6 imply lower bounds on the invisible branching ratio of the SM-like Higgs boson and the universal deviation from unity of the Higgs signal-strengths. For $v'/v = 3$, the left panel of figure 5 shows that this branching ratio is typically in the range of 0.05-0.15 that can be probed by high luminosity running of the LHC. For $v'/v = 5$, the invisible branching ratio is reduced; the right panel of figure 5 shows that in some cases the signal is as small as 0.002 - 0.01, that could be probed by ILC. Essentially the entire parameter space can be probed.

If the kinetic mixing parameter ϵ is large enough to affect the thermal history of the two sectors near the QCD' phase transition, the allowed range of the light mirror fermion masses is enlarged. In section 6 we showed that the upper bound on $m_{f'}$ could be extended as high as 50 GeV, while mirror leptons could be as light as 2 GeV. A mirror quark can be lighter than 2 GeV provided the lightest mirror meson is heavier than 2 GeV. Even including kinetic mixing, the lower bound on ΔN_{eff} remains about 0.3, but the enlarged range of light mirror fermion masses is important for mirror dark matter.

Mirror baryons and leptons are natural candidates for dark matter. Dark matter can be composed of mirror neutrons, mirror atoms, or even a mixture of the two. Such dark matter can be directly detected via Higgs exchange, as illustrated in figure 8. For low v'/v , PandaX and LUX have recently excluded large values for the dark matter mass. In the absence of kinetic mixing, the region of dark matter masses allowed by present limits on dark radiation can be fully explored, up to the $1(2)\sigma$ limit by the XENON1T (LZ) experiment. Adding kinetic mixing to the thermal cosmological history, the limit on ΔN_{eff} is consistent with lighter dark masses that XENON1T and LZ are unable to probe. For mirror neutron dark matter masses in the (1-10) GeV region, present direct detection limits bound $\epsilon < 10^{-3} - 10^{-4}$ from scattering via a dipole moment. Mirror atomic dark matter with a sizable kinetic mixing such that the thermal history is affected is excluded.

In the Minimal Mirror Twin Higgs the seesaw mechanism yields light neutrino masses for both sectors. These neutrinos can be Majorana with those in the mirror sector heavier by a factor $(v'/v)^2$ than the observed neutrinos, leading to important effects in both CMB and LSS. The effective sum on neutrino masses relevant for cosmological data is at least 2.3 times greater than in the Standard Model. Small mixing between these Majorana standard and mirror neutrinos could lead to mirror neutrinos being observed as massive sterile neutrinos. Alternatively the seesaw could lead to Dirac neutrinos, with the mirror states as right-handed neutrinos degenerate with the observed left-handed states, leading to only very small effects on CMB and LSS. The predictions of eqs. (4.4), (4.5) for the masses of mirror neutrinos, however, rely on the assumption that Z_2 breaking does not substantially affect either the neutrino Yukawa couplings or right-handed neutrino masses, and therefore our predictions for the effective sum of neutrino masses, $(\sum m_\nu)_{\text{eff}}$, are less robust than the predictions for ΔN_{eff} , $\Gamma(h \rightarrow \text{inv})$ and the mirror baryon dark matter direct detection cross section.

Additional Z_2 breaking from $B - L$ vevs. A large $B - L$ vev can implement the seesaw mechanism for the known neutrinos, while a small or zero $B' - L'$ vev can lead to large Dirac mirror neutrino masses. Without light mirror neutrinos, the minimal ΔN_{eff} is lowered to 0.08 (0.17) without (with) a massless $B' - L'$ gauge field. Furthermore, there are new possibilities for mirror dark matter. If there is a massless $B' - L'$ gauge boson, dark matter could be $B'_{udd}\nu'$ or $B'_{ddd}\bar{e}'$ atoms, and constraints from self interactions and the ionized fraction are weakened if $\alpha'_{B-L} \gtrsim \alpha$. Without a massless $B' - L'$ gauge boson, dark matter could be a mixture of $B'_{uud}e'$ atoms and mirror neutrons B'_{udd} ; constraints from self interactions and the ionized fraction are relaxed.

Acknowledgments

The work of L. J. Hall and K. Harigaya was supported in part by the Department of Energy, Office of Science, Office of High Energy Physics, under contract No. DE-AC02-05CH11231, and by the National Science Foundation under grants PHY-1316783 and PHY-1521446. R. Barbieri wants to thank Dr. Max Rössler, the Walter Haefner Foundation and the ETH Zurich Foundation for support. The work of L. J. Hall was performed in part at the Institute for Theoretical Studies ETH and at the Aspen Center for Physics, which is supported by National Science Foundation grant PHY-1066293.

A Heavy axion and Z_2 symmetry breaking in Froggatt Nielsen sector

One of the advantages of the Mirror World is the possibility of a heavy visible QCD axion [65–68]. If the Standard Model and mirror sectors have a common Peccei-Quinn symmetry, the axion mass is given by the mirror QCD dynamics and becomes heavier. The constraint from colliders and astrophysics can be evaded even if the Peccei-Quinn symmetry breaking scale is around the TeV scale. If the physical theta angles of mirror QCD and QCD are identical, $\bar{\theta}' = \bar{\theta}$, the theory still solves the strong CP problem. Due to the small Peccei-Quinn symmetry breaking scale and the large axion mass, the Peccei-Quinn symmetry can be easily understood as an accidental symmetry [65, 68]. (See [69–75] for models of an invisible QCD axion with an accidental Peccei-Quinn symmetry.)

Is it not clear whether this idea can be incorporated into models with Mirror Twin Higgs. As we have shown in this paper, the Z_2 symmetry breaking in the Yukawa couplings is mandatory. This may lead to $\bar{\theta}' \neq \bar{\theta}$, ruining the Peccei-Quinn solution to the strong CP problem [76]. Here we show that if the Z_2 symmetry is broken by VEVs of FN symmetry breaking fields we regain $\bar{\theta}' = \bar{\theta}$. We note that this mechanism is not peculiar to Twin Higgs models, but is generically applicable to Mirror World scenarios.

Let us denote the FN symmetry breaking field as ϕ . The determinants of the mass matrix of $SU(3)_c$ and $SU(3)'_c$ charged fermions are proportional to

$$\det_{SU(3)_c} M \propto \phi^A f(|\phi|), \quad \det_{SU(3)'_c} M' \propto \phi'^A f(|\phi'|), \quad (\text{A.1})$$

where A is the anomaly coefficient of (FN symmetry)- $SU(3)_c$ - $SU(3)_c$, and f is a function. Note that the phases of the mass matrices do not depend on the absolute value of the

VEVs of ϕ and ϕ' . The difference of the FN symmetry breaking scales itself does not ruin the axion solution to the strong CP problem.

If the FN symmetry has no color anomaly (i.e. $A = 0$), the determinants of the mass matrices have the same phases for any phases of the ϕ and ϕ' VEVs, giving $\bar{\theta}' = \bar{\theta}$. If the FN symmetry has a color anomaly, the theta angles may or may not be identical, depending on how the phases of the VEVs of ϕ and ϕ' are determined. For example, suppose that the phase directions of ϕ and ϕ' are determined by explicit breaking of the continuous FN symmetry to a discrete Z_N subgroup, so that vacua are given by

$$\langle\phi\rangle = |\langle\phi\rangle| \times \exp\left(2\pi i \frac{k}{N}\right), \quad \langle\phi'\rangle = |\langle\phi'\rangle| \times \exp\left(2\pi i \frac{k'}{N}\right) \quad (k, k' = 0, 1, \dots, N-1). \quad (\text{A.2})$$

If A/N is an integer, the theta angles remain identical in any vacua. If not, the theta angles remain identical in specific vacua.

Assuming that mirror quark masses are larger than the dynamical scale of mirror QCD, the mass of the QCD axion is given by

$$m_a \simeq m_{\eta'} \left(\frac{\Lambda'_{\text{QCD}}}{\Lambda_{\text{QCD}}}\right)^2 \frac{f_\pi}{f_a} \simeq 1 \text{ MeV} \times \left(\frac{\Lambda'_{\text{QCD}}/\Lambda_{\text{QCD}}}{10}\right)^2 \frac{10 \text{ TeV}}{f_a}. \quad (\text{A.3})$$

Comparing this formula with figures 1 and 2 of [68], constraints from colliders and ΔN_{eff} are satisfied for $f_a \gtrsim 10 \text{ TeV}$ or $f_a \lesssim 100 \text{ GeV}$. For $f_a \gtrsim 10 \text{ TeV}$, however, the QCD axion decays into photons and mirror photons after BBN begins, giving a slight difference between the baryon-to-photon ratio during BBN and during recombination. To conclude whether such a possibility is excluded or not, a calculation of BBN with decaying QCD axions, as well as a calculation of the CMB spectrum including the effect of ΔN_{eff} and non-zero neutrino masses, are required, which is beyond the scope of this paper. For $f_a \lesssim 100 \text{ GeV}$, there should be $SU(3)_c$ particles with masses of $O(100) \text{ GeV}$, which is excluded by hadron colliders.¹²

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] T.D. Lee and C.-N. Yang, *Question of parity conservation in weak interactions*, *Phys. Rev.* **104** (1956) 254 [[INSPIRE](#)].
- [2] I. Yu. Kobzarev, L.B. Okun and I. Ya. Pomeranchuk, *On the possibility of experimental observation of mirror particles*, *Sov. J. Nucl. Phys.* **3** (1966) 837 [*Yad. Fiz.* **3** (1966) 1154] [[INSPIRE](#)].
- [3] Z. Chacko, H.-S. Goh and R. Harnik, *The twin Higgs: natural electroweak breaking from mirror symmetry*, *Phys. Rev. Lett.* **96** (2006) 231802 [[hep-ph/0506256](#)] [[INSPIRE](#)].

¹²If the Peccei-Quinn symmetry is broken by strong dynamics, $SU(3)_c$ charged fermions may be as heavy as $O(1) \text{ TeV}$. However, it is difficult to avoid light $SU(3)_c$ charged pseudo Nambu-Goldstone bosons. If the domain wall number is $O(10)$, all $SU(3)_c$ charged particles can be as heavy as $O(1) \text{ TeV}$.

- [4] R. Barbieri, T. Gregoire and L.J. Hall, *Mirror world at the Large Hadron Collider*, [hep-ph/0509242](#) [INSPIRE].
- [5] S. Chang, L.J. Hall and N. Weiner, *A supersymmetric twin Higgs*, *Phys. Rev. D* **75** (2007) 035009 [[hep-ph/0604076](#)] [INSPIRE].
- [6] N. Craig and K. Howe, *Doubling down on naturalness with a supersymmetric twin Higgs*, *JHEP* **03** (2014) 140 [[arXiv:1312.1341](#)] [INSPIRE].
- [7] P. Batra and Z. Chacko, *A composite twin Higgs model*, *Phys. Rev. D* **79** (2009) 095012 [[arXiv:0811.0394](#)] [INSPIRE].
- [8] M. Geller and O. Telem, *Holographic twin Higgs model*, *Phys. Rev. Lett.* **114** (2015) 191801 [[arXiv:1411.2974](#)] [INSPIRE].
- [9] R. Barbieri, D. Greco, R. Rattazzi and A. Wulzer, *The composite twin Higgs scenario*, *JHEP* **08** (2015) 161 [[arXiv:1501.07803](#)] [INSPIRE].
- [10] M. Low, A. Tesi and L.-T. Wang, *Twin Higgs mechanism and a composite Higgs boson*, *Phys. Rev. D* **91** (2015) 095012 [[arXiv:1501.07890](#)] [INSPIRE].
- [11] D. Buttazzo, F. Sala and A. Tesi, *Singlet-like Higgs bosons at present and future colliders*, *JHEP* **11** (2015) 158 [[arXiv:1505.05488](#)] [INSPIRE].
- [12] N. Craig, A. Katz, M. Strassler and R. Sundrum, *Naturalness in the dark at the LHC*, *JHEP* **07** (2015) 105 [[arXiv:1501.05310](#)] [INSPIRE].
- [13] ATLAS and CMS collaborations, *Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **08** (2016) 045 [[arXiv:1606.02266](#)] [INSPIRE].
- [14] A. Manohar and H. Georgi, *Chiral quarks and the nonrelativistic quark model*, *Nucl. Phys. B* **234** (1984) 189 [INSPIRE].
- [15] M.A. Luty, *Naive dimensional analysis and supersymmetry*, *Phys. Rev. D* **57** (1998) 1531 [[hep-ph/9706235](#)] [INSPIRE].
- [16] G. 't Hooft, *A planar diagram theory for strong interactions*, *Nucl. Phys. B* **72** (1974) 461 [INSPIRE].
- [17] S. Borsányi et al., *Calculation of the axion mass based on high-temperature lattice quantum chromodynamics*, *Nature* **539** (2016) 69 [[arXiv:1606.07494](#)] [INSPIRE].
- [18] T. Yanagida, *Horizontal symmetry and masses of neutrinos*, *Conf. Proc.* **C 7902131** (1979) 95 [INSPIRE].
- [19] M. Gell-Mann, P. Ramond and R. Slansky, *Complex spinors and unified theories*, *Conf. Proc.* **C 790927** (1979) 315 [[arXiv:1306.4669](#)] [INSPIRE].
- [20] P. Minkowski, *$\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays?*, *Phys. Lett. B* **67** (1977) 421 [INSPIRE].
- [21] PLANCK collaboration, P.A.R. Ade et al., *Planck 2015 results. XIII. Cosmological parameters*, *Astron. Astrophys.* **594** (2016) A13 [[arXiv:1502.01589](#)] [INSPIRE].
- [22] C.D. Froggatt and H.B. Nielsen, *Hierarchy of quark masses, Cabibbo angles and CP-violation*, *Nucl. Phys. B* **147** (1979) 277 [INSPIRE].
- [23] ATLAS collaboration, *Prospects for new physics in Higgs couplings studies with the ATLAS detector at the HL-LHC*, [ATL-PHYS-PUB-2014-017](#), CERN, Geneva Switzerland (2014).

- [24] D.M. Asner et al., *ILC Higgs white paper*, [arXiv:1310.0763](#) [INSPIRE].
- [25] CP-PACS collaboration, M. Okamoto et al., *Equation of state for pure SU(3) gauge theory with renormalization group improved action*, *Phys. Rev. D* **60** (1999) 094510 [[hep-lat/9905005](#)] [INSPIRE].
- [26] S. Borsányi, G. Endrodi, Z. Fodor, S.D. Katz and K.K. Szabo, *Precision SU(3) lattice thermodynamics for a large temperature range*, *JHEP* **07** (2012) 056 [[arXiv:1204.6184](#)] [INSPIRE].
- [27] CMS collaboration, *Projected performance of an upgraded CMS detector at the LHC and HL-LHC: contribution to the Snowmass process*, [arXiv:1307.7135](#) [INSPIRE].
- [28] S. Davidson, B. Campbell and D.C. Bailey, *Limits on particles of small electric charge*, *Phys. Rev. D* **43** (1991) 2314 [INSPIRE].
- [29] J. Jaeckel, M. Jankowiak and M. Spannowsky, *LHC probes the hidden sector*, *Phys. Dark Univ.* **2** (2013) 111 [[arXiv:1212.3620](#)] [INSPIRE].
- [30] A. Haas, C.S. Hill, E. Izaguirre and I. Yavin, *Looking for milli-charged particles with a new experiment at the LHC*, *Phys. Lett. B* **746** (2015) 117 [[arXiv:1410.6816](#)] [INSPIRE].
- [31] H. Goldberg and L.J. Hall, *A new candidate for dark matter*, *Phys. Lett. B* **174** (1986) 151 [INSPIRE].
- [32] S.I. Blinnikov and M. Yu. Khlopov, *On possible effects of ‘mirror’ particles*, *Sov. J. Nucl. Phys.* **36** (1982) 472 [*Yad. Fiz.* **36** (1982) 809] [INSPIRE].
- [33] S.I. Blinnikov and M. Khlopov, *Possible astronomical effects of mirror particles*, *Sov. Astron.* **27** (1983) 371 [*Astron. Zh.* **60** (1983) 632] [INSPIRE].
- [34] E.W. Kolb, D. Seckel and M.S. Turner, *The shadow world*, *Nature* **314** (1985) 415 [INSPIRE].
- [35] A. Sommerfeld, *Über die Beugung und Bremsung der Elektronen* (in German), *Annalen Phys.* **403** (1931) 257.
- [36] J. Hisano, S. Matsumoto and M.M. Nojiri, *Unitarity and higher order corrections in neutralino dark matter annihilation into two photons*, *Phys. Rev. D* **67** (2003) 075014 [[hep-ph/0212022](#)] [INSPIRE].
- [37] J.M. Cline, Z. Liu, G. Moore and W. Xue, *Composite strongly interacting dark matter*, *Phys. Rev. D* **90** (2014) 015023 [[arXiv:1312.3325](#)] [INSPIRE].
- [38] J.M. Cline, Z. Liu, G. Moore and W. Xue, *Scattering properties of dark atoms and molecules*, *Phys. Rev. D* **89** (2014) 043514 [[arXiv:1311.6468](#)] [INSPIRE].
- [39] M. Kaplinghat, S. Tulin and H.-B. Yu, *Dark matter halos as particle colliders: unified solution to small-scale structure puzzles from dwarfs to clusters*, *Phys. Rev. Lett.* **116** (2016) 041302 [[arXiv:1508.03339](#)] [INSPIRE].
- [40] F.-Y. Cyr-Racine and K. Sigurdson, *Cosmology of atomic dark matter*, *Phys. Rev. D* **87** (2013) 103515 [[arXiv:1209.5752](#)] [INSPIRE].
- [41] S.L. Dubovsky and D.S. Gorbunov, *Small second acoustic peak from interacting cold dark matter?*, *Phys. Rev. D* **64** (2001) 123503 [[astro-ph/0103122](#)] [INSPIRE].
- [42] S.L. Dubovsky, D.S. Gorbunov and G.I. Rubtsov, *Narrowing the window for millicharged particles by CMB anisotropy*, *JETP Lett.* **79** (2004) 1 [*Pisma Zh. Eksp. Teor. Fiz.* **79** (2004) 3] [[hep-ph/0311189](#)] [INSPIRE].

- [43] A.D. Dolgov, S.L. Dubovsky, G.I. Rubtsov and I.I. Tkachev, *Constraints on millicharged particles from Planck data*, *Phys. Rev. D* **88** (2013) 117701 [[arXiv:1310.2376](#)] [[INSPIRE](#)].
- [44] L. Chuzhoy and E.W. Kolb, *Reopening the window on charged dark matter*, *JCAP* **07** (2009) 014 [[arXiv:0809.0436](#)] [[INSPIRE](#)].
- [45] N. Craig and A. Katz, *The fraternal WIMP miracle*, *JCAP* **10** (2015) 054 [[arXiv:1505.07113](#)] [[INSPIRE](#)].
- [46] I. García García, R. Lasenby and J. March-Russell, *Twin Higgs asymmetric dark matter*, *Phys. Rev. Lett.* **115** (2015) 121801 [[arXiv:1505.07410](#)] [[INSPIRE](#)].
- [47] M. Farina, *Asymmetric twin dark matter*, *JCAP* **11** (2015) 017 [[arXiv:1506.03520](#)] [[INSPIRE](#)].
- [48] A. Djouadi and M. Drees, *QCD corrections to neutralino nucleon scattering*, *Phys. Lett. B* **484** (2000) 183 [[hep-ph/0004205](#)] [[INSPIRE](#)].
- [49] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Remarks on Higgs boson interactions with nucleons*, *Phys. Lett. B* **78** (1978) 443 [[INSPIRE](#)].
- [50] ETM collaboration, A. Abdel-Rehim et al., *Direct evaluation of the quark content of nucleons from lattice QCD at the physical point*, *Phys. Rev. Lett.* **116** (2016) 252001 [[arXiv:1601.01624](#)] [[INSPIRE](#)].
- [51] J.M. Alarcon, J. Martin Camalich and J.A. Oller, *The chiral representation of the πN scattering amplitude and the pion-nucleon sigma term*, *Phys. Rev. D* **85** (2012) 051503 [[arXiv:1110.3797](#)] [[INSPIRE](#)].
- [52] J.M. Alarcon, L.S. Geng, J. Martin Camalich and J.A. Oller, *The strangeness content of the nucleon from effective field theory and phenomenology*, *Phys. Lett. B* **730** (2014) 342 [[arXiv:1209.2870](#)] [[INSPIRE](#)].
- [53] D.S. Akerib et al., *Results from a search for dark matter in the complete LUX exposure*, [arXiv:1608.07648](#) [[INSPIRE](#)].
- [54] PANDAX-II collaboration, A. Tan et al., *Dark matter results from first 98.7 days of data from the PandaX-II experiment*, *Phys. Rev. Lett.* **117** (2016) 121303 [[arXiv:1607.07400](#)] [[INSPIRE](#)].
- [55] XENON collaboration, E. Aprile et al., *Physics reach of the XENON1T dark matter experiment*, *JCAP* **04** (2016) 027 [[arXiv:1512.07501](#)] [[INSPIRE](#)].
- [56] LZ collaboration, D.S. Akerib et al., *LUX-ZEPLIN (LZ) conceptual design report*, [arXiv:1509.02910](#) [[INSPIRE](#)].
- [57] J. Billard, L. Strigari and E. Figueroa-Feliciano, *Implication of neutrino backgrounds on the reach of next generation dark matter direct detection experiments*, *Phys. Rev. D* **89** (2014) 023524 [[arXiv:1307.5458](#)] [[INSPIRE](#)].
- [58] E. Del Nobile, G.B. Gelmini, P. Gondolo and J.-H. Huh, *Direct detection of light anapole and magnetic dipole DM*, *JCAP* **06** (2014) 002 [[arXiv:1401.4508](#)] [[INSPIRE](#)].
- [59] CRESST collaboration, G. Angloher et al., *Probing low WIMP masses with the next generation of CRESST detector*, [arXiv:1503.08065](#) [[INSPIRE](#)].
- [60] SUPERCDMS collaboration, R. Calkins, *The SuperCDMS Soudan high threshold WIMP search and the planned SuperCDMS SNOLAB experiment*, *J. Phys. Conf. Ser.* **718** (2016) 042009 [[INSPIRE](#)].

- [61] J.M. Cline, Z. Liu and W. Xue, *Millicharged atomic dark matter*, *Phys. Rev. D* **85** (2012) 101302 [[arXiv:1201.4858](#)] [[INSPIRE](#)].
- [62] CRESST collaboration, G. Angloher et al., *Results on light dark matter particles with a low-threshold CRESST-II detector*, *Eur. Phys. J. C* **76** (2016) 25 [[arXiv:1509.01515](#)] [[INSPIRE](#)].
- [63] I. Affleck and M. Dine, *A new mechanism for baryogenesis*, *Nucl. Phys. B* **249** (1985) 361 [[INSPIRE](#)].
- [64] M. Dine, L. Randall and S.D. Thomas, *Baryogenesis from flat directions of the supersymmetric Standard Model*, *Nucl. Phys. B* **458** (1996) 291 [[hep-ph/9507453](#)] [[INSPIRE](#)].
- [65] V.A. Rubakov, *Grand unification and heavy axion*, *JETP Lett.* **65** (1997) 621 [[hep-ph/9703409](#)] [[INSPIRE](#)].
- [66] Z. Berezhiani, L. Gianfagna and M. Giannotti, *Strong CP problem and mirror world: the Weinberg-Wilczek axion revisited*, *Phys. Lett. B* **500** (2001) 286 [[hep-ph/0009290](#)] [[INSPIRE](#)].
- [67] A. Hook, *Anomalous solutions to the strong CP problem*, *Phys. Rev. Lett.* **114** (2015) 141801 [[arXiv:1411.3325](#)] [[INSPIRE](#)].
- [68] H. Fukuda, K. Harigaya, M. Ibe and T.T. Yanagida, *Model of visible QCD axion*, *Phys. Rev. D* **92** (2015) 015021 [[arXiv:1504.06084](#)] [[INSPIRE](#)].
- [69] S.M. Barr and D. Seckel, *Planck scale corrections to axion models*, *Phys. Rev. D* **46** (1992) 539 [[INSPIRE](#)].
- [70] R. Holman, S.D.H. Hsu, T.W. Kephart, E.W. Kolb, R. Watkins and L.M. Widrow, *Solutions to the strong CP problem in a world with gravity*, *Phys. Lett. B* **282** (1992) 132 [[hep-ph/9203206](#)] [[INSPIRE](#)].
- [71] M. Dine, *Problems of naturalness: some lessons from string theory*, [hep-th/9207045](#) [[INSPIRE](#)].
- [72] A.G. Dias, V. Pleitez and M.D. Tonasse, *Naturally light invisible axion in models with large local discrete symmetries*, *Phys. Rev. D* **67** (2003) 095008 [[hep-ph/0211107](#)] [[INSPIRE](#)].
- [73] L.M. Carpenter, M. Dine and G. Festuccia, *Dynamics of the Peccei Quinn scale*, *Phys. Rev. D* **80** (2009) 125017 [[arXiv:0906.1273](#)] [[INSPIRE](#)].
- [74] K. Harigaya, M. Ibe, K. Schmitz and T.T. Yanagida, *Peccei-Quinn symmetry from a gauged discrete R symmetry*, *Phys. Rev. D* **88** (2013) 075022 [[arXiv:1308.1227](#)] [[INSPIRE](#)].
- [75] K. Harigaya, M. Ibe, K. Schmitz and T.T. Yanagida, *Peccei-Quinn symmetry from dynamical supersymmetry breaking*, *Phys. Rev. D* **92** (2015) 075003 [[arXiv:1505.07388](#)] [[INSPIRE](#)].
- [76] A. Albaid, M. Dine and P. Draper, *Strong CP and SUZ₂*, *JHEP* **12** (2015) 046 [[arXiv:1510.03392](#)] [[INSPIRE](#)].