

What is the importance of islands to environmental conservation?

Review Article**Author(s):**

Kueffer, Christoph; Kinney, Kealohanuiopuna

Publication date:

2017-12

Permanent link:

<https://doi.org/10.3929/ethz-b-000220360>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Environmental Conservation 44(4), <https://doi.org/10.1017/S0376892917000479>

What is the importance of islands to environmental conservation?

THEMATIC SECTION

Humans and Island
EnvironmentsCHRISTOPH KUEFFER*¹ AND KEALOHANUIOPUNA KINNEY²¹*Institute of Integrative Biology, ETH Zurich, Universitätsstrasse 16, CH-8092 Zurich, Switzerland and*²*Institute of Pacific Islands Forestry, US Forest Service, 60 Nowelo St. Hilo, HI, USA*

Date submitted: 15 May 2017; Date accepted: 8 August 2017

SUMMARY

This article discusses four features of islands that make them places of special importance to environmental conservation. First, investment in island conservation is both urgent and cost-effective. Islands are threatened hotspots of diversity that concentrate unique cultural, biological and geophysical values, and they form the basis of the livelihoods of millions of islanders. Second, islands are paradigmatic places of human–environment relationships. Island livelihoods have a long tradition of existing within spatial, ecological and ultimately social boundaries and are still often highly dependent on local resources and social cohesion. Island cultures and their rich biocultural knowledge can be an important basis for revitalizing and innovating sustainable human–nature relationships. Third, islands form a global web that interlinks biogeographic regions and cultural spaces. They are nodes in a global cultural network: as multicultural island societies, through diaspora islander communities on continents and through numerous political and trade relationships among islands and between islands and countries on continents. Fourth, islands can serve as real-world laboratories that enable scientific innovation, integration of local and generalized knowledge and social learning and empowerment of local actors. We conclude that island systems can serve as globally distributed hubs of innovation, if the voices of islanders are better recognized.

Keywords: oceanic island, biocultural, biodiversity, sustainability, model system, networks, conservation, heritage, social innovation, insularity

INTRODUCTION

This article gives a broad overview of the relevance of islands to environmental conservation from an interdisciplinary perspective. We consider only islands in the ocean; not islands in freshwater bodies or habitat islands. Among these

islands of the world's oceans, we cover both islands close to continents and others isolated far out in the oceans, and the full range from small to very large islands. Small and isolated islands represent unique cultural and biological values and the environmental challenges of insularity in its most pronounced form. However, as we will demonstrate, all islands and island people share enough concerns to consider them together (Baldacchino 2007; Royle 2008; Gillespie & Clague 2009; Baldacchino & Niles 2011; Royle 2014).

Islands are hotspots of cultural, biological and geophysical diversity, and as such they form the basis of the livelihoods of millions of islanders (Menard 1986; Nunn 1994; Royle 2008; Gillespie & Clague 2009; Royle 2014; Kueffer *et al.* 2016). In total, 43 out of the world's 195 countries are islands or archipelagos, and over two-thirds include islands (global island database, http://www.globalislands.net/about/gid_functions.php, last accessed 3 April 2017). Today, islands are amongst the most vulnerable places on Earth threatened by both environmental and socioeconomic challenges (Baldacchino 2007; Caujapé-Castells *et al.* 2010; Baldacchino & Niles 2011; Walker & Bellingham 2011; Connell 2013). These threats include climate change (and especially sea-level rise), unsustainable use of local resources, overpopulation, the end of cheap energy with its consequences for global travel and transport on which many islands depend (e.g. tourism and import of many goods including food) and exposure of small-island economies to the vagaries of an unpredictable global economy (Baldacchino 2007; Baldacchino & Niles 2011; Walker & Bellingham 2011; Connell 2013; Royle 2014).

Given the high concentration of biological and human values per land area and the high threat level, it might be expected that islands are listed amongst the highest priorities of international environmental conservation. However, analyses of global spending show that island conservation efforts are underfunded relative to continents (e.g. Waldron *et al.* 2013), and island people have struggled to make their voices heard in international policy processes (Dahl 2017). The vast majority of endemic and threatened species in countries such as France, the UK or the USA are situated on islands associated with these countries, but conservation spending and institutional capacity on these islands are very small compared to the respective mainland (e.g. Leonard 2008). This might have to do with the often weak power of islanders (Royle 2014). The continued

*Correspondence: Prof Dr Christoph Kueffer email: kueffer@env.ethz.ch




Figure 1 Islands are important to environmental conservation for at least four interconnected reasons: (i) they are global hotspots of cultural, biological and geophysical diversity and uniqueness; (ii) they are paradigmatic places of human–environment relationships; (iii) they form a global cultural and ecological web; and (iv) they serve as laboratories for scientific and social innovation.

dependence on former colonial powers is much higher among islands than on continents; most islands in the Arctic region, Mediterranean Sea and Atlantic Ocean and many islands in the Central and Eastern Pacific are part of or dependencies of continental countries, and there also remain many continental dependencies in the Caribbean Sea and Indian Ocean. Islands that gained independence are often small, highly dependent on other countries and exposed to fluctuations and shifts in the world economy. Islands might also be overlooked due to their small total area (5–6% of Earth's land area) and human population size (about 10% of the world's population).

In this article, we argue that islands are fundamental to international environmental conservation, despite their small total land area and population size, for at least four interconnected reasons. First, as hotspots of biological and geophysical diversity and uniqueness, the world's islands together represent a major proportion of the natural riches on the Earth. Second, islands are paradigmatic places of human–environment relationships. Island cultures around the world hold rich cultural knowledge and experience about living within spatial, ecological and ultimately social boundaries. The corresponding biocultural knowledge is valuable in its own right as an important compound of cultural heritage and contemporary human life on Earth, and it is also a pivotal resource for revitalizing and innovating sustainable human–nature relationships on islands and beyond. Third, given their global distribution across all geographic regions and from the highest latitudes to the equator, islands sample and represent the ecological and cultural diversity of the planet comprehensively and form a global web that interlinks biogeographic regions and cultural spaces. Fourth, islands can serve as real-world laboratories that enable innovation in both the sciences and culture. We review in this article each of these four interconnected dimensions that together contribute to the unique importance of islands to environmental conservation (Fig. 1).

HOTSPOTS OF GEOPHYSICAL AND BIOLOGICAL DIVERSITY AND UNIQUENESS

Islands are widely recognized as hotspots of geophysical and biological diversity and uniqueness (Fig. 1). A third of all identified global biodiversity hotspots consist mostly or entirely of islands (Zachos & Habel 2011): several Pacific Ocean regions, several island groups in Southeast Asia, the Mediterranean Basin, the Caribbean islands and Madagascar and Indian Ocean islands. Several other hotspot regions include islands (e.g. Indo-Burma, Western Ghats and Sri Lanka and the Guinean forests of Western Africa). Equally, many UNESCO World Heritage sites are situated on islands, including, among others, many last wilderness areas such as Aldabra Atoll, Macquarie Island or Phoenix Islands, but also many sites are on highly populated islands. Some sites are mainly protected for their biodiversity, while others are protected for their geophysical uniqueness (e.g. the volcanic landscapes of La Réunion, Tenerife, the Aeolian Islands, Surtsey and Hawaii).

In addition to terrestrial biodiversity, islands are also important as coastal areas for marine biodiversity. Islands are characterized by a particularly high ratio of coastline to land area. Therefore, the total island coastline is disproportionately higher than is suggested by their global land area. In addition, even small islands have large, exclusive economic zones with huge territorial claims to surrounding oceans. Island people are stewards for a sixth of the Earth's surface. As a result, more than half of the world's marine biodiversity and seven of the world's ten coral reef hotspots belong to island territories (CBD 2014). The marine side of islands has important implications for the global ecosystem. While terrestrial island ecosystems have, due to their small overall size, a negligible effect on global Earth system processes such as element cycles or climate, islands are central to the global marine ecosystem as widely scattered coastal areas that serve as breeding and feeding sites for marine animals (including coral reefs) and as terrestrial nesting sites for seabirds and marine mammals (Mulder *et al.* 2011). Because the high productivity of marine ecosystems is largely confined to coastal areas, and nesting sites (rather than food availability) can be limiting for marine animals, islands have an ecological influence on oceans that is disproportionately higher than is suggested by their land area.

Marine island ecosystems are biodiversity hotspots because many of them harbour a very high density of species per area. In contrast, in terrestrial island ecosystems, the species number per area is low (Kier *et al.* 2009) due to their isolation from the species pools of other land areas. Instead, what makes terrestrial island biodiversity valuable is its uniqueness and great variability: across the world's islands, between islands within archipelagos and islands groups and within islands (Fig. 1). In this sense, islands are paradigmatic cases of the conservation policy concept of biodiversity hotspots. They concentrate a broad sample of unique and diverse natural features into small land areas. Islands can be considered living museums of natural heritage.

Diversity across the world's islands

The islands of the world's oceans are very diverse in their abiotic characteristics (Menard 1986; Nunn 1994; Gillespie & Clague 2009). Islands occur in all oceanic regions and at all latitudes, their areas and topographies vary greatly, and their geological origins are diverse. There have been attempts to assemble global databases of physical characteristics of islands such as climate, size and isolation (Weigelt *et al.* 2013; Weigelt & Krefl 2013).

Depending on definition, the smallest islands have a size of only a few square metres (i.e. a single rock), while the largest landmasses that are typically considered islands – Greenland and New Guinea – have areas of over 2 million and almost 800 000 km², respectively (i.e. if Australia is classified as a continent). Together, islands in the ocean make up *c.* 5–6% of total land area (excluding mostly ice-covered Antarctica and Greenland). There are at least 20 000 islands >1 km² in area, close to 0.5 million >1 ha in area and there are probably billions of islands if all sizes are considered (Royle 2014). More specifically, 18 islands are larger than 100 000 km² (e.g. New Guinea, Borneo, Madagascar, Baffin and Great Britain), another 53 are larger than 10 000 km² and *c.* 1500 are between 100 and 10 000 km² (Depraetere & Dahl 2007). Equally diverse is the topography of islands. Low-lying islands such as atolls reach only a few metres above sea level (m asl), while the tallest ones are major mountains that can be higher than 4000 m asl; in the subtropics and tropics, 14 islands reach the alpine zone with an elevation higher than 2750 m asl (Juvik *et al.* 2014).

Although isolation by a surrounding ocean is the defining characteristic that is common to all islands in the ocean, this factor also varies greatly. Some land-bridge islands (continental peninsulas that, due to interglacial sea-level rise, lost their connection to continents) are isolated by less than 1 km from the nearest continental landmass, while the most isolated islands are at a distance of over 5000 km from any continent (e.g. different islands in French Polynesia) (Weigelt *et al.* 2013). The geography of islands has changed repeatedly and strongly over geological timescales due to continental shift, ontogeny of islands and sea-level changes (e.g. Fernández-Palacios *et al.* 2016). Because of sea-level change, some islands that are isolated in interglacial times (high sea level) are connected during glacial times (low sea level). Examples are the central island group of the Hawaiian archipelago around Maui, many islands in Southeast Asia or the Seychelles granitic islands (now *c.* 40 islands of a total land area of *c.* 235 km², they formed a micro-continent of more than 40 000 km² during the last glacial maximum only some 20 000 years ago).

A fourth major factor that contributes to the high abiotic diversity of islands is their diverse geological origin (Menard 1986; Nunn 1994; Juvik & Juvik 1998; Gillespie & Clague 2009; Kueffer *et al.* 2016). Oceanic islands in a strict sense are volcanic islands that formed through the accumulation of submarine magma and were never connected to continents. However, not all islands are of volcanic origin. Land-

bridge islands and other islands on a continental shelf, as well as continental fragments or micro-continent (originally continental areas, but now isolated in the ocean through continental drift; e.g. Madagascar or the Seychelles), are built on a continental underground and consequently are characterized by a different geology (e.g. granites). While volcanic islands reach a maximal age of *c.* 5–10 million years, continental ones can be much older. The Seychelles granitic islands for instance have been isolated from any other landmass for *c.* 65 million years, and they are composed of rocks that are several 100 million years old.

The world's islands equally demonstrate high biodiversity. Islands of a given area tend to support fewer species than continental regions of the same area due to their isolation and small total land area. However, the world's islands together make a disproportionately large contribution to the world's total biodiversity due to their high level of endemism (i.e. a high proportion of species on a particular island – or in an island group – occur naturally nowhere else on Earth, neither on a different island nor on a continent). For instance, between 20% and 25% of the world's bird and plant species are endemic to islands and occur nowhere else (Johnson & Stattersfield 1990; Caujapé-Castells *et al.* 2010), and very high numbers of island endemics are found in many other taxonomic groups, including, for instance, snails (Chiba & Cowie 2016). The biodiversity value of islands is, however, not fully captured only by the statistics of species numbers (Gillespie & Clague 2009; Kueffer *et al.* 2016). Islands are also important because they comprehensively represent the biogeography and climate zones of the world, and therefore demonstrate a high diversity of different phylogenetic lineages from all continents (Weigelt *et al.* 2015).

Further, islands are showcases of evolutionary processes. Evolutionary adaptation to island life has led to some of the most unique life forms. One general adaptation on islands is that animals and plants tend to lose their dispersability. Consequently, there are many examples of flightless birds and insects on islands, as well as plants with reduced dispersability, including the plant with the largest seed – the palm *Lodoicea maldivica* from the Seychelles (Edwards *et al.* 2015). Also sizes of island organisms change: small species tend to become bigger and large ones smaller (Lomolino 2005). Insular dwarfism has, for instance, been documented for dinosaurs, mammoths, elephants and humans, and insular gigantism has been documented for rats, lagomorphs, lemurs, tortoises and birds (e.g. the moa and dodo). Other recurrent evolutionary adaptations to island life include woodiness of plant groups that are herbaceous on continents, diminution of clutch size in vertebrates, diminution of defensive behaviour in animals and plants and development of sexual dimorphism in plants. Because island floras and faunas are small, species that are typical for occupying a particular ecological niche are often missing. As a result, ecological functions on islands are often taken up by unusual taxonomic groups; for instance, giant tortoises as large grazers, Komodo dragons as top predators or

widespread pollination by lizards (Gillespie & Clague 2009). In a certain sense, island evolution widened the range of species that can maintain a certain ecological function.

Diversity within archipelagos and within islands

Islands tend to be clustered by forming chains, groups and regional assemblies (Depraetere & Dahl 2007). Geophysical diversity is often high within island groups, partly because they are composed of different types of islands (e.g. volcanic ones and others of continental origin) and especially because they include volcanic islands of different ages. In the first phase of their ontogeny, characterized by volcanic activity, volcanic islands grow to tall mountains that, at their climax, typically reach over 4000 m above sea level and up to 10 000 m from the seabed (Mauna Kea in the Hawaiian archipelago) (Juvik & Juvik 1998). Once volcanic activity stops, the work of erosion becomes increasingly visible and shapes island landforms. Old volcanic islands are eventually characterized by deep valleys, rugged topography and steep slopes carved by erosion. Ultimately, a volcanic island is completely eroded and remains only as a flat island of an elevation of a few metres – and in the case of a tropical island, it may be surrounded by a coral reef. Former tropical volcanic island in these last stages are called atolls. Because volcanic activity follows lines of tectonic rupture, volcanic islands and atolls occur in chains in a predictable way, with, at one end, new islands forming through volcanic activity still under the sea, followed by the highest and youngest islands, and finally, at the other end, the oldest volcanic islands and atolls. The Hawaiian archipelago, for instance, stretches from the southeast at 155°W, with the youngest island, Big Island, reaching over 4000 m asl and still with active volcanism, to the low-lying atolls of the northwestern Hawaiian islands at 180°W. Southeast of the Big Island, the next island (Loihi) forms as a growing volcano below sea level. Northwest of the northwestern Hawaiian Islands, the submersed former islands (guyots) of the Emperor Guyots chain reach almost to the Kamchatka peninsula in the northeast of Eurasia, which indicates the former extent of the archipelago. Soil conditions change with island age from nitrogen limitation on the youngest islands to phosphorus limitation on the oldest ones (Vitousek 2004).

These geophysical forces working over the lifetime of a volcanic island produce high topographic diversity on a particular island. Older islands are characterized by a highly rugged topography with steep slopes. Younger, tall islands often have highly contrasting climates on the rain-fed windward and the dry leeward sides in the shade of the mountain peak(s). Annual rainfall can vary by a factor of 10–20 over a distance of typically less than 100 km. Such topographic and associated climatic heterogeneity leads to very high habitat diversity, so that on a single (sub)tropical island, amongst the driest, most humid and coldest tropical habitats can be present – with their associated specialized floras and faunas, and often also specialized traditional or modern land-use systems

(Vitousek *et al.* 2004; Rolett 2008). In some ways, islands are representations of whole continents at a micro-scale.

Such high geophysical variation – and associated climate, soil and habitat variation – within archipelagos and within islands is a primary driver of the production of island biodiversity *in situ* (i.e. of speciation on islands). Indeed, particularly spectacular features of island biodiversity are adaptive radiations: macro-evolutionary processes that produce many different and ecologically diverse species from a single ancestor species (Gillespie & Clague 2009). Species that were formed through adaptive radiation contribute a disproportionately high share to the total species richness on islands. Such evolutionary radiations can unfold along island chains over time periods longer than the life of a single island (Gillespie & Clague 2009).

Threatened island hotspots

Hotspots of natural heritage on islands are highly threatened. In the case of geophysical features, housing and tourism development particularly threaten the integrity of unique landscape features, as exemplified by the struggle against the further development of astronomical infrastructure in the summit area of Mauna Kea volcano (Hawaii archipelago), which is sacred to native Hawaiians.

Biodiversity loss is particularly severe on islands (Grenyer *et al.* 2006; Whittaker & Fernández-Palacios 2007; Gillespie & Clague 2009; Caujapé-Castells *et al.* 2010; Chiba & Cowie 2016). The majority of documented species extinctions to date are from islands, including such prominent cases as the dodo and other giant birds, most honeycreeper species in Hawaii, many other birds, dwarf mammals from Mediterranean islands, many snail species across the Pacific and many plant species (e.g. Juvik & Juvik 1998; Whittaker & Fernández-Palacios 2007; Gillespie & Clague 2009; Caujapé-Castells *et al.* 2010; Alcover *et al.* 2015; Chiba & Cowie 2016). In addition, the proportion of threatened species is also particularly high on islands. In the Hawaiian Islands, for instance, most of the remaining honeycreeper species might be extinct within the next few decades (Paxton *et al.* 2016).

In addition to past threats, new ones are emerging. New building development triggered by population growth and economic development – including tourism – encroaches on natural areas and increases pollution levels (especially in coastal areas). Climate change will have direct negative impacts on island biodiversity in particular because habitats are often very small and highly fragmented (both naturally and anthropogenically), leaving not much space for species to track their climate to new areas (Harter *et al.* 2015). Climate change might worsen other threats (e.g. biological invasions spreading to higher elevations or increased fire risk in already highly fragmented lowland dry habitats). Major impacts can also be expected in marine ecosystems, especially on coral reefs. Besides direct impacts, indirect impacts will be important, especially when human populations are displaced by sea-level rise from coastal areas and might settle in the

as-yet less developed interiors of islands (Wetzel *et al.* 2012). Lastly, many populations of island species are already too small to survive over the long term without the addition of further threats; a phenomenon called ‘extinction debt’ (Triantis *et al.* 2010). Kueffer and Kaiser-Bunbury (2014), for instance, report the following examples: 44.2% of the Hawaiian flora is restricted to 20 or fewer populations in the wild, and more than 100 species are known to have 20 or fewer remaining individuals. In Haiti, less than 4% of the land area is covered with forest, and no primary forest remains. The flora of Haiti is composed of more than 5000 vascular plants with an endemism of about a third, as well as more than 2000 animal species (most of them originally forest species). In New Caledonia, less than 2% of lowland dry vegetation remains intact, mostly as patches of less than 5 ha, the largest one being 200 ha. Such a reduction and fragmentation of remaining habitat will have major consequences not only for well-documented organism groups such as plants or birds, but also for invertebrates, for which data are often lacking (e.g. beetles in the Azores; Terzopoulou *et al.* 2015).

PARADIGMATIC PLACES OF HUMAN–ENVIRONMENT RELATIONSHIPS

Islands can be considered paradigmatic places of human–environment relationships (Fig. 1). Island societies often build on long traditions of living within spatial, ecological and ultimately social boundaries and are still often highly dependent on local resources and social cohesion. In addition, islanders had to learn how to deal with rapid and major environmental and socioeconomic transformations that often had severe consequences for biodiversity, the environment and livelihoods. At present, many environmental problems are particularly acute and tangible on islands, and in this sense islands are at the forefront of contemporary environmental conservation (e.g. Kelman 2010).

Human history on islands includes some of the most impressive human achievements. In the Pacific, for instance, the greatest marine migration in human history happened, being one of the most impressive examples of human determination, creativity and adaptability (Gillespie & Clague 2009; Low 2013). Much later, the mutineers of HMS *Bounty* found their way to the tiny and very isolated island of Pitcairn, where they established a colony, adapting their European knowledge to the new challenges and apparently profiting from plants that remained on the island from earlier Polynesian settlements (e.g. breadfruits and sweet potatoes) (Gillespie & Clague 2009). Similar stories of human exploration and adaptation happened in the Arctic, or in the other oceans.

Upon arrival, humans had to adapt to a life on a small and isolated piece of land, often characterized by limited resources. Such a recurrent experiment of human adaptation happened in very different climates and geographic zones and on very diverse islands, including atolls and young to old volcanic islands with contrasting topographies and soil

conditions. On high tropical islands, they had to find different agricultural solutions for the very wet windward compared with the very dry leeward sites (Vitousek *et al.* 2004). In some cases, the island’s terrestrial flora and fauna provided a rich food source (e.g. in the Mediterranean Basin or on Southeast Asian islands), while in other cases, humans depended mostly on the few plants and animals that they carried with them on their journeys across the oceans, plus marine resources (e.g. on many islands in the Pacific) (Juvik & Juvik 1998; Gillespie & Clague 2009; Whistler 2009). Human colonization was conducted by very different cultural groups, and at very different times, sometimes in prehistory, while in other cases not before the colonial expansion of Europeans. Many of these human explorations of island life have in common that people depended on scarce resources, small land areas, few opportunities to migrate and establish a new life elsewhere (as individuals or groups), a life at the interface of the sea and the land and limited exchange with other societies on other islands or continents (Fitzpatrick & Keegan 2007; Rolett 2008; Royle 2014). On some islands, people had to adapt to a harsh climate (e.g. at the highest latitudes in the Arctic or on dry islands such as in the Socotra archipelago in the Indian Ocean).

Indigenous people or early colonial settlers had to invent long-term subsistence systems based on local resources and restricted land (Rolett 2008; Gillespie & Clague 2009; Whistler 2009; Koshiha *et al.* 2014; McMillen *et al.* 2014; Royle 2014; Braje *et al.* 2017). Practices involved, for instance, terracing and stonewall enclosures, polycultural cultivation and fallowing, water management, agriculture and agroforestry adapted to local conditions, aquaculture and sustainable harvesting practices of wild plants and animals. These diverse practices were and still often are embedded in local knowledge systems, institutional arrangements, customs and mythologies. Initial land use on newly settled islands typically led to substantially transformed coastal and lowland anthropogenic landscapes and different degrees of exploitation of interior habitats (Juvik & Juvik 1998; Rolett & Diamond 2004; Fitzpatrick & Keegan 2007; De Nascimento *et al.* 2009; Gillespie & Clague 2009; Braje *et al.* 2017). While some islands had to be abandoned at some point due to overexploitation, on other islands, traditional land-use systems persisted for thousands of years (Rolett & Diamond 2004; Fitzpatrick & Keegan 2007; Rolett 2008; Gillespie & Clague 2009; Koshiha *et al.* 2014; Braje *et al.* 2017). Indigenous people also contributed to species extinctions, such as among birds (Duncan *et al.* 2013; Braje *et al.* 2017), dwarf mammals (Hadjisterkotis *et al.* 2000; Braje *et al.* 2017), megafauna (Braje *et al.* 2017) or tree species (e.g. De Nascimento *et al.* 2009).

However, on most islands, the environmental impacts of indigenous people were much smaller than habitat degradation and biodiversity loss triggered through the arrival of Western people during colonial times (e.g. Fitzpatrick & Keegan 2007; Cheke & Hume 2008; Gillespie & Clague 2009; Royle 2014; Braje *et al.* 2017). In many cases, islands were largely deforested within a few decades after arrival, which reduced native habitats to tiny pockets in inaccessible places and

mostly at higher elevations. As a consequence, today, on many islands, the only habitats that are still represented at a substantial scale in a relatively undisturbed state are montane or alpine habitats (Juvik *et al.* 2014). Besides habitat destruction (including through fires), overexploitation of animal and plant populations (especially through hunting and for timber) and the introduction of invasive alien animals and plants were often major threat factors (Sodhi & Brook 2006; Caujapé-Castells *et al.* 2010; Kueffer & Kaiser-Bunbury 2014).

Biocultural heritage of islands

As a consequence of their socioecological histories, island cultures around the world represent a unique biocultural heritage and hold rich local knowledge (Loh & Harmon 2005; Royle 2008; Gillespie & Clague 2009; Hong 2013; McMillen *et al.* 2014; Kealiikanakaoleohaililani & Giardina 2016; Lauer 2017). The local knowledge of islanders encompasses far more than only a static collection of natural history facts (Lauer 2017); it also includes knowledge that is tacit, transmitted in oral form and encoded in mythologies, social innovations or land-use practices (Rolett 2008; Kealiikanakaoleohaililani & Giardina 2016; Lauer 2017). Local knowledge is embedded in particular worldviews, epistemologies, languages, symbolic systems and cultural practices. In addition, biocultural knowledge is dynamic. It is continuously adapting to environmental and social change, and it is evolving through learning and the incorporation of knowledge from outsiders (Lauer 2017). This means that much of local knowledge cannot simply be subsumed under Western thinking (Kealiikanakaoleohaililani & Giardina 2016; Lauer 2017). However, non-hierarchical and reciprocal dialogue between Western and local perspectives is still often lacking (Lauer 2017). A more equal relationship is possible, as demonstrated by the Cook/Forster Collection held at the Georg-August-University in Goettingen (Germany), which has been reinterpreted from a Māori perspective, thereby acknowledging the value of alternative epistemologies from islands (Schorch *et al.* 2016).

The biocultural history of islands also has great potential to inspire the imaginations of islanders and outsiders. Sailing across the Pacific using a replica of a Polynesian voyaging vessel, the *Hōkūle'a*, has occurred several times since the 1970s, illustrating how ancient traditions inspire modern hopes (Low 2013). Modern literature from the Caribbean and Pacific oceans tell stories of environmental loss and hope, and transport alternative ways of thinking about nature, non-human living beings and human–nature relationships to our time (DeLoughrey 2007). Environmentalism needs stories of the manifold possibilities of human life and future society that transgress the promise of technoscientific progress, and island stories provide one rooting of alternative narratives of current and future human–nature relationships. Teresia K. Teaiwa (an I-Kiribati and American poet and academic) proposed to make ‘island’ a verb (Baldacchino 2007, p. 514). She argued

that we have to learn how to ‘island’ the world – how to “behave as if we were all living on islands.”

Threatened livelihoods and the biocultural heritage of islands

Island societies and cultures and their biocultural heritage are threatened (Juvik & Juvik 1998; Baldacchino 2007; Baldacchino & Niles 2011; Connell 2013; Royle 2014). During colonial times, armed force and the introduction of diseases led to the massive reduction of indigenous populations on many islands, including the loss of power and cultural identity and practices. In the Hawaiian archipelago, for instance, the population size of native Hawaiians was reduced by *c.* 90% during the 19th century (Juvik & Juvik 1998). In post-colonial times, the vulnerability and powerlessness of islanders have remained high (Royle 2014). In particular, the dependence on former colonial powers remains much higher among island territories than on continents: most islands in the Arctic region, Mediterranean Sea and Atlantic Ocean, as well as many islands in the Central and Eastern Pacific, are part of or dependencies of continental countries, and there also remain many continental dependencies in the Caribbean Sea and Indian Ocean. Further, small-island states are often characterized by an economy with a low diversification and high dependence on the outside, and they are therefore highly exposed to external interests and influences. An increasing ratio of tourists to the local population and an increasing foreign ownership of land on islands can further weaken local identities (e.g. Royle 2014). Ultimately, unsustainable use of land and resources, as well as pollution and poor waste management, threaten the viability of many island societies (Juvik & Juvik 1998; Baldacchino & Niles 2011; Walker & Bellingham 2011; Connell 2013).

Recognition of the value of the biocultural heritage of islands

Recognition of the value of the biocultural heritage of islands has, in recent decades, been growing both locally (e.g. a renaissance of Polynesian culture in Hawaii) and internationally. Local knowledge is today considered a crucial component of any effective ecosystem management programme on an island (e.g., Jupiter *et al.* 2017; Lauer 2017). International organizations are paying attention to the biocultural diversity on islands (Hong 2013; Dahl 2017). The Representative List of the Intangible Cultural Heritage of Humanity of UNESCO listed many cultural innovations on islands; for instance, the Maloya dance music that originated among African and Malagasy slaves in La Réunion and the Sega dance in Mauritius; woven handcraft of the people of Papua; sand drawings in Vanuatu; or singing traditions in Corsica. There are also many cultural island landscapes listed on the UNESCO World Heritage List of cultural or mixed sites, such as the subsistence agriculture and sheep farming landscape on St Kilda in the Hebrides (Scotland), the vineyard




Figure 2 The global distribution of the world's islands. Black dots indicate the location of 17 883 islands listed by Weigelt *et al.* (2013); most of them are located along coastlines of continents or larger islands. Seventy islands or island groups that are mentioned in the text or are otherwise of special significance are indicated with a number (from west to east): 1: Tonga; 2: Northwestern Hawaiian Islands; 3: Samoa; 4: Aleutian Islands; 5: Hawaii Archipelago; 6: Tahiti (French Polynesia); 7: Marquesas Islands; 8: Pitcairn; 9: California Channel Islands; 10: Rapa Nui (Easter Island); 11: Galapagos; 12: Cuba; 13: Juan Fernandez Islands (Robinson Crusoe, Alejandro Selkirk, Santa Clara); 14: Jamaica; 15: Bahamas; 16: Hispaniola (Haiti and Dominican Republic); 17: Puerto Rico; 18: Guadeloupe; 19: Trinidad and Tobago; 20: Saint Lucia; 21: Falkland Islands; 22: Azores; 23: Cape Verde; 24: Iceland (including Surtsey); 25: Madeira; 26: Canary Islands (e.g. Tenerife, El Hierro); 27: Ascension; 28: Gough; 29: Ireland; 30: Saint Helena; 31: Balearic Islands (e.g. Majorca); 32: Gulf of Guinea Islands; 33: Corsica; 34: Eolie (Aeolian Islands); 35: Svalbard; 36: Aegean Islands (Greece); 37: Cyprus; 38: Zanzibar; 39: Comoros; 40: Aldabra Atoll; 41: Madagascar; 42: Socotra; 43: Seychelles (main group of mostly continental islands); 44: Mascarenes (Mauritius, La Réunion, Rodrigues); 45: Kerguelen; 46: Diego Garcia; 47: Maldives; 48: Sri Lanka; 49: Andaman and Nicobar Islands; 50: Sumatra (Indonesia); 51: Borneo; 52: Sulawesi (Indonesia); 53: Taiwan; 54: Philippines; 55: Okinawa (Ryukyu Islands, Japan); 56: Moluccas; 57: Palau; 58: Japan; 59: New Guinea; 60: Guam; 61: Tasmania; 62: Macquarie; 63: Solomon Islands; 64: New Caledonia; 65: Nauru; 66: Vanuatu; 67: New Zealand; 68: Phoenix; 69: Tuvalu; 70: Fiji.

culture on Pico Island in the Azores (dating back to the 15th century), the Le Morne Cultural Landscape of Mauritius that was used as a shelter by runaway slaves throughout the 18th and early 19th centuries or the Southern Lagoon Rock Islands in Palau with remains of a *c.* 3000-year-old culture that was abandoned in the 17th and 18th centuries (illustrating both the persistence and vulnerability of small-island communities).

GLOBAL CULTURAL AND ECOLOGICAL WEB

Islands occur in all world regions and at all latitudes (Fig. 2). By latitude, the highest number and density of islands is found between 50°N and 80°N, where they also make up *c.* 60–70% of the total land at these latitudes. In contrast, they make up only *c.* 20% of the total land in the subtropics and tropics, and less than 5% between –40°N and –65°N (Depraetere & Dahl 2007). At the highest latitudes, the Arctic islands (in the northern hemisphere) and the sub-Antarctic islands (in the southern hemisphere) host some of the last wilderness areas on the planet. The largest ocean is the Pacific, ranging from Southeast Asia in the West to the Americas in the

East. It encompasses almost half of the surface of the world's oceans and about a third of the total surface of the Earth. Thousands of islands are scattered across the Pacific, with most of them confined to subtropical and tropical latitudes. In contrast, the mid-latitudes of the Atlantic Ocean between Africa and the east coast of the Americas are relatively free of islands. The only island groups occurring in this region are those of Macaronesia in the northwest of Africa, some islands close to the coasts of Africa and South America and the isolated British islands of Ascension and Saint Helena. Other oceanic regions include the Caribbean Sea, Mediterranean Sea, Western Indian Ocean region and the archipelagos of Southeast Asia. For a bibliography of key reference literature for the different world's oceanic regions, see Kueffer *et al.* (2016). Gillespie and Clague (2009) include short entries about different island groups and oceanic regions. As a consequence of their global distribution, islands represent all climate zones, biogeographic regions and many human cultural groups. This global representation and association with the different world oceanic regions is essential for the manifold cultural and ecological riches of islands. The islands of the world can be

imagined as a global web of local nodes of high biological and cultural value (Fig. 1).

The web-like nature of islands becomes more accentuated when considering the interconnectedness of island peoples and cultures. Many island societies are today melting pots of different cultures: indigenous, early seafaring people (e.g. Arabs or Vikings), colonial settlers, slaves and workers that arrived on islands for different economic opportunities (first jobs on plantations, later in tourism or trade, now, for example, also banking). As a result, European influences are recognizable on islands in the Caribbean Sea, as well as in the Indian, Atlantic or Pacific oceans. In the Bonin (Ogasawara) Islands – an island group in the Pacific that was settled by immigrants from many different countries and now belongs to Japan – inhabitants speak a pidgin version of English with elements from different European and Austronesian languages and Japanese (Long 2007). Indian communities are established in the Caribbean, as well as in Mauritius (Indian Ocean) or Fiji (Pacific Ocean). African slaves were brought to La Réunion and Madagascar in the Indian Ocean, as well as to the islands of the Caribbean Sea, and as a consequence of French and African influences, islands at opposing sides of the planet share similar creole languages. Workers that came in the late 19th century to Hawaii for plantation work were Chinese, Japanese, Okinawans, Koreans, Puerto Ricans, Portuguese from the Azores archipelago in the middle of the Atlantic Ocean and Filipinos (Juvik & Juvik 1998). All of them tried to maintain some of their lifestyle from home (often other islands) with what was available in Hawaii, while adopting new elements from their new home; for instance, in cooking (Laudan 1996). In return, there is a growing diaspora community of islanders now living on continents (and in many cases, the money that these diaspora communities send to their relatives represents a major proportion of an island's income). Lastly, many islands still maintain close relationships with former colonial powers or are still part of continental countries, and there are increasingly regional and global networks among island nations.

Thus, island societies are globally interconnected through multiple and interwoven threads. Islands are places where people and traditions meet in a particular and bounded place that allows for multicultural exchange, but also enables the focusing of the creativity of individuals and groups. Not surprisingly, islands have been innovation hubs of world culture, such as Reggae in Jamaica, Haitian Voodoo tradition, Cuban music, Björk Guðmundsdóttir from Iceland, Jane Campion's films from New Zealand, or the Nobel Prize in Literature winner Derek Walcott from the Caribbean island of Saint Lucia.

Environmentalism promotes strengthening local actors so that they can speak up for their particular needs and perspectives, but it is also evident that local concerns and environmental issues need a strong voice internationally (Heise 2008; Turnhout *et al.* 2012). The need to strengthen the local voices of islanders at an international level is increasingly recognized in environmental politics (Kelman

2010; Dahl 2017). Similarly, environmental sciences struggle with integrating detailed knowledge about the complexities of local environmental issues with much coarser global data and analyses (Kueffer 2012; Sagarin & Pauchard 2012; Greschke & Tischler 2013; Lauer 2017). Islands are widely replicated and share many common environmental concerns, providing a unique opportunity to deal with this tension between the local and the global. Regional and global networking and multi-island research and policy programmes could help to pool institutional capacity and enable cross-island learning (Baldacchino 2007; Kueffer 2012; Kueffer *et al.* 2014). Indeed, networked conservation action can be more cost effective than isolated efforts, as has been demonstrated in the Mediterranean Basin, for instance (Kark *et al.* 2009).

There are global networks such as the Alliance of Small Island States (AOSIS), the Global Island Partnership (GLISPA), the Small Island Developing States Network (SIDSnet), the Society for Island Biology (SIB), the International Small Islands Studies Association (ISISA) and the Network of Island Universities (RETI), and there are numerous local and regional networks focused on specific islands, archipelagos and oceanic regions (Baldacchino 2007; Kelman 2010; Kueffer *et al.* 2016; Dahl 2017). There are also several examples of multi-campus universities and shared teaching and research programmes that link experts on several islands, such as the University of the West Indies (UWI), the University of the South Pacific (USP) and the University of the Highlands and Islands (UHI). Together, these networks can assist in linking global support of environmental policies on islands with regional and local action and the expertise of island-based stakeholders, policy-makers and experts. Emerging opportunities for sharing knowledge among environmentalists on islands include internet-based communication tools such as online courses or video-conferences (Norder & Rijdsdijk 2016).

ISLANDS AS LABORATORIES

Islands have often been and still are the basis of major scientific breakthroughs in disciplines ranging from anthropology and the social sciences to geosciences, biogeography, ecology and evolutionary biology, and they have inspired indigenous cultures as well as modern literature and arts for centuries (DeLoughrey 2007; Royle 2008; Gillespie & Clague 2009; Kueffer *et al.* 2014; Royle 2014; Kueffer *et al.* 2016; Braje *et al.* 2017). Islands are used as model systems in a wide range of disciplines, including ecology, evolutionary biology and biogeography (Vitousek 2004; Kueffer *et al.* 2014), systems ecology (Davies *et al.* 2016), conservation biology (Kueffer 2012; Ewel *et al.* 2013; Kueffer & Kaiser-Bunbury 2014), the environmental sciences (Walker & Bellingham 2011), historical ecology (Rolett & Diamond 2004; Vitousek *et al.* 2004; Braje *et al.* 2017), sustainability science (McDaniel & Gowdy 2000; Diamond 2005; Kelman *et al.* 2015), socioecological research (Chertow *et al.* 2013) and anthropology and the social sciences (Royle 2014; Coulthard

et al. 2017). Some of the reasons why islands are attractive as model systems are related to the three general features of islands discussed above. As hotspots of unique biological and cultural features, they provide many opportunities for discovery. Evolution on islands, for instance, has inspired modern evolutionary biology since its formulation through Charles Darwin and Alfred Russel Wallace (Kueffer *et al.* 2014). High geophysical variability between islands and within islands enabled the studying of how ecosystem processes respond to varying abiotic conditions (Vitousek 2004). As paradigmatic places of human–environment relationships, they provide a wealth of case studies of how past societies in different ecological, social and cultural contexts dealt with limited resources (e.g. Braje *et al.* 2017). Today, many environmental problems are already very acute, which forces agents on islands to act. This enables scholars working on the adaptation to and mitigation of environmental problems in fields ranging from conservation biology and restoration ecology (Ewel *et al.* 2013; Kueffer *et al.* 2014) to ecosystem management (Jupiter *et al.* 2017), sustainability science (Kelman *et al.* 2015) and social wellbeing (Coulthard *et al.* 2017) to work in situations where environmental problems are already real that are only predicted in other places. On islands, global challenges become concrete: we know which species might die out within decades; we know which areas will be threatened by sea-level rise; we can see coral reef bleaching; on many islands it has become obvious that traffic, energy consumption, waste and population densities have reached the carrying capacity of the land; and we can guess what it will mean for island economies when cheap energy is not available anymore.

The global distribution of islands allows for the replication of studies and datasets, which was central for the development of MacArthur and Wilson's dynamic theory of island biogeography, for instance – by far the most widely discussed theory in biogeography (Kueffer *et al.* 2014). Multi-island comparative studies also allow for the exploitation of differences between islands as natural experiments; for instance, enabling us to understand how different political systems on otherwise-similar islands affected sustainable land use (Diamond 2005). In addition, practitioners on different islands can learn from each other's experiences (Kueffer 2012).

Another reason why islands are considered to be good model systems is their small size and isolation. This makes it easier to define boundaries for analysis, collect comprehensive or almost complete data across whole systems and compile and integrate perspectives of multiple disciplines and many datasets on the same phenomenon. Davies *et al.* (2016), for instance, attempt to build computer simulations of entire social–ecological systems based on data from Moorea, French Polynesia. The deCODE Genetics database in Iceland is built on the vision of compiling a comprehensive genetics and health data database of the Icelandic population (Greenhough 2006). Islands are also ideal locations for long-term place-based research (Kueffer 2006; Chertow *et al.* 2013). As with any model, framing islands as model systems is always a

simplification and a way of favouring some understandings and valuations of a particular island (or islands in general) over others (e.g. Greenhough 2006).

Seeing islands as laboratories encompasses more than understanding them only as model systems. It positions them as places for experimentation and imagination. In Western literature and arts, islands have, for thousands of years, been places of social experimentation and utopia or dystopia, e.g. Homer's *Odyssey* in ancient Greek times, Thomas More's *Utopia* from 1516, *Robinson Crusoe* by Daniel Defoe from 1719, *Lord of the Flies* by William Golding from 1954, T.C. Boyle's *San Miguel* from 2012, Craigie Horsfield's artwork from El Hierro (Canary Islands) or many 'real' individuals or groups that tried to find a place removed from the world on islands (Clark 2001; Royle 2014). Both T.C. Boyle and Craigie Horsfield portray islands as the ultimate frontier of the human exploration of space. Today, islands are often discussed as experimental spaces for the resurrection and innovation of sustainable living (Baldacchino 2007; Rolett 2008; Baldacchino & Niles 2011; Kelman *et al.* 2015; Kealiikanakaoleohailani & Giardina 2016). Due to their small size, it is often easier to bring stakeholders and experts from different political or economic sectors together to work on a common issue. Islands have also become laboratories for experimenting with new biodiversity conservation strategies for highly disturbed novel ecosystems and landscapes (Ewel *et al.* 2013; Kueffer & Kaiser-Bunbury 2014). Island conservationists have been leaders in bringing species back from the brink of extinction through *ex situ* conservation in zoos and botanical gardens and later reintroduction to the wild. There are also many successful examples of habitat restoration (e.g. Florens & Baider 2013), including on small offshore islands, where all invasive species can be eradicated, which then can become new safe places for threatened species (Townes *et al.* 1990; Zino 2008; Kueffer & Kaiser-Bunbury 2014). In addition, there have been first experiments with designing new nature, such as through rewilding (i.e. the introduction of alien species to the wild that are functional analogues of extinct native species, with the idea that they can replace lost ecological functions; Hansen 2010). All of these new conservation strategies will grow in importance on continents, but have first been recognized and addressed and are most evident on islands (Kueffer & Kaiser-Bunbury 2014).

The use of islands as laboratories also has a problematic history. External forces used islands recurrently to do their dangerous experiments far away from home, including for nuclear weapon tests (Royle 2014). In addition, framing islands as model systems or laboratories has frequently imposed external understandings and values on local island peoples. In environmental conservation in particular, islands have in the past too often been the playgrounds of foreign countries and international organizations and their predefined environmental management approaches (compare, for example, Rodríguez *et al.* 2007). The idea of islands as real-world laboratories for innovating environmental solutions

is therefore only valid if such experiments are guided through local communities. Local and traditional knowledge on islands has to be recognized as the basis for environmental conservation with external knowledge complementing it, not vice versa. Relying on the biocultural wisdom of islanders might often mean that current environmental policies on islands are fundamentally reconsidered. Environmental problems must be framed according to local needs and perspectives, and knowledge must be represented, encoded and communicated in locally sensitive forms (i.e. by using local forums of communication; e.g. oral communication, or embedded in ritual settings), relying on local metaphors, stories, language, words and images, and by reflecting upon who is doing the research, communicating it and acting upon it. On islands, such a change of perspective is particularly pertinent, both because of the rich cultural heritage of island people and the too frequent and too prominent neglect of local perspectives by international conservation actors and bodies.

ISLAND FUTURES

Islands are globally distributed, but share through their insularity many common values and threats. They are hotspots of cultural, biological and geophysical riches, and they can serve as globally connected laboratories of environmental conservation, sustainable living and sociocultural innovation. We have highlighted the threats affecting the unique values of islands. Much of the cultural and biological heritage on islands might disappear within the 21st century without major action. There is therefore an immediate need to demonstrate on islands how sustainability and environmental conservation can be achieved in real places, and now.

However, beyond their vulnerabilities, islands should foremost be seen as places of hope and innovation. This will require that local livelihoods and their actions and biocultural knowledge and heritage are better recognized. In the coming decades, the environmental crisis will likely develop into a perfect storm of multiple major challenges – climate change, overexploitation of natural resources and associated food and energy crises, amongst others. These crises are too big to be tackled in their entirety, which gives many people a feeling of hopelessness. Islands have been used as illustrations of the severity of our environmental crisis – as examples of past or imminent environmental collapse and of major biodiversity and cultural loss. But islands can also be places of hope, where new ways of sustainable living and environmental conservation might be developed at a manageable scale.

ACKNOWLEDGEMENTS

An early version of this manuscript benefitted from helpful comments from Nicholas Polunin, Paulo Borges and two anonymous reviewers.

FINANCIAL SUPPORT

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

CONFLICT OF INTEREST

None.

ETHICAL STANDARDS

None.

References

- Alcover, J. A., Pieper, H., Pereira, F. & Rando, J. C. (2015) Five new extinct species of rails (Aves: Gruiformes: Rallidae) from the Macaronesian Islands (North Atlantic Ocean). *Zootaxa* **4057**(2): 151–190.
- Baldacchino, G., ed. (2007) *A World of Islands. An Island Studies Reader*. Luqa, Malta: Agenda Academic Publishers.
- Baldacchino, G. & Niles, D., eds. (2011) *Island Futures. Conservation and Development Across the Asia-Pacific Region*. New York, NY, USA: Springer.
- Braje, T. J., Leppard, T. P., Fitzpatrick, S. M. & Erlandson, J. M. (2017) Archaeology, historical ecology and anthropogenic island ecosystems. *Environmental Conservation*, in press. doi: 10.1017/S0376892917000261.
- Caujapé-Castells, J., Tye, A., Crawford, D. J., Santos-Guerra, A., Sakai, A., Beaver, K., Lobin, W., Florens, F. B. V., Moura, M., Jardim, R., Gomes, I. & Kueffer, C. (2010) Conservation of oceanic island floras: present and future global challenges. *Perspectives in Plant Ecology Evolution and Systematics* **12**(2): 107–130.
- CBD (2014) *Island Biodiversity – Island Bright Spots in Conservation & Sustainability*. Montreal, Canada: Convention on Biological Diversity.
- Cheke, A. & Hume, J. (2008) *Lost Land of the Dodo: The Ecological History of Mauritius, Reunion, and Rodrigues*. New Haven, CT, USA: Yale University Press.
- Chertow, M., Fugate, E. & Ashton, W. (2013) The intimacy of human–nature interactions on islands. In: *Long Term Socio-Ecological Research*, eds. S. J. Singh, H. Haberl, M. Chertow, M. Mirtl, & M. Schmid, pp. 315–337. Berlin, Germany: Springer.
- Chiba, S. & Cowie, R. H. (2016) Evolution and extinction of land snails on oceanic islands. *Annual Review of Ecology, Evolution, and Systematics* **47**: 123–141.
- Clark, T. (2001) *Searching for Crusoe*. New York, NY, USA: Random House.
- Connell, J. (2013) *Islands at Risk?: Environments, Economies and Contemporary Change*. Cheltenham, UK: Edward Elgar Publishing.
- Coulthard, S., Evans, L., Turner, R., Mills, D., Foale, S., Abernethy, K., Hicks, C. & Monnereau, I. (2017) Exploring ‘islandness’ and the impacts of nature conservation through the lens of wellbeing. *Environmental Conservation*, in press. doi: 10.1017/S0376892917000273.
- Dahl, A. L. (2017) Island conservation issues in international conventions and agreements. *Environmental Conservation*, in press.

- Davies, N., Field, D., Gavaghan, D., Holbrook, S. J., Planes, S., Troyer, M., Bonsall, M., Claudet, J., Roderick, G. & Schmitt, R. J. (2016) Simulating social-ecological systems: the Island Digital Ecosystem Avatars (IDEA) consortium. *GigaScience* 5(1): 14.
- De Nascimento, L., Willis, K. J., Fernández-Palacios, J. M., Criado, C. & Whittaker, R. J. (2009) The long-term ecology of the lost forests of La Laguna, Tenerife (Canary Islands). *Journal of Biogeography* 36(3): 499–514.
- DeLoughrey, E. M. (2007) *Routes and Roots: Navigating Caribbean and Pacific Island Literatures*. Honolulu, HI, USA: University of Hawaii Press.
- Depraetere, C. & Dahl, A. L. (2007) Island locations and classifications. In: *A World of Islands. An Island Studies Reader*, ed. G. Baldacchino, pp. 57–105. Luqa, Malta: Agenda Academic Publishers.
- Diamond, J. (2005) *Collapse: How Societies Choose to Fail or Succeed*. New York, NY, USA: Viking Press.
- Duncan, R. P., Boyer, A. G. & Blackburn, T. M. (2013) Magnitude and variation of prehistoric bird extinctions in the Pacific. *Proceedings of the National Academy of Sciences of the United States of America* 110(16): 6436–6441.
- Edwards, P. J., Fleischer-Dogley, F. & Kaiser-Bunbury, C. N. (2015) The nutrient economy of *Lodoicea maldivica*, a monodominant palm producing the world's largest seed. *New Phytologist* 206(3): 990–999.
- Ewel, J. J., Mascaro, J., Kueffer, C., Lugo, A. E., Lach, L. & Gardener, M. R. (2013) Islands: where novelty is the norm. In: *Novel Ecosystems. Intervening in the New Ecological World Order*, eds. R. J. Hobbs, E. S. Higgs & C. M. Hall, pp. 29–44. Oxford, UK: Wiley-Blackwell.
- Fernández-Palacios, J. M., Rijdsdijk, K. F., Norder, S. J., Otto, R., Nascimento, L., Fernández-Lugo, S., Tjørve, E. & Whittaker, R. J. (2016) Towards a glacial-sensitive model of island biogeography. *Global Ecology and Biogeography* 25(7): 817–830.
- Fitzpatrick, S. M. & Keegan, W. F. (2007) Human impacts and adaptations in the Caribbean Islands: an historical ecology approach. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* 98(1): 29–45.
- Florens, F. B. V. & Baider, C. (2013) Ecological restoration in a developing island nation: how useful is the science? *Restoration Ecology* 21(1): 1–5.
- Gillespie, R. G. & Clague, D. A., eds. (2009) *Encyclopedia of Islands*. Berkeley, CA, USA: University of California Press.
- Greenhough, B. (2006) Tales of an island-laboratory: defining the field in geography and science studies. *Transactions of the Institute of British Geographers* 31(2): 224–237.
- Grenyer, R., Orme, C. D. L., Jackson, S. F., Thomas, G. H., Davies, R. G., Davies, T. J., Jones, K. E., Olson, V. A., Ridgely, R. S. & Rasmussen, P. C. (2006) Global distribution and conservation of rare and threatened vertebrates. *Nature* 444(7115): 93–96.
- Greschke, H. M. & Tischler, J., eds. (2013) *The Challenges of Global Climate Change. Locally-Grounded Interdisciplinary Approaches*. Berlin, Germany: Springer.
- Hadjisterkotis, E., Masala, B. & Reese, D. S. (2000) The origin and extinction of the large endemic Pleistocene mammals of Cyprus. *Biogeographia* 21 [www document]. URL <http://escholarship.org/uc/item/40b208r7>
- Hansen, D. M. (2010) On the use of taxon substitutes in rewilding projects on islands. In: *Islands and Evolution*, eds. V. Pérez-Mellado & M. M. Ramon, pp. 111–146. Menorca, Spain: Institut Menorquí d'Estudis.
- Harter, D. E., Irl, S. D., Seo, B., Steinbauer, M. J., Gillespie, R., Triantis, K. A., Fernández-Palacios, J.-M. & Beierkuhnlein, C. (2015) Impacts of global climate change on the floras of oceanic islands – projections, implications and current knowledge. *Perspectives in Plant Ecology, Evolution and Systematics* 17(2): 160–183.
- Heise, U. (2008) *Sense of Place and Sense of Planet*. Oxford, UK: Oxford University Press.
- Johnson, T. & Stattersfield, A. (1990) A global review of island endemic birds. *Ibis* 132(2): 167–180.
- Hong, S.-K. (2013) Biocultural diversity conservation for island and islanders: necessity, goal and activity. *Journal of Marine and Island Cultures* 2(2): 102–106.
- Jupiter, S. D., Wenger, A., Klein, C. J., Albert, S., Mangubhai, S., Nelson, J., Teneva, L., Tulloch, V. J., White, A. T. & Watson, J. E. (2017) Opportunities and constraints for implementing integrated land–sea management on islands. *Environmental Conservation*, in press. doi: 10.1017/S0376892917000091.
- Juvik, J. O., Kueffer, C. & Juvik, S. (2014) Losing the high ground: rapid transformation of tropical island alpine and subalpine environments. *Arctic, Antarctic, and Alpine Research* 46(4): 705–708.
- Juvik, S. P. & Juvik, J. O. (1998) *Atlas of Hawaii. 3rd Edition*. Honolulu, HI, USA: University of Hawai'i Press.
- Kark, S., Levin, N., Grantham, H. S. & Possingham, H. P. (2009) Between-country collaboration and consideration of costs increase conservation planning efficiency in the Mediterranean Basin. *Proceedings of the National Academy of Sciences of the United States of America* 106(36): 15368–15373.
- Kealiikanakaoleohaililani, K. & Giardina, C. P. (2016) Embracing the sacred: an indigenous framework for tomorrow's sustainability science. *Sustainability Science* 11(1): 57–67.
- Kelman, I. (2010) Hearing local voices from small island developing states for climate change. *Local Environment* 15(7): 605–619.
- Kelman, I., Burns, T. R. & des Johansson, N. M. (2015) Islander innovation: a research and action agenda on local responses to global issues. *Journal of Marine and Island Cultures* 4(1): 34–41.
- Kier, G., Kreft, H. & Lee, T. M. (2009) A global assessment of endemism and species richness across island and mainland regions. *Proceedings of the National Academy of Sciences of the United States of America* 106: 9322–9327.
- Koshiha, S., Besebes, M., Soaladaob, K., Ngiraingas, M., Isechal, A. L., Victor, S. & Golbuu, Y. (2014) 2000 years of sustainable use of watersheds and coral reefs in Pacific Islands: a review for Palau. *Estuarine, Coastal and Shelf Science* 144: 19–26.
- Kueffer, C. (2006) Integrative ecological research: case-specific validation of ecological knowledge for environmental problem solving. *Gaia* 15(2): 115–120.
- Kueffer, C. (2012) The importance of collaborative learning and research among conservationists from different oceanic islands. *Revue d'Ecologie (Terre et Vie) Suppl. 11*: 125–135.
- Kueffer, C., Drake, D. & Fernández-Palacios, J. M. (2016) Island biology. In: *Oxford Bibliographies in Ecology*, ed. D. Gibson. Oxford, UK: Oxford University Press [www document]. URL <http://www.oxfordbibliographies.com/view/document/obo-9780199830060/obo-9780199830060-0149.xml?rskey=lzfTBH&result=63>
- Kueffer, C., Drake, D. R. & Fernández-Palacios, J. M. (2014) Island biology: looking towards the future. *Biology Letters* 10(10): 20140719.

- Kueffer, C. & Kaiser-Bunbury, C. (2014) Reconciling conflicting perspectives for biodiversity conservation in the Anthropocene. *Frontiers in Ecology and Environment* 12(2): 131–137.
- Laudan, R. (1996) *The Food of Paradise: Exploring Hawaii's Culinary Heritage*. Honolulu, HI, USA: University of Hawaii Press.
- Lauer, M. (2017) Changing understandings of local knowledge in island environments. *Environmental Conservation*, in press. doi: 10.1017/S0376892917000303.
- Leonard, D. L. (2008) Recovery expenditures for birds listed under the US Endangered Species Act: the disparity between mainland and Hawaiian taxa. *Biological Conservation* 141(8): 2054–2061.
- Loh, J. & Harmon, D. (2005) A global index of biocultural diversity. *Ecological Indicators* 5(3): 231–241.
- Lomolino, M. V. (2005) Body size evolution in insular vertebrates: generality of the island rule. *Global Ecology & Biogeography* 32: 1683–1699.
- Long, D. (2007) *English on the Bonin (Ogasawara) Islands*. Durham, NC, USA: Duke University Press.
- Low, S. (2013) *Hawaiiki Rising: Hōkūle'a, Nainoa Thompson, and the Hawaiian Renaissance*. Waipahu, HI, USA: Island Heritage Publishing.
- McDaniel, C., & Gowdy, J. (2000) *Paradise for Sale: A Parable of Nature*. Berkeley, CA, USA: University of California Press.
- McMillen, H. L., Ticktin, T., Friedlander, A., Jupiter, S. D., Thaman, R., Campbell, J., Veitayaki, J., Giambelluca, T., Nihmei, S., Rupeni, E., Apis-Overhoff, L., Aalbersberg, W. & Orcheron, D. F. (2014) Small islands, valuable insights: systems of customary resource use and resilience to climate change in the Pacific. *Ecology and Society* 19(4): 44.
- Menard, W. (1986) *Islands*. New York, NY, USA: Scientific American.
- Mulder, C. P., Anderson, W. B., Towns, D. R. & Bellingham, P. J. (2011) *Seabird Islands: Ecology, Invasion, and Restoration*. Oxford, UK: Oxford University Press.
- Zachos, F. E. & Habel, J. C., eds. (2011) *Biodiversity Hotspots. Distribution and Protection of Conservation Priority Areas*. Berlin, Germany: Springer.
- Norder, S. & Rijdsdijk, K. (2016) Interdisciplinary island studies: connecting the social sciences, natural sciences and humanities. *Island Studies Journal* 11(2): 673–686.
- Nunn, P. D. (1994) *Oceanic Islands*. Oxford, UK: Blackwell.
- Paxton, E. H., Camp, R. J., Gorresen, P. M., Crampton, L. H., Leonard, D. L. & VanderWerf, E. A. (2016) Collapsing avian community on a Hawaiian island. *Science Advances* 2(9): e1600029.
- Rodríguez, J., Taber, A., Daszak, P., Sukumar, R., Valladares-Padua, C., Padua, S., Aguirre, L., Medellín, R., Acosta, M. & Aguirre, A. (2007) Globalization of conservation: a view from the South. *Science* 317(5839): 755.
- Rolett, B. & Diamond, J. (2004) Environmental predictors of pre-European deforestation on Pacific islands. *Nature* 431(7007): 443–446.
- Rolett, B. V. (2008) Avoiding collapse: pre-European sustainability on Pacific islands. *Quaternary International* 184(1): 4–10.
- Royle, S. A. (2008) *Geography of Islands: Small Islands Insularity*. London, UK, and New York, NY, USA: Routledge.
- Royle, S. A. (2014) *Islands: Nature and Culture*. London, UK: Reaktion.
- Sagarin, R. & Pauchard, A. (2012) *Observation and Ecology. Broadening the Scope of Science to Understand a Complex World*. Washington, DC, USA: Island Press.
- Schorch, P., McCarthy, C. & Hakiwai, A. (2016) Globalizing Māori museology: reconceptualizing engagement, knowledge, and virtuality through Mana Taonga. *Museum Anthropology* 39(1): 48–69.
- Sodhi, N. S. & Brook, B. W. (2006) *Southeast Asian Biodiversity in Crisis*. Cambridge, UK: Cambridge University Press.
- Terzopoulou, S., Rigal, F., Whittaker, R. J., Borges, P. A. V. & Triantis, K. A. (2015) Drivers of extinction: the case of Azorean beetles. *Biology Letters* 11: 20150273.
- Towns, D.R., Daugherty, H. & Atkinson, I.A.E., eds. (1990) *Ecological Restoration of New Zealand Islands*. Wellington, New Zealand: Department of Conservation.
- Triantis, K. A., Borges, P. A. V., Ladle, R. J., Hortal, J., Cardoso, P., Gaspar, C., Dinis, F., Mendonca, E., Silveira, L. M. A., Gabriel, R., Melo, C., Santos, A. M. C., Amorim, I. R., Ribeiro, S. P., Serrano, A. R. M., Quartau, J. A. & Whittaker, R. J. (2010) Extinction debt on oceanic islands. *Ecography* 33(2): 285–294.
- Turnhout, E., Bloomfield, B., Hulme, M., Vogel, J. & Wynne, B. (2012) Listen to the voices of experience. *Nature* 488: 454–455.
- Vitousek, P. M. (2004) *Nutrient Cycling and Limitation. Hawai'i as a Model System*. Princeton, NJ, USA: Princeton University Press.
- Vitousek, P. M., Ladefoged, T. N., Kirch, P. V., Hartshorn, A. S., Graves, M. W., Hotchkiss, S. C., Tuljapurkar, S. & Chadwick, O. A. (2004) Soils, agriculture, and society in precontact Hawaii. *Science* 304(5677): 1665–1669.
- Waldron, A., Mooers, A. O., Miller, D. C., Nibbelink, N., Redding, D., Kuhn, T. S., Roberts, J. T. & Gittleman, J. L. (2013) Targeting global conservation funding to limit immediate biodiversity declines. *Proceedings of the National Academy of Sciences of the United States of America* 110(29): 12144–12148.
- Walker, L. R. & Bellingham, P. (2011) *Island Environments in a Changing World*. Cambridge, UK: Cambridge University Press.
- Weigelt, P., Jetz, W. & Kreft, H. (2013) Bioclimatic and physical characterization of the world's islands. *Proceedings of the National Academy of Sciences of the United States of America* 110(38): 15307–15312.
- Weigelt, P., Kissling, W. D. & Kisel, Y. (2015) Global patterns and drivers of phylogenetic structure in island floras. *Scientific Reports* 5: 12213.
- Weigelt, P. & Kreft, H. (2013) Quantifying island isolation: insights from global patterns of insular plant species richness. *Ecography* 36: 417–429.
- Wetzel, F. T., Kissling, W., Beissmann, H. & Penn, D. (2012) Future climate change driven sea-level rise: secondary consequences from human displacement for island biodiversity. *Global Change Biology* 18: 2707–2719.
- Whistler, W. A. (2009) *Plants of the Canoe People. An Ethnobotanical Voyage Through Polynesia*. Lawai, Kauai, HI, USA: National Tropical Botanical Garden.
- Whittaker, R. J. & Fernández-Palacios, J. M. (2007) *Island Biogeography. Ecology, Evolution, and Conservation. 2nd Edition*. Oxford, UK: Oxford University Press.
- Zino, F., Hounscome, M.V., Buckle, A.P. & Biscoito, M. (2008) Was the removal of rabbits and house mice from Selvagem Grande beneficial to the breeding of Cory's shearwaters *Calonectris diomedea borealis*? *Oryx* 42: 151–154.