# **ETH** zürich

# Search for single production of vector-like quarks decaying to a b quark and a Higgs boson

#### **Journal Article**

#### **Author(s):**

CMS Collaboration; Sirunyan, Albert M.; Backhaus, Malte; Bäni, Lukas; Berger, Pirmin; Casal, Bruno; Dissertori, Günther; Dittmar, Michael; Donegà, Mauro; Dorfer, Christian; [Grab, Christophorus](https://orcid.org/0000-0002-6182-3380) D; Heidegger, Constantin; Hits, Dmitry; Hoss, Jan; Kasieczka, Gregor; Klijnsma, Thomas; Lustermann, Werner; Mangano, Boris; Marionneau, Matthieu; Meinhard, Maren T.; Meister, Daniel; Micheli, Francesco; Musella, Pasquale; Nessi-Tedaldi, Francesca; Pandolfi, Francesco; Pata, Joosep; Pauss, Felicitas; Perrin, Gaël; Perrozzi, Luca; Quittnat, Milena; Reichmann, Michael; Sanz Becerra, Diego A.; Schönenberger, Myriam; Shchutska, Lesya; Tavolaro, Vittorio R.; Theofilatos, Konstantinos; Vesterbacka Olsson, Minna L.; [Wallny, Rainer](https://orcid.org/0000-0001-8038-1613) D; Zhu, De H.; et al.

# **Publication date:**

2018-06

#### **Permanent link:** <https://doi.org/10.3929/ethz-b-000271730>

**Rights / license:** [Creative Commons Attribution 4.0 International](http://creativecommons.org/licenses/by/4.0/)

#### **Originally published in:**

Journal of High Energy Physics 2018(6), [https://doi.org/10.1007/JHEP06\(2018\)031](https://doi.org/10.1007/JHEP06(2018)031)

PUBLISHED FOR SISSA BY 2 SPRINGER

Received: February 5, 2018 Revised: May 1, 2018 Accepted: May 15, 2018 PUBLISHED: June 5, 2018

# Search for single production of vector-like quarks decaying to a b quark and a Higgs boson



# The CMS collaboration

E-mail: [cms-publication-committee-chair@cern.ch](mailto:cms-publication-committee-chair@cern.ch)

Abstract: A search is presented for single production of heavy vector-like quarks (B) that decay to a Higgs boson and a b quark, with the Higgs boson decaying to a highly boosted  $b\bar{b}$  pair reconstructed as a single collimated jet. The analysis is based on data collected by the CMS experiment in proton-proton collisions at  $\sqrt{s} = 13 \text{ TeV}$ , corresponding to an integrated luminosity of 35.9 fb−<sup>1</sup> . The data are consistent with background expectations, and upper limits at 95% confidence level on the product of the B quark cross section and the branching fraction are obtained in the range 1.28–0.07 pb, for a narrow B quark with a mass between 700 and 1800 GeV. The production of B quarks with widths of 10, 20 and 30% of the resonance mass is also considered, and the sensitivities obtained are similar to those achieved in the narrow width case. This is the first search at the CERN LHC for the single production of a B quark through its fully hadronic decay channel, and the first study considering finite resonance widths of the B quark.

Keywords: Beyond Standard Model, Hadron-Hadron scattering (experiments), vectorlike quarks

ArXiv ePrint: [1802.01486](https://arxiv.org/abs/1802.01486)



#### Contents



## 1 Introduction

With the discovery of the Higgs boson  $(H)$  by the ATLAS  $[1]$  and CMS  $[2, 3]$  experiments at the CERN LHC, the standard model (SM) of particle physics has now been completely confirmed. However, the SM does not address, for example, problems related to the nature of the electroweak symmetry breaking and the hierarchy between the electroweak and the Planck mass scales. Several extensions of the SM address such issues through the introduction of new particles that allow the cancellation of loop corrections to the mass of the Higgs boson [4]. Supersymmetric theories propose bosonic partners of the top quark to address the hierarchy problem; other models such as Little Higgs or Composite Higgs boson models [5–8] overcome the hierarchy problem by introducing heavy fermionic resonances called vector-like quarks (VLQs) [4, 9–11]. The vector-like nature of these quarks does not exclude their having a fundamental mass, in contrast to chiral fermions, which acquire mass via electroweak symmetry breaking in the SM. The VLQs are therefore not excluded by present searches, unlike a fourth generation of SM quarks that is ruled out by electroweak precision measurements [12, 13], and by the measured properties of the SM Higgs boson [14–16]. Previous searches for VLQs have been performed by the ATLAS [17–22] and CMS [23–29] experiments in proton-proton collisions recorded at centre-of-mass energies of 7, 8, and 13 TeV.

We present a search for electroweak production of single vector-like B quarks with electrical charge  $-1/3e$ , with e the proton charge, that decay to a bottom (b) quark and a Higgs boson. The search uses pp events collected by the CMS experiment at a centre-of-mass energy of 13 TeV, corresponding to an integrated luminosity of  $35.9 \text{ fb}^{-1}$ .

We study the fully hadronic final state with the Higgs boson decaying to a pair of b quarks. Figure 1 illustrates the electroweak production of a B quark in association with a b and a light-flavour quark, typically emitted into the forward region of the detector.

The B decay channel considered in this analysis is  $B \to Hb$ . However, the B quark can also decay into Zb, Wt, and possibly into lighter states predicted in models beyond the SM that have model-dependent branching fractions. Our results are interpreted assuming that the B quark belongs to a singlet or doublet representation and that it decays exclusively to SM particles. The singlet branching fractions of the B quark into Hb, Zb, and Wt are  $\mathcal{B} \approx 25, 25, \text{ and } 50\%, \text{ and the doublet branching fractions are 50, 50, and 0\%, and all}$ depend on the vector-like quark mass  $m_{\rm B}$ .

Previous CMS searches for vector-like B quarks relied on the assumption of a decay width that is narrow compared to the experimental resolution. The present analysis, in addition to searching for B quarks with narrow decay widths, also explores the possibility that B quarks have a non-negligible width, with values up to 30% of the resonance mass. In comparison, the experimental resolution in the reconstructed B mass, defined as the ratio between the root-mean-square width of the peak and its mean position, ranges between 8 and 15%, depending on the mass hypothesis. In addition to broadening the width of the observed signal, the intrinsic width of the resonance would modify the kinematic distributions of the final state, thus changing the selection efficiency. These effects are taken into account in this analysis.

The cross section for single production of a B quark depends on  $m<sub>B</sub>$  and its electroweak couplings to SM particles. The kinematic distributions depend only on the total width of the B quark. The benchmark model in this analysis assumes a weak coupling of the B quark to the Z boson and b quark. Because of the mixing between B and the SM bottom quark in models where B is a singlet or part of a doublet, the BbZ electroweak coupling has a predominant chirality, respectively, right- or left-handed. The coupling chirality can potentially affect the kinematic distributions. We explicitly checked and found that these effects are negligible for the channel discussed in this work, and our results can therefore be interpreted in both singlet and doublet models.

# 2 The CMS detector and particle reconstruction

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. A silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two end sections, reside within the solenoid. Forward calorimeters extend the pseudorapidity  $(\eta)$  coverage provided by the barrel and end detectors. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system and kinematic variables, can be found in ref. [30].

Events of interest are selected using a two-tiered trigger system [31]. The first level, composed of specialized hardware processors, uses information from the calorimeters and



Figure 1. The leading-order Feynman diagram for the production of a single vector-like B quark in association with a b quark and light-flavour quark, and its decay to a Higgs boson and a b quark.

muon detectors to select events at a rate of  $\approx$ 100 kHz within a time interval of less than 4  $\mu$ s. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event-reconstruction software optimized for fast processing that reduces the event rate to  $\approx$ 1 kHz before data storage.

Event reconstruction is based on the CMS particle-flow (PF) algorithm [32], which reconstructs and identifies each individual particle through an optimized combination of information from the various elements of the CMS detector. The energy of electrons is defined through the combination of the electron momentum at the primary interaction vertex determined in the tracker, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track from the primary pp collision vertex. The energy of muons is obtained from the curvature of the corresponding track. The reconstructed energy of charged hadrons is extracted from the reconstructed tracks in the tracker and their matching energy depositions in ECAL and HCAL. Energy depositions are corrected for ignoring calorimeter readouts that are close to threshold (zero suppression) and for the response function of calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

Jets are reconstructed by clustering PF candidates using the anti- $k_T$  algorithm [33]. Prior to clustering, the charged-hadron subtraction algorithm [34] is applied to the event to reduce the effects of pileup (i.e. additional pp collisions occurring within the same or neighbouring LHC bunch crossings).

This algorithm discards charged hadrons not originating from the primary vertex, which is defined as the reconstructed vertex with the largest value of summed  $p_T^2$  of charged hadrons contributing to jets. The jets are clustered using the jet finding algorithm [33, 35], which defines the associated missing transverse momentum taken as the negative of the vector sum of the  $p<sub>T</sub>$  of those jets. We consider jets with different distance parameter of

 $\Delta R =$ √  $(\Delta y)^2 + (\Delta \phi)^2 = 0.4$  or 0.8, with y the rapidity, referred to as AK4 and AK8 jets, respectively.

The residual pileup contamination from neutral hadrons is subtracted, assuming that it is proportional to the event energy density and the jet area, estimated using the FASTJET package [35]. Jet momenta are determined from the vectorial sum of all the individual PF particles in the jet. The energy scale calibrations obtained from Monte Carlo (MC) simulation are applied to correct the four-momenta of jets. Residual corrections, accounting for remaining discrepancies between jet response in data and in simulated events, are applied to the former. The jet energy resolution for simulated jets is degraded slightly to reproduce the resolution in data. The AK4 jet candidates are required to have  $p_T > 30$  GeV and  $|\eta| < 4$ , and to satisfy a stringent set of identification criteria designed to reject spurious detector and reconstruction effects [36]. The jets with  $|\eta| > 2.4$  are referred to as forward jets. The AK8 jets, used to identify and reconstruct Higgs boson candidates, are selected to have  $p_T > 300$  GeV and  $|\eta| < 2.4$ .

A multivariate b tagging algorithm (CSVv2) [37] is used to identify central jets (with  $|\eta| < 2.4$ ) arising from the hadronization of b quarks. Parameters are chosen for the CSVv2 discriminant such that the tagging efficiency for b quark jets is  $\approx 70\%$  while the identification probability averaged over the jet kinematics in  $t\bar{t}$  events is ≈1% for light flavour jets with  $p_T > 30$  GeV.

The Higgs boson candidates are identified using the heavy-flavour content of the AK8 jet. A pruning algorithm [38] is applied that uses the Cambridge-Aachen (CA) algorithm [39] to recluster each AK8 jet starting from all its original constituents and to discard soft and wide-angle radiation inside the jet in each step of the iterative procedure. The procedure defines a pruned-jet mass, computed from the sum of the four-momenta of the constituents that have not been removed by the pruning algorithm, which achieves a better mass resolution. The pruned mass of the jet is then used as a discriminant to reject quark and gluon jets and to select Higgs bosons, by requiring its mass to be within the window of 105–135 GeV. Two subjets are obtained using the soft drop declustering algorithm [40, 41], and these are required to pass the same CSVv2 discriminant threshold used for the AK4 jets.

#### 3 Modelling and simulation

The production and decay of high mass  $B \to Hb$ , with  $H \to b\overline{b}$ , provides a signature with multiple jets rich in heavy-flavour content, and characterized by a highly boosted Higgs boson. The dominant background in this search is from SM events comprised of jets produced through the strong quantum chromodynamic (QCD) interaction, referred to as multijet events. Additional contributions arise from tt events, and minor backgrounds are associated with the production of W or Z bosons in association with jets.

Simulated events are used throughout the analysis to define selection strategy and to determine the expected sensitivity to vector-like quarks. The background from multijet events is estimated using data in control regions. Simulation is also used to cross-check the multijet background prediction and to evaluate its validity. The contributions from other

backgrounds, such as  $t\bar{t}$  events and W or Z boson production in association with jets, are estimated through MC simulation.

Multijet events, as well as electroweak backgrounds from virtual or on-mass shell Z or  $\gamma$ +jets and W+jets production, are simulated at leading order (LO) using the MADGRAPH5 aMC@NLO 2.2.2 generator [42], interfaced to PYTHIA 8.2 [43] with the CUETP8M1 [44, 45] underlying-event tune for parton-shower simulation and evolution. The background  $t\bar{t}$  events are generated using POWHEG v2 at next-to-leading order (NLO)  $[46-49]$ , also interfaced to PYTHIA. The mass of the top quark is set to 172.5 GeV, and the cross section is calculated at next-to-next-to-leading order (NNLO) in perturbative QCD using a next-to-next-to-leading-logarithmic (NNLL) soft-gluon approximation (NNLO+NNLL) in the Top++ 2.0 program [50]. The cross sections for Z or  $\gamma$ +jets and W+jets processes are calculated at NNLO using the fewz MC program [51].

The B  $\rightarrow$  Hb  $\rightarrow$  bbb events are simulated at LO, modelled using the universal FeynRules output [52, 53] and the MC generator MadGraph5 amc@nlo, interfaced to pythia 8 for parton-shower simulation. Several mass hypotheses are considered for signals in the range  $700 < m<sub>B</sub> < 1800 \,\text{GeV}$ , in steps of  $100 \,\text{GeV}$  for total decay widths of 1 GeV, representing the narrow-width categories. Signal events for B quarks with large widths (10, 20, or 30\%) of the mass hypothesis) are also generated in the same mass range. All B quarks are generated with left-handed chirality, but the effect on the kinematic distributions of only considering one chirality is found to be negligible. Interference between the signal and the SM background is negligible.

Simulations using LO and NLO calculations, respectively, use the LO and NLO NNPDF3.0 [54] sets of parton distribution functions (PDFs). All signal and background events are processed using GEANT  $4\left[55\right]$  to provide a full simulation of the CMS detector. The generated events are also reweighted to account for the dependence of the reconstruction efficiency on the number of pileup interactions in the collisions.

#### 4 Interpretation framework

The total cross section for the single production and decay of a B quark with final state X can be written as:

$$
\sigma(C_1, C_2, m, \Gamma_B, X) = C_1^2 C_2^2 \hat{\sigma}_{AW}(m, \Gamma_B), \qquad (4.1)
$$

where  $C_1$  and  $C_2$  are the production and decay couplings corresponding to the interactions through which a B quark is produced and decays, and  $\hat{\sigma}_{AW}$  is the reduced cross section for a resonance of arbitrary width (AW). This width can be written as  $\Gamma_{\rm B} = \Gamma(C_i, m_{\rm B}, m_{\rm decays}),$ as it depends on the B quark mass, on the masses of all its decay products, and on its couplings to all decay channels,  $C_i$ .

Equation (4.1) is valid in all width regimes. However, when  $\Gamma_B/m_B$  approaches zero, it is possible to factorize production and decay and to write the cross section as:

$$
\sigma(C_1, C_2, m_\text{B}, \Gamma_\text{B}) = \sigma_{\text{prod}}(C_1, m_\text{B}) \mathcal{B}_{\text{B}\to\text{X}} = C_1^2 \hat{\sigma}_{\text{NWA}}(m_\text{B}) \mathcal{B}_{\text{B}\to\text{X}},\tag{4.2}
$$

where  $C_1$  is the B production coupling, and information for the parameters  $C_2$  and  $\Gamma_\text{B}$  are included in the branching fraction for the specific B quark decay, in this case  $\mathcal{B}_{B\to X}$ , while  $\hat{\sigma}_{\text{NWA}}(m_{\text{B}})$  is the reduced cross section in the narrow-width approximation (NWA).

Our assumptions have the B quark decaying into Hb, Zb, and Wt with branching fractions that are specified in the model. The couplings of the B quark to SM bosons and quarks can be parametrized as:  $c_Z = e/(2c_w s_w \kappa_Z)$ ,  $c_W = e/(\sqrt{2s_w \kappa_W})$ , and  $c_H =$  $(m_{\text{BKH}})/v$ , where e is the electric charge of the proton,  $v = 246 \text{ GeV}$  is the vacuumexpectation value for the field of the Higgs boson,  $c_w$  and  $s_w$  are the cosine and sine of the weak mixing angle  $\theta_W$ , and  $\kappa$  is a coupling strength that can be fixed to obtain the desired width. Numerically,  $e/(2c_w s_w) = 0.370$ , and  $c_w = e/(\sqrt{2s_w}) = 0.458$ . For the process under consideration, we can set  $C_1 \equiv c_Z$  and  $C_2 \equiv c_H$ .

The  $\kappa$  values can be related to the mixing angle between the vector-like B quark and the b quark [56], and correspond to left- and right-handed couplings, which are the dominant chiralities for a singlet or part of a doublet B quark, respectively. For small values of  $\kappa$ , corresponding to the NWA regime, the following relations hold to excellent approximation: for a B singlet  $\kappa_Z \approx \kappa_H \approx \kappa_W \approx \kappa$ , while for a (T,B) doublet (where T is a vector-like quark with electrical charge 2/3) with no vector-like top quark Yukawa coupling,  $\kappa_Z \approx \kappa_H \approx \kappa$ , and  $\kappa_{\rm W} = 0$ . By imposing these relations among the  $\kappa$  values, and fixing the  $\Gamma_{\rm B}/m_{\rm B}$  ratio to 1%,  $\kappa$  is  $\approx 0.1$  in the whole range of explored masses. Table 1 provides the values for  $\hat{\sigma}_{\text{NWA}}$  and the physical cross sections in the NWA for the pp  $\rightarrow$  Bbq process. The CTEQ6L PDF set  $[57]$  is used in this calculation.

To interpret the results in a model-independent way, the mechanism through which the B quarks achieve large widths is not specified, and  $\Gamma_B$  is considered as a free parameter. The relations among the  $\kappa_X$  (with X = W, Z, H), corresponding to the NWA limit ( $\kappa_Z$  =  $\kappa_{\rm H} = \kappa_{\rm W} = \kappa$ ), are imposed for the large-width regime. With this assumption, the total width  $\Gamma_B$  is always proportional to  $\kappa^2$ , and therefore  $\kappa$  can be chosen to obtain a specific  $\Gamma_{\rm B}/m_{\rm B}$  ratio. However, with the assumption relaxed, in a simplified model, new physics can be invoked to generate the required couplings.

Table 2 reports the cross sections integrated over the phase space of q and b, the particles produced in association with the B quark (see figure 1), for fixed values of  $\Gamma_B/m_B$ , with configurations of  $\kappa$  corresponding to singlet ( $\sigma_S$ ) and doublet ( $\sigma_D$ ) representations. Given the yields for a doublet in the Zb and Hb decay modes, these couplings at fixed width are larger than for singlets, and as a consequence  $\sigma_{\text{D}} > \sigma_{\text{S}}$ .

#### 5 Event selection

This analysis searches for a Higgs boson and a bottom quark arising from the decay of a B quark, and the decay of the Higgs boson into a pair of b quarks. An additional light-flavour quark, resulting from the production mechanism and produced in the forward direction (see figure 1), is also present. For values of  $m<sub>B</sub>$  much larger than the Higgs boson mass, the decay products of the B quark are expected to have large  $p<sub>T</sub>$ . The two b quarks originating from the Higgs boson tend therefore to emerge very close to each other in  $\eta$ - $\phi$ space, resulting in a single large jet.

			Singlet model				Doublet model				
$m_{\rm B}$ (GeV)	$\hat{\sigma}_{\text{NWA}}$ (pb)		$\kappa$				$B_{B\to Wt}$ $B_{B\to Zb}$ $B_{B\to Hb}$ $\sigma_{NWA}$ (pb)	$\kappa$			$\mathcal{B}_{B\to Zb}$ $\mathcal{B}_{B\to Hb}$ $\sigma_{NWA}$ (pb)
700	$31.30_{\ \ -20\%}^{\ +28\%}$		0.18	0.466	0.271	0.263	0.1631	0.25	0.499	0.501	0.5720
800	$21.50$ $^{+29\%}_{-21\%}$		0.16	0.474	0.276	0.260	0.0830		$0.22 \quad 0.499$	0.501	0.3003
900	$15.10_{\phantom{0}-21\%}^{\phantom{0}+30\%}$		0.14	0.489	0.263	0.258	0.0451	0.19	0.500	0.500	0.1666
1000	$10.80$ $^{+31\%}_{-23\%}$		0.13	0.483	0.261	0.256	0.0257	0.17	0.500	0.500	0.0962
1100	$7.85$ $^{+32\%}_{-22\%}$		0.11	0.486	0.259	0.255	0.0153		$0.16$ $0.500$	0.500	0.0580
1200	$5.77$ $^{+33\%}_{-23\%}$		0.10	0.489	0.257	0.254	0.0094	0.15	0.500	0.500	0.0358
1300	$4.29_{\phantom{0}-23\%}^{\phantom{0}+34\%}$		0.10	0.490	0.256	0.254	0.0059	0.13	0.500	0.500	0.0227
1400	$3.23_{\phantom{0}-23\%}^{\phantom{0}+34\%}$		0.09	0.492	0.255	0.253	0.0038		$0.12 \quad 0.500$	0.500	0.0147
1500	$2.45\text{ }^{+35\%}_{-25\%}$		0.08	0.493	0.254	0.253	0.0025	0.12	0.500	0.500	0.0097
1600	$1.86$ $^{+36\%}_{-24\%}$		0.08	0.494	0.254	0.252	0.0017	0.11	0.500	0.500	0.0065
1700	$1.44$ $^{+37\%}_{-24\%}$		0.07	0.494	0.254	0.252	0.0011	0.10	0.500	0.500	0.0044
1800	1.11	$^{+37 \%}_{-25 \%}$	0.07	0.495	0.253	0.252	0.0008	0.10	0.500	0.500	0.0031

**Table 1.** Cross sections for pp  $\rightarrow$  Bbq, with the ratio  $\Gamma_B/m_B$  fixed to 1% (NWA). The couplings and branching fractions in simplified models are calculated using the equations in the text. The uncertainties in the production cross sections correspond to the halving and doubling of the QCD renormalization and factorization scales.

	$\Gamma_{\rm B}/m_{\rm B}=10\%$				$\Gamma_{\rm B}/m_{\rm B}=20\%$		$\Gamma_{\rm B}/m_{\rm B}=30\%$		
$m_{\rm B}$ (GeV)	$\tilde{\sigma}_{AW}(pb)$	$\sigma_{\rm S}$ (fb) $(\kappa)$	$\sigma_{\rm D}$ (fb) $(\kappa)$	$\tilde{\sigma}_{AW}(pb)$	$\sigma_{\rm S}$ (fb) $(\kappa)$	$\sigma_{\rm D}$ (fb) $(\kappa)$	$\tilde{\sigma}_{AW}(pb)$	$\sigma_{\rm S}$ (fb) $(\kappa)$	$\sigma_{\rm D}$ (fb) $(\kappa)$
700	3.01	400(0.588)	1378 (0.8010)	1.43	759 (0.832)	2616(1.130)	0.899		$1074(1.020)$ 3703 $(1.390)$
800	2.10	203(0.508)	726 (0.699)	1.00	386 (0.719)	1377 (0.9880)	0.634	552 (0.880)	1968(1.210)
900	1.51	111(0.448)	406(0.619)	0.719	212 (0.633)	775 (0.876)	0.454	301 (0.776)	1101(1.070)
1000	1.09	63.7(0.401)	237(0.556)	0.523	122(0.567)	453 (0.787)	0.331	174 (0.694)	647 (0.964)
1100	0.807	38.2(0.363)	144(0.505)	0.386	73.2(0.513)	276 (0.714)	0.246	105(0.628)	394 (0.875)
1200	0.601	23.6(0.331)	89.7 (0.463)	0.290	45.5(0.468)	173 (0.654)	0.185	65.2(0.574)	248 (0.801)
1300	0.451	14.9(0.305)	57.1 (0.427)	0.220	29.0(0.431)	111(0.603)	0.141	41.9(0.528)	160(0.739)
1400	0.342	9.70(0.283)	37.2 (0.396)	0.167	18.9(0.400)	72.9(0.560)	0.108	27.5(0.489)	106(0.686)
1500	0.262	6.42(0.263)	24.9(0.369)	0.129	12.6(0.372)	48.9(0.522)	0.0836	18.4(0.456)	71.3(0.640)
1600	0.203	4.34(0.246)	16.9(0.346)	0.101	8.61(0.349)	33.5(0.489)	0.0651	12.5(0.427)	48.7(0.599)
1700	0.158	2.99(0.232)	11.6(0.326)	0.0788	5.94(0.328)	23.2(0.460)	0.0514	8.71(0.401)	34.0(0.564)
1800	0.124	2.08(0.219)	8.13(0.307)	0.0621	4.16(0.309)	16.3(0.435)	0.0408	6.14(0.379)	24.0(0.532)

**Table 2.** Cross sections for  $pp \rightarrow Bbg$  for three values of the  $\Gamma_B/m_B$  ratio. The conditions assume that singlets and doublets have  $\kappa_W = \kappa_Z = \kappa_H \equiv \kappa$ ,  $\kappa_W = 0$  and  $\kappa_Z = \kappa_H \equiv \kappa$ , respectively. For each  $\Gamma_{\rm B}/m_{\rm B}$ , we provide the values of  $\tilde{\sigma}_{\rm AW}$  and of the physical cross sections for both the singlet and doublet models,  $\sigma_S$  and  $\sigma_D$  respectively. The uncertainties in the production cross sections correspond to the halving and doubling of the QCD renormalization and factorization scales. The values of  $\kappa$  are listed in the parentheses.



Figure 2. The b-tagged subjet multiplicity of AK8 jets in events passing preselection criteria. The lower panel shows the ratio of data to the MC background prediction. The normalization of the contributions from signals at  $m<sub>B</sub> = 1$  and 1.8 TeV is multiplied by a factor of 500. Background events are normalized to data. Only the statistical uncertainties are taken into consideration here, and they are too small to be visible.

The data are collected through an online selection (trigger) based on jet activity  $H<sub>T</sub>$ , defined as the scalar  $p_T$  sum of all AK4 jets with  $p_T > 30$  GeV and  $|\eta| < 3$ . The jet activity threshold for this trigger is 900 GeV. Collisions containing at least one jet reconstructed through the HLT system with  $p_T > 450 \text{ GeV}$  are also selected, to increase the HLT efficiency. At the analysis level,  $H_T$  is recalculated using AK4 jets with  $p_T > 50$  GeV and  $|\eta| < 2.4$ , and  $H_{\rm T} > 950$  GeV is required. This offline selection corresponds to a trigger efficiency in excess of 87%.

Events are preselected if they contain three or more AK4 jets with  $p_T > 30$  GeV and  $|\eta|$  < 4, among which there must be at least one b-tagged jet with  $|\eta|$  < 2.4. A veto is applied to events with one or more leptons to ensure that the selection criteria do not overlap with those used for searches for the B quark in leptonic final states. Selected events are further required to have at least one large Higgs-tagged AK8 jet, fulfilling the Higgs boson tagging requirements as described in section 2. The Higgs boson tagging efficiency is 10–20%, depending on the value of  $m<sub>B</sub>$ . Figure 2 compares to data the b-tagged subjet multiplicity expected for simulated background and for signal processes.

The B quark is reconstructed from the Higgs jet candidate along with a nonoverlapping b-tagged jet. The b quark from B quark decay is usually highly energetic  $(p_T > 200 \,\text{GeV})$ , thus the b jet with the highest  $p<sub>T</sub>$  is chosen, and this reduces significantly the combinatorial background. Furthermore, to reduce overlaps with the decay products of the Higgs boson, a condition is applied on the distance between the two objects in  $(\eta, \phi)$ , requiring  $\Delta R(\text{b}, \text{H}) > 1.2.$ 



Figure 3. Multiplicity of forward jets before event categorization. The normalization of the signal contributions is multiplied by a factor of 500. All the contributions to background are obtained from Monte Carlo simulation and are normalized to data. The lower panel shows the ratio of data to background. We show only the statistical uncertainties.

To further reduce the multijet background and the contamination from gluon-like jets,  $H<sub>T</sub>$  is required to be in excess of 950 GeV for smaller mass values of 700  $< m<sub>B</sub> < 1500$  GeV, while for  $1500 < m<sub>B</sub> < 1800$  GeV, a trigger with a threshold of  $H<sub>T</sub> > 1250$  GeV is chosen. In what follows, we refer to the former as the "low-mass analysis" and to the latter as the "high-mass analysis".

The signal to background discrimination is enhanced by exploiting the distinctive presence of a forward jet. Events are therefore separated into categories based on the forward-jet multiplicity. A high-purity category is obtained by requiring at least one forward jet. A second category that contains a large fraction of events from both signal and background, is defined requiring no forward jets. The forward-jet multiplicity expected for background and signal events after preselection is compared to data in figure 3. After all the selections are implemented, we reach signal efficiencies ranging from 2% or less at low masses, to larger values at larger  $m_B$ , as a result of the optimization of the analysis for highly-boosted topologies. The disagreement between data and simulation at large forward-jet multiplicities does not affect the analysis, as the background contribution in the signal region is estimated from data. Moreover, the effect on the measurement is negligible since the majority of vector-like B quark events contain less than 2 forward jets, for which the simulated and observed yields are consistent after preselection.

#### 6 Signal extraction

A potential signal would manifest itself as a localized excess over the expected background in the spectrum of the reconstructed mass  $m<sub>bH</sub>$ . A binned maximum likelihood fit is performed to the  $m<sub>bH</sub>$  distribution to extract a signal, exploiting the characteristic structure of the reconstructed B quark mass spectrum.

Multijet events constitute the dominant source of background in this search. An additional contribution of  $5-7\%$  arises from  $t\bar{t}$  events. To reduce the dependence of the maximum-likelihood fit on the modelling of the multijet background in simulation, the contribution from this background is obtained from data. The procedure we use to estimate the yield of such events in the signal region is referred to as the "ABCD method" (discussed below), but its dependence on  $m<sub>bH</sub>$  is taken from a background-enriched control region in data. A minor contribution  $(\approx 1\%)$  to the background arises from other SM sources, such as Z+jets and W+jets events. Both  $t\bar{t}$  events and these minor backgrounds are estimated from simulation. The normalization of multijet events in the signal region is estimated using three data control regions, enriched in background events. These regions, in addition to the one enriched in signal events, are sampled in a two-dimensional phase space defined by two variables: the b-tagged subjet multiplicity of the Higgs jet and its reconstructed mass, mJ. From a check on the simulation, the number of b-tagged subjets is not correlated with  $m_1$ . The four regions used to define the ABCD method are: (i) region A, with two b-tagged subjets, and  $105 < m<sub>J</sub> < 135$  GeV, (ii) region B, with two b-tagged subjets, and 75  $\lt m_l \lt 105$  GeV or  $m_l > 135$  GeV, (iii) region C, with one b-tagged subjet, and  $105 < m<sub>J</sub> < 135$  GeV, and (iv) region D, with one b-tagged subjet, and  $75 < m_{\rm J} < 105 \,\text{GeV}$  or  $> 135 \,\text{GeV}$ .

Region A is the signal region, defined by the selection criteria described in the previous section. The multijet background yield in the signal region is obtained from regions B, C, and D, which are background enriched. Assuming that the b-tagged subjet multiplicity and the Higgs boson mass are uncorrelated, the number of background events in the four regions follows the relationship:

$$
N_{\rm A}/N_{\rm C} = N_{\rm B}/N_{\rm D},\tag{6.1}
$$

where  $N_A$ ,  $N_B$ ,  $N_C$ , and  $N_D$  are the yields in regions A, B, C, and D, respectively. Thus, the number of background events in the signal region A is given by:

$$
N_{\rm A} = N_{\rm C} N_{\rm B} / N_{\rm D},\tag{6.2}
$$

after subtracting the  $t\bar{t}$  contribution predicted in the MC simulation. The contributions from Z+jets and W+jets backgrounds are not subtracted as they are negligible.

The  $m<sub>B</sub>$  distribution of the multijet background in the signal region is estimated from the  $m_{\text{bH}}$  distribution in region C, since the reconstructed  $m_{\text{bH}}$  spectrum is not expected to be correlated with the b jet multiplicity. The compatibility of the distributions in regions A and C is verified using simulated multijet events, and cross-checked in data.

In addition, the method is validated using a signal-depleted region from sidebands at large mass. Here, two regions  $(A'$  and  $C'$ ) are defined, similar to A and C in the mass region  $135 < m<sub>J</sub> < 165$  GeV. Two control regions (B' and D') are defined requiring  $75 < m<sub>J</sub> < 105 \text{ GeV}$  or  $m<sub>J</sub> > 165 \text{ GeV}$ , respectively, with 2 or 1 b-tagged subjets. The background distribution estimated in region A', using the method described above, agrees

with the observed data in region A'. The systematic uncertainty in the normalization of the multijet background is taken to be equal to the observed difference between the predicted and the measured yields in region A'. It amounts to 10% in the high purity category, and 5% in the category with no forward jets.

#### 7 Systematic uncertainties

The systematic effect of each source of uncertainty is evaluated by propagating the uncertainty in the input parameters to the reconstructed B quark mass distribution and to the event yield. Then, the uncertainties in the event yield and in the  $m<sub>bH</sub>$  distribution for signal and background processes are taken into account as "nuisance" parameters that are integrated over in the statistical process of inferring the resultant parameters.

The statistical uncertainties in the background estimate of multijet production from control samples in data are propagated to  $m<sub>bH</sub>$  in the signal region by changing the observed event yields in regions B and D, up and down by one standard deviation, and recalculating the expected distribution in the signal region. As the expected multijet distribution in  $m<sub>bH</sub>$  is estimated from region C, its statistical uncertainty in this region is considered in the signal extraction. In addition to the normalization, this uncertainty affects the distribution of the background  $m<sub>bH</sub>$  in the signal region. Therefore, a systematic uncertainty in the estimated shape of this distribution, arising from the limited number of events in the observed  $m<sub>hH</sub>$  spectrum in region C, was derived by allowing the content of each bin to fluctuate independently according to Poisson statistics.

An additional systematic uncertainty in the estimated multijet background is obtained from the difference between the observed and predicted yields in the check, in the validation step that uses large-mass sideband regions, described in section  $6$ , and corresponds to  $\approx$  5–10%.

The systematic uncertainties from the limited number of simulated events and background estimates from simulation are also included by fluctuating each bin of the  $m<sub>bH</sub>$  distribution independently, according to Poisson statistics.

Additional systematic uncertainties in simulated signal and background distributions originate from the corrections applied to rescale simulated distributions to data. Other such uncertainties are listed below. An uncertainty of 2.5% [58] in the measured integrated luminosity is used just to account for the total event yields.

The corrections to account for the difference between the b tagging efficiency measured in data and in simulation are changed up and down by their uncertainties in both AK4 jets and subjets. The reconstructed four-momenta of the AK4 and AK8 jets are also shifted by  $\pm 1$  standard deviation in the jet energy scale and resolution, and propagated to  $m_{\text{bH}}$ . In addition, the pruned mass scale and resolution of the Higgs-tagged jet are changed within their uncertainties, affecting the  $m_{\text{bH}}$  spectrum by 0.5–5.5%.

All simulated events are weighted to match the distribution of pileup interactions. The corresponding uncertainty is obtained by changing the total inelastic cross section by  $\pm 4.6\%$ , which is used to calculate the pileup distribution in data. Scale factors are applied to account for differences between the trigger efficiency measured in data and in

Source	Effect
Luminosity	$2.5\%$
b tagging efficiency	$0 - 9\%$
Misidentification efficiency	$0 - 2\%$
Pileup modelling	$0 - 12\%$
Trigger	${<}0.5\%$
<b>PDF</b>	$1.0 - 4.5\%$
$\mu_{\rm R}$ and $\mu_{\rm F}$	$15 - 25\%$
Jet energy scale	$1 - 7\%$
Jet energy resolution	$1.0 - 1.5\%$
Jet mass scale.	$0 - 5\%$
Jet mass resolution	$0 - 4\%$
MC Statistical accuracy	$1 - 4\%$
Mismodelling of forward jets	$0.5/2.0\%$
Background estimation	$5 - 10\%$

Table 3. Summary of systematic uncertainties in background events. The quantification of the effects quoted in the table reflects the uncertainties in the event yields. All uncertainties are considered in the simulated background events, except the one on background estimation that affects only the data-based estimate of the multijet process. All the systematic uncertainties apply to both categories of forward-jet multiplicity, except for the case of the modelling of the forward jets, where the first entry corresponds to the category with no forward jets, and the second entry to the category with at least one jet in the forward region.

simulated events, with the uncertainties in the scale factors applied as a function of  $H<sub>T</sub>$  and propagated to the  $m<sub>bH</sub>$  distribution.

An additional uncertainty is applied to the simulated signal and backgrounds to account for discrepancies in the modelling of the forward jet multiplicity. The magnitude of this effect is obtained by considering the difference between the event yield in data and in MC, and results in an uncertainty of 0.5% for the category with no forward jets, and 2.0% for the category with at least one jet in the forward region.

The uncertainties from the choice of factorization and renormalization scales,  $\mu_F$  and  $\mu_{\rm R}$ , are taken into account by halving and doubling the nominal values and using the combination of  $\mu_F$  and  $\mu_R$  leading to the maximal change. The resulting uncertainty in signal acceptance is as small as 1.3%, depending on the mass hypothesis. Larger effects (15–25%) are observed in the overall normalization and acceptance in simulated backgrounds. In addition, the uncertainty from the choice of PDF is estimated by reweighting the simulated signal and background events using the NNPDF3.0 [59–61] set of eigenvectors.

A summary of the systematic uncertainties considered in this analysis, along with their effect when propagated to the reconstructed B mass, is presented in table 3.

#### 8 Results

A binned maximum likelihood fit is performed to the  $m<sub>bH</sub>$  distribution in figure 4, where the dominant multijet background is estimated from data, as discussed in section 6. The fitted  $m<sub>hH</sub>$  distributions are presented in figure 5, while the expected yields are listed in table 4 for the backgrounds, and for two signal hypotheses ( $m<sub>bH</sub> = 1000$  and 1800 GeV), together with their observed yields. The observed distributions are consistent with the background-only hypothesis in all the categories. Upper limits are set therefore on the product of the cross section and branching fraction of a B quark decaying to Hb, produced in association with another b quark and a light-flavoured quark, as a function of  $m<sub>bH</sub>$ . Exclusion limits at 95% confidence level (CL) are calculated using a modified frequentist approach and a profile likelihood ratio as test statistic, in an asymptotic approximation  $[62-64]$ . The combination of the two forward-jet multiplicity-based categories increases the sensitivity of the analysis by up to 20% relative to that obtained when only requiring at least one jet in the  $|\eta| > 2.4$ region of the detector.

Systematic uncertainties described in section 7 are treated as nuisance parameters affecting the rate of the expected  $m<sub>bH</sub>$  distribution. Both the uncertainties affecting the normalization, modelled using log-normal priors, and uncertainties in distributions are included in the fit [65].

The observed and expected combined upper limits from the two categories are given in figure 6. Assuming a narrow width, values of  $\sigma$   $\mathcal{B}$ (Hb) between 0.07–1.28 pb are excluded at the 95% confidence level, for masses in the range 700–1800 GeV. Upper limits are compared with the predictions calculated at NLO [53] for both singlet and doublet B quark models, assuming narrow widths and  $\mathcal{B}(Hb) \approx 25\%$ . Figure 6 also shows the observed and expected upper limits on the product of the cross section and branching fraction for B quarks with intrinsic widths fixed to  $\Gamma_B/m_B = 10, 20,$  and 30%. Sensitivities similar to those for negligible widths are observed for exclusion limits that lie between 0.08 and 1.97, 0.11 and 1.32, and 0.10 and 1.22 pb, respectively, for the 10, 20, and 30%  $\Gamma_{\rm B}/m_{\rm B}$  values.

#### 9 Summary

A search has been presented for electroweak production of vector-like B quarks with charge  $-1/3e$ , decaying to a bottom quark and a Higgs boson (H). The analysis uses a data sample corresponding to an integrated luminosity of  $35.9 \text{ fb}^{-1}$ , collected in pp collisions at  $\sqrt{s} = 13 \,\text{TeV}.$ 

No significant deviations are observed relative to the standard model prediction, and upper limits are placed on the product of the cross section and the branching fraction of the B quark.

Expected and observed limits at 95% confidence level vary from 1.20 to 0.07 pb and from 1.28 to 0.07 pb, respectively, for B quark masses in the range considered, which extends from 700 to 1800 GeV. The search is performed under the hypothesis of a singlet or doublet B quark of narrow width decaying to Hb with a branching fraction of approximately 25%. The possibility of having non-negligible resonant widths is also studied. Limits obtained



Figure 4. Distribution in the reconstructed B quark mass, after applying all selections to events with no forward jets (left) and to events with at least one forward jet (right), compared to the background distributions estimated before fitting. The upper and lower plots refer to the lowand high-mass  $m<sub>B</sub>$  analyses, respectively. The expectations for signal MC events are given by the blue histogram lines. The different background contributions are indicated by the colour-filled histograms, and are obtained from Monte Carlo simulation, except for the multijets component, which is derived from data. The grey-hatched error band shows total uncertainties in the background expectation. The ratios of observations to background expectations are given in the lower panels, together with the total uncertainties prior to fitting, indicated by the grey-hatched band.



Figure 5. Distribution in the reconstructed B quark mass after applying all selections to events with no forward jets (left) and to events with at least one forward jet (right), compared to the background distributions estimated after fitting. The upper and lower plots refer to the lowand high- $m_B$  analyses, respectively. The expectations for signal MC events are given by the blue lines. The different background contributions are indicated by the colour-filled histograms, and are obtained from Monte Carlo simulation, except for the multijets component, which is derived from data. The grey-hatched error band shows total uncertainties in the background expectation. The ratios of the observations to background expectations are given in the lower panels, together with the total uncertainties after fitting, indicated by the grey-hatched band.

Category	Source		$700 < m_{\rm B} < 1500 \,\text{GeV}$ $1500 < m_{\rm B} < 1800 \,\text{GeV}$		
	$t\overline{t}$	$394 \pm 46$	$117 \pm 18$		
	$W+jets$	$29 \pm 13$	$10.5 \pm 4.3$		
	$Z + jets$	$43 \pm 15$	$23 \pm 23$		
No forward jets	Multijets	$5416 \pm 60$	$1612 \pm 24$		
	Total background	$5882 \pm 42$	$1762 \pm 26$		
	Observed in data	$5886 \pm 77$	$1753 \pm 42$		
	Expected signal	$7.3 \pm 0.3$	$0.27 \pm 0.01$		
	$t\overline{t}$	$163 \pm 20$	$58 \pm 17$		
	$W+{\rm jets}$	$11.5 \pm 4.2$	$4.3 \pm 1.4$		
	$Z + jets$	$2^{+10}_{-2}$			
$>0$ forward jets	Multijets	$1938 \pm 23$	$549 \pm 10$		
	Total background	$2115 \pm 21$	$612 \pm 15$		
	Observed in data	$2107 \pm 46$	$608 \pm 25$		
	Expected signal	$11.5 \pm 0.3$	$0.51 \pm 0.01$		

**Table 4.** Observed and expected fitted number of events in the signal ranges of 700  $< m<sub>B</sub> < 1500$ and  $1500 < m<sub>B</sub> < 1800 \text{ GeV}$ , and expected signal at  $m<sub>B</sub> = 1000$  and 1800 GeV. The multijet background is obtained from data, while the yields for the other sources of background are obtained from MC simulation. The combined statistical and systematic uncertainties correspond to the quadrature of the statistical and systematic uncertainties.

on the production of B quarks with widths of 10, 20, and 30% of the resonance mass are comparable to those found for the narrow-width approximation. This search extends existing knowledge on vector-like quarks, by interpreting the results in a new theoretical framework with non-negligible resonance widths, and investigating the final state with a bottom quark and a Higgs boson for the first time.

# Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC and the CMS detector provided by the following funding agencies: BMWFW and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COL-CIENCIAS (Colombia); MSES and CSF (Croatia); RPF (Cyprus); SENESCYT (Ecuador); MoER, ERC IUT, and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece);



Figure 6. The median observed and expected 95% CL upper limits on the product of the B quark production cross section and branching fraction as a function of the signal mass, assuming narrowwidth resonances (upper-left) and widths of 10 (upper-right), 20 (lower-left), and  $30\%$  (lower-right) of the resonance mass for the B quark. The results are shown for the combination of 0 and  $>0$ forward-jet categories. The continuous red curves correspond to the theoretical expectations for singlet and doublet models.

OTKA and NIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, RosAtom, RAS, RFBR and RAEP (Russia); MESTD (Serbia); SEIDI, CPAN, PCTI and FEDER (Spain); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR, and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU and SFFR (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract No. 675440 (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science - EOS" - be.h project n. 30820817; the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Council of Science and Industrial Research, India; the HOMING PLUS program of the Foundation for Polish Science, cofinanced from European Union, Regional Development Fund, the Mobility Plus program of the Ministry of Science and Higher Education, the National Science Center (Poland), contracts Harmonia 2014/14/M/ST2/00428, Opus 2014/13/B/ST2/02543, 2014/15/B/ST2/03998, and 2015/19/B/ST2/02861, Sonata-bis 2012/07/E/ST2/01406; the National Priorities Research Program by Qatar National Research Fund; the Programa Severo Ochoa del Principado de Asturias; the Thalis and Aristeia programs cofinanced by EU-ESF and the Greek NSRF; the Rachadapisek Sompot Fund for Postdoctoral Fellowship, Chulalongkorn University and the Chulalongkorn Academic into Its 2nd Century Project Advancement Project (Thailand); the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License [\(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

#### References

- [1] ATLAS collaboration, Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2012.08.020) **B 716** (2012) 1 [[arXiv:1207.7214](https://arxiv.org/abs/1207.7214)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1207.7214)].
- [2] CMS collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2012.08.021) **B** 716 (2012) 30  $\left[\text{arXiv:1207.7235}\right]$  $\left[\text{arXiv:1207.7235}\right]$  $\left[\text{arXiv:1207.7235}\right]$   $\left[\text{INSPIRE}\right]$  $\left[\text{INSPIRE}\right]$  $\left[\text{INSPIRE}\right]$ .
- [3] ATLAS and CMS collaborations, Combined measurement of the Higgs boson mass in pp collisions at  $\sqrt{s}$  = 7 and 8 TeV with the ATLAS and CMS experiments, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.114.191803) 114 [\(2015\) 191803](https://doi.org/10.1103/PhysRevLett.114.191803) [[arXiv:1503.07589](https://arxiv.org/abs/1503.07589)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1503.07589)].
- [4] J.A. Aguilar-Saavedra, R. Benbrik, S. Heinemeyer and M. Pérez-Victoria, Handbook of vectorlike quarks: mixing and single production, Phys. Rev. D 88 [\(2013\) 094010](https://doi.org/10.1103/PhysRevD.88.094010)  $\vert$ [arXiv:1306.0572](https://arxiv.org/abs/1306.0572) $\vert$  [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1306.0572)].
- [5] M. Schmaltz and D. Tucker-Smith, *Little Higgs review, [Ann. Rev. Nucl. Part. Sci.](https://doi.org/10.1146/annurev.nucl.55.090704.151502)* 55 (2005) [229](https://doi.org/10.1146/annurev.nucl.55.090704.151502) [[hep-ph/0502182](https://arxiv.org/abs/hep-ph/0502182)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+hep-ph/0502182)].
- [6] D.B. Kaplan and H. Georgi,  $SU(2) \times U(1)$  breaking by vacuum misalignment, [Phys. Lett.](https://doi.org/10.1016/0370-2693(84)91177-8) **B** 136 [\(1984\) 183](https://doi.org/10.1016/0370-2693(84)91177-8) [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22Phys.Lett.,B136,183%22)].
- [7] H. Georgi, D.B. Kaplan and P. Galison, *Calculation of the composite Higgs mass, [Phys. Lett.](https://doi.org/10.1016/0370-2693(84)90823-2)* B 143 [\(1984\) 152](https://doi.org/10.1016/0370-2693(84)90823-2) [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22Phys.Lett.,B143,152%22)].
- [8] M.J. Dugan, H. Georgi and D.B. Kaplan, Anatomy of a composite Higgs model, [Nucl. Phys.](https://doi.org/10.1016/0550-3213(85)90221-4) B 254 [\(1985\) 299](https://doi.org/10.1016/0550-3213(85)90221-4) [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22Nucl.Phys.,B254,299%22)].
- [9] J.A. Aguilar-Saavedra, Identifying top partners at LHC, JHEP 11 [\(2009\) 030](https://doi.org/10.1088/1126-6708/2009/11/030) [[arXiv:0907.3155](https://arxiv.org/abs/0907.3155)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:0907.3155)].
- [10] A. De Simone, O. Matsedonskyi, R. Rattazzi and A. Wulzer, A first top partner hunter's guide, JHEP 04 [\(2013\) 004](https://doi.org/10.1007/JHEP04(2013)004) [[arXiv:1211.5663](https://arxiv.org/abs/1211.5663)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1211.5663)].
- [11] M. Buchkremer, G. Cacciapaglia, A. Deandrea and L. Panizzi, Model independent framework for searches of top partners, [Nucl. Phys.](https://doi.org/10.1016/j.nuclphysb.2013.08.010)  $\bf{B} 876$  (2013) 376  $\left[$ [arXiv:1305.4172](https://arxiv.org/abs/1305.4172) $\right]$  [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1305.4172)].
- [12] G.D. Kribs, T. Plehn, M. Spannowsky and T.M.P. Tait, Four generations and Higgs physics, Phys. Rev. **D 76** [\(2007\) 075016](https://doi.org/10.1103/PhysRevD.76.075016) [[arXiv:0706.3718](https://arxiv.org/abs/0706.3718)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:0706.3718)].
- [13] S. Banerjee, M. Frank and S.K. Rai, Higgs data confronts sequential fourth generation fermions in the Higgs triplet model, Phys. Rev.  $\bf{D}$  89 [\(2014\) 075005](https://doi.org/10.1103/PhysRevD.89.075005) [[arXiv:1312.4249](https://arxiv.org/abs/1312.4249)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1312.4249)].
- [14] ATLAS and CMS collaborations, Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at  $\sqrt{s} = 7$  and 8 TeV, JHEP 08 [\(2016\) 045](https://doi.org/10.1007/JHEP08(2016)045) [[arXiv:1606.02266](https://arxiv.org/abs/1606.02266)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1606.02266)].
- [15] O. Eberhardt et al., Joint analysis of Higgs decays and electroweak precision observables in the Standard Model with a sequential fourth generation, Phys. Rev. D 86 [\(2012\) 013011](https://doi.org/10.1103/PhysRevD.86.013011) [[arXiv:1204.3872](https://arxiv.org/abs/1204.3872)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1204.3872)].
- [16] CMS collaboration, *Searches for Higgs bosons in pp collisions at*  $\sqrt{s} = 7$  and 8 TeV in the context of four-generation and fermiophobic models, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2013.06.043) B 725 (2013) 36 [[arXiv:1302.1764](https://arxiv.org/abs/1302.1764)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1302.1764)].
- [17] ATLAS collaboration, Search for vector-like B quarks in events with one isolated lepton, missing transverse momentum and jets at  $\sqrt{s} = 8$  TeV with the ATLAS detector, [Phys. Rev.](https://doi.org/10.1103/PhysRevD.91.112011) D 91 [\(2015\) 112011](https://doi.org/10.1103/PhysRevD.91.112011) [[arXiv:1503.05425](https://arxiv.org/abs/1503.05425)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1503.05425)].
- [18] ATLAS collaboration, Search for pair production of a new heavy quark that decays into a W boson and a light quark in pp collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector, [Phys. Rev.](https://doi.org/10.1103/PhysRevD.92.112007) D 92 [\(2015\) 112007](https://doi.org/10.1103/PhysRevD.92.112007) [[arXiv:1509.04261](https://arxiv.org/abs/1509.04261)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1509.04261)].
- [19] ATLAS collaboration, Search for pair and single production of new heavy quarks that decay to a Z boson and a third-generation quark in pp collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector, JHEP 11 [\(2014\) 104](https://doi.org/10.1007/JHEP11(2014)104)  $\left[$ [arXiv:1409.5500](https://arxiv.org/abs/1409.5500) $\right]$  [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1409.5500)].
- [20] ATLAS collaboration, Search for single production of a vector-like quark via a heavy gluon in the 4b final state with the ATLAS detector in pp collisions at  $\sqrt{s} = 8 \text{ TeV}$ , [Phys. Lett.](https://doi.org/10.1016/j.physletb.2016.04.061) B 758 [\(2016\) 249](https://doi.org/10.1016/j.physletb.2016.04.061) [[arXiv:1602.06034](https://arxiv.org/abs/1602.06034)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1602.06034)].
- [21] ATLAS collaboration, Search for single production of vector-like quarks decaying into Wb in pp collisions at  $\sqrt{s} = 8 \text{ TeV}$  with the ATLAS detector, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-016-4281-8) C 76 (2016) 442 [[arXiv:1602.05606](https://arxiv.org/abs/1602.05606)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1602.05606)].
- [22] ATLAS collaboration, Search for the production of single vector-like and excited quarks in the Wt final state in pp collisions at  $\sqrt{s} = 8 \text{ TeV}$  with the ATLAS detector, JHEP 02 [\(2016\)](https://doi.org/10.1007/JHEP02(2016)110) [110](https://doi.org/10.1007/JHEP02(2016)110) [[arXiv:1510.02664](https://arxiv.org/abs/1510.02664)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1510.02664)].
- [23] CMS collaboration, *Search for vector-like charge* 2/3 T quarks in proton-proton collisions at  $\sqrt{s} = 8 \text{ TeV},$  Phys. Rev. **D 93** [\(2016\) 012003](https://doi.org/10.1103/PhysRevD.93.012003) [[arXiv:1509.04177](https://arxiv.org/abs/1509.04177)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1509.04177)].
- [24] CMS collaboration, Search for pair-produced vectorlike B quarks in proton-proton collisions  $\alpha t \sqrt{s} = 8 \text{ TeV}, \text{Phys. Rev. } \mathbf{D}$  93 [\(2016\) 112009](https://doi.org/10.1103/PhysRevD.93.112009) [[arXiv:1507.07129](https://arxiv.org/abs/1507.07129)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1507.07129)].
- [25] CMS collaboration, Search for top-quark partners with charge 5/3 in the same-sign dilepton final state, [Phys. Rev. Lett.](https://doi.org/10.1103/PhysRevLett.112.171801) 112 (2014) 171801 [[arXiv:1312.2391](https://arxiv.org/abs/1312.2391)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1312.2391)].
- [26] CMS collaboration, Search for single production of vector-like quarks decaying to a Z boson and a top or a bottom quark in proton-proton collisions at  $\sqrt{s} = 13 \text{ TeV}$ , JHEP 05 [\(2017\)](https://doi.org/10.1007/JHEP05(2017)029) [029](https://doi.org/10.1007/JHEP05(2017)029) [[arXiv:1701.07409](https://arxiv.org/abs/1701.07409)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1701.07409)].
- [27] CMS collaboration, Search for single production of a heavy vector-like T quark decaying to a Higgs boson and a top quark with a lepton and jets in the final state, [Phys. Lett.](https://doi.org/10.1016/j.physletb.2017.05.019) B 771 [\(2017\) 80](https://doi.org/10.1016/j.physletb.2017.05.019) [[arXiv:1612.00999](https://arxiv.org/abs/1612.00999)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1612.00999)].
- [28] CMS collaboration, Search for electroweak production of a vector-like quark decaying to a top quark and a Higgs boson using boosted topologies in fully hadronic final states, [JHEP](https://doi.org/10.1007/JHEP04(2017)136) 04 [\(2017\) 136](https://doi.org/10.1007/JHEP04(2017)136) [[arXiv:1612.05336](https://arxiv.org/abs/1612.05336)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1612.05336)].
- [29] CMS collaboration, Search for single production of vector-like quarks decaying into a b quark and a W boson in proton-proton collisions at  $\sqrt{s} = 13 \text{ TeV}$ , [Phys. Lett.](https://doi.org/10.1016/j.physletb.2017.07.022) **B 772** (2017) 634 [[arXiv:1701.08328](https://arxiv.org/abs/1701.08328)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1701.08328)].
- [30] CMS collaboration, The CMS experiment at the CERN LHC, 2008 JINST 3 [S08004](https://doi.org/10.1088/1748-0221/3/08/S08004) [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22JINST,3,S08004%22)].
- [31] CMS collaboration, The CMS trigger system, 2017 JINST 12 [P01020](https://doi.org/10.1088/1748-0221/12/01/P01020) [[arXiv:1609.02366](https://arxiv.org/abs/1609.02366)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1609.02366)].
- [32] CMS collaboration, Particle-flow reconstruction and global event description with the CMS detector, 2017 JINST 12 [P10003](https://doi.org/10.1088/1748-0221/12/10/P10003)  $arXiv:1706.04965$  [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1706.04965)].
- [33] M. Cacciari, G.P. Salam and G. Soyez, The anti- $k_t$  jet clustering algorithm, JHEP 04 [\(2008\)](https://doi.org/10.1088/1126-6708/2008/04/063) [063](https://doi.org/10.1088/1126-6708/2008/04/063) [[arXiv:0802.1189](https://arxiv.org/abs/0802.1189)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:0802.1189)].
- [34] CMS collaboration, *Pileup removal algorithms*, [CMS-PAS-JME-14-001](http://cds.cern.ch/record/1751454) [IN[SPIRE](https://inspirehep.net/search?p=find+R+CMS-PAS-JME-14-001)].
- [35] M. Cacciari, G.P. Salam and G. Soyez, FastJet user manual, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-012-1896-2) C 72 (2012) 1896 [[arXiv:1111.6097](https://arxiv.org/abs/1111.6097)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1111.6097)].
- [36] CMS collaboration, Jet algorithms performance in 13 TeV data, [CMS-PAS-JME-16-003](http://cds.cern.ch/record/2256875) [IN[SPIRE](https://inspirehep.net/search?p=find+R+CMS-PAS-JME-16-003)].
- [37] CMS collaboration, *Identification of heavy-flavour jets with the CMS detector in pp* collisions at 13 TeV,  $2018$  JINST 13 [P05011](https://doi.org/10.1088/1748-0221/13/05/P05011) [[arXiv:1712.07158](https://arxiv.org/abs/1712.07158)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1712.07158)].
- [38] S.D. Ellis, C.K. Vermilion and J.R. Walsh, *Techniques for improved heavy particle searches* with jet substructure, Phys. Rev. D 80 [\(2009\) 051501](https://doi.org/10.1103/PhysRevD.80.051501)  $\text{arXiv:}$ 0903.5081  $\text{[insPIRE]}$ .
- [39] S. Catani, Y.L. Dokshitzer, M.H. Seymour and B.R. Webber, *Longitudinally invariant*  $K_t$ clustering algorithms for hadron hadron collisions, [Nucl. Phys.](https://doi.org/10.1016/0550-3213(93)90166-M) B 406 (1993) 187 [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22Nucl.Phys.,B406,187%22)].
- [40] M. Dasgupta, A. Fregoso, S. Marzani and G.P. Salam, Towards an understanding of jet substructure, JHEP 09 [\(2013\) 029](https://doi.org/10.1007/JHEP09(2013)029) [[arXiv:1307.0007](https://arxiv.org/abs/1307.0007)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1307.0007)].
- [41] A.J. Larkoski, S. Marzani, G. Soyez and J. Thaler, Soft drop, JHEP 05 [\(2014\) 146](https://doi.org/10.1007/JHEP05(2014)146) [[arXiv:1402.2657](https://arxiv.org/abs/1402.2657)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1402.2657)].
- [42] J. Alwall et al., The automated computation of tree-level and next-to-leading order differential cross sections and their matching to parton shower simulations, JHEP 07 [\(2014\)](https://doi.org/10.1007/JHEP07(2014)079) [079](https://doi.org/10.1007/JHEP07(2014)079) [[arXiv:1405.0301](https://arxiv.org/abs/1405.0301)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1405.0301)].
- [43] T. Sjöstrand et al., An introduction to PYTHIA 8.2, [Comput. Phys. Commun.](https://doi.org/10.1016/j.cpc.2015.01.024) 191 (2015) [159](https://doi.org/10.1016/j.cpc.2015.01.024) [[arXiv:1410.3012](https://arxiv.org/abs/1410.3012)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1410.3012)].
- [44] P. Skands, S. Carrazza and J. Rojo, Tuning PYTHIA 8.1: the Monash 2013 tune, [Eur. Phys.](https://doi.org/10.1140/epjc/s10052-014-3024-y) J. C 74 [\(2014\) 3024](https://doi.org/10.1140/epjc/s10052-014-3024-y) [[arXiv:1404.5630](https://arxiv.org/abs/1404.5630)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1404.5630)].
- [45] CMS collaboration, Event generator tunes obtained from underlying event and multiparton scattering measurements, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-016-3988-x) C  $76$  (2016) 155 [[arXiv:1512.00815](https://arxiv.org/abs/1512.00815)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1512.00815)].
- [46] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, JHEP 11 [\(2004\) 040](https://doi.org/10.1088/1126-6708/2004/11/040) [[hep-ph/0409146](https://arxiv.org/abs/hep-ph/0409146)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+hep-ph/0409146)].
- [47] S. Frixione, P. Nason and G. Ridolfi, A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction, JHEP  $\overline{09}$  [\(2007\) 126](https://doi.org/10.1088/1126-6708/2007/09/126) [[arXiv:0707.3088](https://arxiv.org/abs/0707.3088)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:0707.3088)].
- [48] S. Frixione, P. Nason and C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method, JHEP 11 [\(2007\) 070](https://doi.org/10.1088/1126-6708/2007/11/070)  $\left[$ [arXiv:0709.2092](https://arxiv.org/abs/0709.2092) $\right]$  [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:0709.2092)].
- [49] S. Alioli, P. Nason, C. Oleari and E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX, JHEP 06 [\(2010\) 043](https://doi.org/10.1007/JHEP06(2010)043) [[arXiv:1002.2581](https://arxiv.org/abs/1002.2581)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1002.2581)].
- [50] M. Czakon and A. Mitov,  $Top++:$  A program for the calculation of the top-pair cross-section at hadron colliders, [Comput. Phys. Commun.](https://doi.org/10.1016/j.cpc.2014.06.021) 185 (2014) 2930 [[arXiv:1112.5675](https://arxiv.org/abs/1112.5675)] [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22Comput.Phys.Commun.,185,2930%22)].
- [51] Y. Li and F. Petriello, Combining QCD and electroweak corrections to dilepton production in FEWZ, Phys. Rev. D 86 [\(2012\) 094034](https://doi.org/10.1103/PhysRevD.86.094034) [[arXiv:1208.5967](https://arxiv.org/abs/1208.5967)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1208.5967)].
- [52] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer and T. Reiter, UFO The Universal FeynRules Output, [Comput. Phys. Commun.](https://doi.org/10.1016/j.cpc.2012.01.022) 183 (2012) 1201 [[arXiv:1108.2040](https://arxiv.org/abs/1108.2040)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1108.2040)].
- [53] O. Matsedonskyi, G. Panico and A. Wulzer, On the interpretation of top partners searches, JHEP 12 [\(2014\) 097](https://doi.org/10.1007/JHEP12(2014)097) [[arXiv:1409.0100](https://arxiv.org/abs/1409.0100)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1409.0100)].
- [54] NNPDF collaboration, R.D. Ball et al., Parton distributions from high-precision collider data, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-017-5199-5) C 77 (2017) 663  $\text{arXiv:1706.00428}$  $\text{arXiv:1706.00428}$  $\text{arXiv:1706.00428}$  [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1706.00428)].
- [55] GEANT4 collaboration, S. Agostinelli et al.,  $GEANT4 a$  simulation toolkit, [Nucl.](https://doi.org/10.1016/S0168-9002(03)01368-8) [Instrum. Meth.](https://doi.org/10.1016/S0168-9002(03)01368-8) A 506 (2003) 250 [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22Nucl.Instrum.Meth.,A506,250%22)].
- [56] C.-Y. Chen, S. Dawson and E. Furlan, *Vectorlike fermions and Higgs effective field theory* revisited, Phys. Rev. D 96 [\(2017\) 015006](https://doi.org/10.1103/PhysRevD.96.015006) [[arXiv:1703.06134](https://arxiv.org/abs/1703.06134)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1703.06134)].
- [57] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P.M. Nadolsky and W.K. Tung, New generation of parton distributions with uncertainties from global QCD analysis, [JHEP](https://doi.org/10.1088/1126-6708/2002/07/012) 07 [\(2002\) 012](https://doi.org/10.1088/1126-6708/2002/07/012) [[hep-ph/0201195](https://arxiv.org/abs/hep-ph/0201195)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+hep-ph/0201195)].
- [58] CMS collaboration, CMS luminosity measurements for the 2016 data taking period, [CMS-PAS-LUM-17-001](http://cds.cern.ch/record/2257069) [IN[SPIRE](https://inspirehep.net/search?p=find+R+CMS-PAS-LUM-17-001)].
- [59] M. Botje et al., The PDF4LHC Working Group Interim Recommendations, [arXiv:1101.0538](https://arxiv.org/abs/1101.0538) [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1101.0538)].
- [60] S. Alekhin et al., The PDF4LHC Working Group Interim Report, [arXiv:1101.0536](https://arxiv.org/abs/1101.0536) [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1101.0536)].
- [61] NNPDF collaboration, R.D. Ball et al., Parton distributions for the LHC Run II, [JHEP](https://doi.org/10.1007/JHEP04(2015)040) 04 [\(2015\) 040](https://doi.org/10.1007/JHEP04(2015)040) [[arXiv:1410.8849](https://arxiv.org/abs/1410.8849)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1410.8849)].
- [62] T. Junk, Confidence level computation for combining searches with small statistics, [Nucl.](https://doi.org/10.1016/S0168-9002(99)00498-2) [Instrum. Meth.](https://doi.org/10.1016/S0168-9002(99)00498-2) A 434 (1999) 435 [[hep-ex/9902006](https://arxiv.org/abs/hep-ex/9902006)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+hep-ex/9902006)].
- [63] A.L. Read, Presentation of search results: The CL<sub>s</sub> technique, J. Phys. **G 28** [\(2002\) 2693](https://doi.org/10.1088/0954-3899/28/10/313) [IN[SPIRE](https://inspirehep.net/search?p=find+J+%22J.Phys.,G28,2693%22)].
- [64] G. Cowan, K. Cranmer, E. Gross and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, [Eur. Phys. J.](https://doi.org/10.1140/epjc/s10052-011-1554-0) C 71 (2011) 1554 [Erratum ibid. C 73 (2013) 2501] [[arXiv:1007.1727](https://arxiv.org/abs/1007.1727)] [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1007.1727)].
- [65] J.S. Conway, Incorporating Nuisance Parameters in Likelihoods for Multisource Spectra, in Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding, CERN,Geneva, Switzerland 17–20 January 2011, pp. 115–120, [arXiv:1103.0354](https://arxiv.org/abs/1103.0354), [IN[SPIRE](https://inspirehep.net/search?p=find+EPRINT+arXiv:1103.0354)].

#### The CMS collaboration

#### Yerevan Physics Institute, Yerevan, Armenia

A.M. Sirunyan, A. Tumasyan

#### Institut für Hochenergiephysik, Wien, Austria

W. Adam, F. Ambrogi, E. Asilar, T. Bergauer, J. Brandstetter, E. Brondolin, M. Dragicevic, J. Erö, A. Escalante Del Valle, M. Flechl, M. Friedl, R. Frühwirth<sup>1</sup>, V.M. Ghete, J. Grossmann, J. Hrubec, M. Jeitler<sup>1</sup>, A. König, N. Krammer, I. Krätschmer, D. Liko, T. Madlener, I. Mikulec, E. Pree, N. Rad, H. Rohringer, J. Schieck<sup>1</sup>, R. Schöfbeck, M. Spanring, D. Spitzbart, A. Taurok, W. Waltenberger, J. Wittmann, C.-E. Wulz<sup>1</sup>, M. Zarucki

#### Institute for Nuclear Problems, Minsk, Belarus

V. Chekhovsky, V. Mossolov, J. Suarez Gonzalez

#### Universiteit Antwerpen, Antwerpen, Belgium

E.A. De Wolf, D. Di Croce, X. Janssen, J. Lauwers, M. Van De Klundert, H. Van Haevermaet, P. Van Mechelen, N. Van Remortel

#### Vrije Universiteit Brussel, Brussel, Belgium

S. Abu Zeid, F. Blekman, J. D'Hondt, I. De Bruyn, J. De Clercq, K. Deroover, G. Flouris, D. Lontkovskyi, S. Lowette, I. Marchesini, S. Moortgat, L. Moreels, Q. Python, K. Skovpen, S. Tavernier, W. Van Doninck, P. Van Mulders, I. Van Parijs

#### Université Libre de Bruxelles, Bruxelles, Belgium

D. Beghin, B. Bilin, H. Brun, B. Clerbaux, G. De Lentdecker, H. Delannoy, B. Dorney, G. Fasanella, L. Favart, R. Goldouzian, A. Grebenyuk, A.K. Kalsi, T. Lenzi, J. Luetic, T. Maerschalk, A. Marinov, T. Seva, E. Starling, C. Vander Velde, P. Vanlaer, D. Vannerom, R. Yonamine, F. Zenoni

#### Ghent University, Ghent, Belgium

T. Cornelis, D. Dobur, A. Fagot, M. Gul, I. Khvastunov<sup>2</sup>, D. Poyraz, C. Roskas, S. Salva, D. Trocino, M. Tytgat, W. Verbeke, M. Vit, N. Zaganidis

#### Université Catholique de Louvain, Louvain-la-Neuve, Belgium

H. Bakhshiansohi, O. Bondu, S. Brochet, G. Bruno, C. Caputo, A. Caudron, P. David, S. De Visscher, C. Delaere, M. Delcourt, B. Francois, A. Giammanco, M. Komm, G. Krintiras, V. Lemaitre, A. Magitteri, A. Mertens, M. Musich, K. Piotrzkowski, L. Quertenmont, A. Saggio, M. Vidal Marono, S. Wertz, J. Zobec

#### Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

W.L. Aldá Júnior, F.L. Alves, G.A. Alves, L. Brito, G. Correia Silva, C. Hensel, A. Moraes, M.E. Pol, P. Rebello Teles

#### Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

E. Belchior Batista Das Chagas, W. Carvalho, J. Chinellato<sup>3</sup>, E. Coelho, E.M. Da Costa, G.G. Da Silveira<sup>4</sup>, D. De Jesus Damiao, S. Fonseca De Souza, L.M. Huertas Guativa, H. Malbouisson, M. Melo De Almeida, C. Mora Herrera, L. Mundim, H. Nogima, L.J. Sanchez Rosas, A. Santoro, A. Sznajder, M. Thiel, E.J. Tonelli Manganote<sup>3</sup>, F. Torres Da Silva De Araujo, A. Vilela Pereira

# Universidade Estadual Paulista <sup>a</sup>, Universidade Federal do ABC  $^b$ , São Paulo, Brazil

S. Ahuja<sup>a</sup>, C.A. Bernardes<sup>a</sup>, T.R. Fernandez Perez Tomei<sup>a</sup>, E.M. Gregores<sup>b</sup>, P.G. Mercadante<sup>b</sup>, S.F. Novaes<sup>a</sup>, Sandra S. Padula<sup>a</sup>, D. Romero Abad<sup>b</sup>, J.C. Ruiz Vargas<sup>a</sup>

# Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov, R. Hadjiiska, P. Iaydjiev, M. Misheva, M. Rodozov, M. Shopova, G. Sultanov

University of Sofia, Sofia, Bulgaria

A. Dimitrov, L. Litov, B. Pavlov, P. Petkov

Beihang University, Beijing, China

W. Fang<sup>5</sup>, X. Gao<sup>5</sup>, L. Yuan

# Institute of High Energy Physics, Beijing, China

M. Ahmad, J.G. Bian, G.M. Chen, H.S. Chen, M. Chen, Y. Chen, C.H. Jiang, D. Leggat, H. Liao, Z. Liu, F. Romeo, S.M. Shaheen, A. Spiezia, J. Tao, C. Wang, Z. Wang, E. Yazgan, H. Zhang, J. Zhao

# State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

Y. Ban, G. Chen, J. Li, Q. Li, S. Liu, Y. Mao, S.J. Qian, D. Wang, Z. Xu, F. Zhang<sup>5</sup>

#### Tsinghua University, Beijing, China

Y. Wang

#### Universidad de Los Andes, Bogota, Colombia

C. Avila, A. Cabrera, C.A. Carrillo Montoya, L.F. Chaparro Sierra, C. Florez, C.F. González Hernández, J.D. Ruiz Alvarez, M.A. Segura Delgado

# University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

B. Courbon, N. Godinovic, D. Lelas, I. Puljak, P.M. Ribeiro Cipriano, T. Sculac

#### University of Split, Faculty of Science, Split, Croatia

Z. Antunovic, M. Kovac

#### Institute Rudjer Boskovic, Zagreb, Croatia

V. Brigljevic, D. Ferencek, K. Kadija, B. Mesic, A. Starodumov<sup>6</sup>, T. Susa

#### University of Cyprus, Nicosia, Cyprus

M.W. Ather, A. Attikis, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis, H. Rykaczewski

#### Charles University, Prague, Czech Republic

M. Finger<sup>7</sup>, M. Finger Jr.<sup>7</sup>

#### Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin

# Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

H. Abdalla<sup>8</sup>, Y. Assran<sup>9,10</sup>, E. El-khateeb<sup>11</sup>

# National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

S. Bhowmik, R.K. Dewanjee, M. Kadastik, L. Perrini, M. Raidal, C. Veelken

#### Department of Physics, University of Helsinki, Helsinki, Finland

P. Eerola, H. Kirschenmann, J. Pekkanen, M. Voutilainen

#### Helsinki Institute of Physics, Helsinki, Finland

J. Havukainen, J.K. Heikkilä, T. Järvinen, V. Karimäki, R. Kinnunen, T. Lampén, K. Lassila-Perini, S. Laurila, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, H. Siikonen, E. Tuominen, J. Tuominiemi

#### Lappeenranta University of Technology, Lappeenranta, Finland

T. Tuuva

#### IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon, F. Couderc, M. Dejardin, D. Denegri, J.L. Faure, F. Ferri, S. Ganjour, S. Ghosh, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, C. Leloup, E. Locci, M. Machet, J. Malcles, G. Negro, J. Rander, A. Rosowsky, M.O. Sahin, M. Titov ¨

# Laboratoire Leprince-Ringuet, Ecole polytechnique, CNRS/IN2P3, Université Paris-Saclay, Palaiseau, France

A. Abdulsalam12, C. Amendola, I. Antropov, S. Baffioni, F. Beaudette, P. Busson, L. Cadamuro, C. Charlot, R. Granier de Cassagnac, M. Jo, I. Kucher, S. Lisniak, A. Lobanov, J. Martin Blanco, M. Nguyen, C. Ochando, G. Ortona, P. Paganini, P. Pigard, R. Salerno, J.B. Sauvan, Y. Sirois, A.G. Stahl Leiton, T. Strebler, Y. Yilmaz, A. Zabi, A. Zghiche

# Universit´e de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

J.-L. Agram13, J. Andrea, D. Bloch, J.-M. Brom, M. Buttignol, E.C. Chabert, C. Collard, E. Conte<sup>13</sup>, X. Coubez, F. Drouhin<sup>13</sup>, J.-C. Fontaine<sup>13</sup>, D. Gelé, U. Goerlach, M. Jansová, P. Juillot, A.-C. Le Bihan, N. Tonon, P. Van Hove

# Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

S. Gadrat

# Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France

S. Beauceron, C. Bernet, G. Boudoul, N. Chanon, R. Chierici, D. Contardo, P. Depasse, H. El Mamouni, J. Fay, L. Finco, S. Gascon, M. Gouzevitch, G. Grenier, B. Ille, F. Lagarde, I.B. Laktineh, M. Lethuillier, L. Mirabito, A.L. Pequegnot, S. Perries, A. Popov<sup>14</sup>, V. Sordini, M. Vander Donckt, S. Viret, S. Zhang

#### Georgian Technical University, Tbilisi, Georgia

 $T.$  Toriashvili $^{15}$ 

#### Tbilisi State University, Tbilisi, Georgia

Z. Tsamalaidze<sup>7</sup>

#### RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

C. Autermann, L. Feld, M.K. Kiesel, K. Klein, M. Lipinski, M. Preuten, C. Schomakers, J. Schulz, M. Teroerde, B. Wittmer, V. Zhukov<sup>14</sup>

#### RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

A. Albert, D. Duchardt, M. Endres, M. Erdmann, S. Erdweg, T. Esch, R. Fischer, A. Güth, T. Hebbeker, C. Heidemann, K. Hoepfner, S. Knutzen, M. Merschmeyer, A. Meyer, P. Millet, S. Mukherjee, T. Pook, M. Radziej, H. Reithler, M. Rieger, F. Scheuch, D. Teyssier, S. Thüer

#### RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

G. Flügge, B. Kargoll, T. Kress, A. Künsken, T. Müller, A. Nehrkorn, A. Nowack, C. Pistone, O. Pooth, A. Stahl $^{16}$ 

#### Deutsches Elektronen-Synchrotron, Hamburg, Germany

M. Aldaya Martin, T. Arndt, C. Asawatangtrakuldee, K. Beernaert, O. Behnke, U. Behrens, A. Bermúdez Martínez, A.A. Bin Anuar, K. Borras<sup>17</sup>, V. Botta, A. Campbell, P. Connor, C. Contreras-Campana, F. Costanza, C. Diez Pardos, G. Eckerlin, D. Eckstein, T. Eichhorn, E. Eren, E. Gallo<sup>18</sup>, J. Garay Garcia, A. Geiser, J.M. Grados Luyando, A. Grohsjean, P. Gunnellini, M. Guthoff, A. Harb, J. Hauk, M. Hempel<sup>19</sup>, H. Jung, M. Kasemann, J. Keaveney, C. Kleinwort, I. Korol, D. Krücker, W. Lange, A. Lelek, T. Lenz, K. Lipka, W. Lohmann<sup>19</sup>, R. Mankel, I.-A. Melzer-Pellmann, A.B. Meyer, M. Missiroli, G. Mittag, J. Mnich, A. Mussgiller, E. Ntomari, D. Pitzl, A. Raspereza, M. Savitskyi, P. Saxena, R. Shevchenko, N. Stefaniuk, G.P. Van Onsem, R. Walsh, Y. Wen, K. Wichmann, C. Wissing, O. Zenaiev

#### University of Hamburg, Hamburg, Germany

R. Aggleton, S. Bein, V. Blobel, M. Centis Vignali, T. Dreyer, E. Garutti, D. Gonzalez, J. Haller, A. Hinzmann, M. Hoffmann, A. Karavdina, R. Klanner, R. Kogler, N. Kovalchuk, S. Kurz, D. Marconi, M. Meyer, M. Niedziela, D. Nowatschin, F. Pantaleo<sup>16</sup>, T. Peiffer, A. Perieanu, C. Scharf, P. Schleper, A. Schmidt, S. Schumann, J. Schwandt, J. Sonneveld, H. Stadie, G. Steinbrück, F.M. Stober, M. Stöver, H. Tholen, D. Troendle, E. Usai, A. Vanhoefer, B. Vormwald

#### Institut für Experimentelle Teilchenphysik, Karlsruhe, Germany

M. Akbiyik, C. Barth, M. Baselga, S. Baur, E. Butz, R. Caspart, T. Chwalek, F. Colombo, W. De Boer, A. Dierlamm, N. Faltermann, B. Freund, R. Friese, M. Giffels, M.A. Harrendorf, F. Hartmann<sup>16</sup>, S.M. Heindl, U. Husemann, F. Kassel<sup>16</sup>, S. Kudella, H. Mildner, M.U. Mozer, Th. Müller, M. Plagge, G. Quast, K. Rabbertz, M. Schröder, I. Shvetsov, G. Sieber, H.J. Simonis, R. Ulrich, S. Wayand, M. Weber, T. Weiler, S. Williamson, C. Wöhrmann, R. Wolf

# Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis, T. Geralis, A. Kyriakis, D. Loukas, I. Topsis-Giotis

#### National and Kapodistrian University of Athens, Athens, Greece

G. Karathanasis, S. Kesisoglou, A. Panagiotou, N. Saoulidou, E. Tziaferi

# National Technical University of Athens, Athens, Greece

K. Kousouris

#### University of Ioánnina, Ioánnina, Greece

I. Evangelou, C. Foudas, P. Gianneios, P. Katsoulis, P. Kokkas, S. Mallios, N. Manthos, I. Papadopoulos, E. Paradas, J. Strologas, F.A. Triantis, D. Tsitsonis

# MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanad, N. Filipovic, G. Pasztor, O. Surányi, G.I. Veres<sup>20</sup>

Wigner Research Centre for Physics, Budapest, Hungary G. Bencze, C. Hajdu, D. Horvath<sup>21</sup>, Á. Hunyadi, F. Sikler, V. Veszpremi, G. Vesztergombi<sup>20</sup>

Institute of Nuclear Research ATOMKI, Debrecen, Hungary N. Beni, S. Czellar, J. Karancsi<sup>22</sup>, A. Makovec, J. Molnar, Z. Szillasi

# Institute of Physics, University of Debrecen, Debrecen, Hungary

M. Bartók<sup>20</sup>, P. Raics, Z.L. Trocsanyi, B. Ujvari

# Indian Institute of Science (IISc), Bangalore, India

S. Choudhury, J.R. Komaragiri

#### National Institute of Science Education and Research, Bhubaneswar, India

S. Bahinipati<sup>23</sup>, P. Mal, K. Mandal, A. Nayak<sup>24</sup>, D.K. Sahoo<sup>23</sup>, N. Sahoo, S.K. Swain

#### Panjab University, Chandigarh, India

S. Bansal, S.B. Beri, V. Bhatnagar, R. Chawla, N. Dhingra, A. Kaur, M. Kaur, S. Kaur, R. Kumar, P. Kumari, A. Mehta, J.B. Singh, G. Walia

#### University of Delhi, Delhi, India

Ashok Kumar, Aashaq Shah, A. Bhardwaj, S. Chauhan, B.C. Choudhary, R.B. Garg, S. Keshri, A. Kumar, S. Malhotra, M. Naimuddin, K. Ranjan, R. Sharma

#### Saha Institute of Nuclear Physics, HBNI, Kolkata, India

R. Bhardwaj<sup>25</sup>, R. Bhattacharya, S. Bhattacharya, U. Bhawandeep<sup>25</sup>, D. Bhowmik, S. Dey,

S. Dutt25, S. Dutta, S. Ghosh, N. Majumdar, A. Modak, K. Mondal, S. Mukhopadhyay,

S. Nandan, A. Purohit, P.K. Rout, A. Roy, S. Roy Chowdhury, S. Sarkar, M. Sharan, B. Singh, S. Thakur<sup>25</sup>

# Indian Institute of Technology Madras, Madras, India

P.K. Behera

#### Bhabha Atomic Research Centre, Mumbai, India

R. Chudasama, D. Dutta, V. Jha, V. Kumar, A.K. Mohanty<sup>16</sup>, P.K. Netrakanti, L.M. Pant, P. Shukla, A. Topkar

#### Tata Institute of Fundamental Research-A, Mumbai, India

T. Aziz, S. Dugad, B. Mahakud, S. Mitra, G.B. Mohanty, N. Sur, B. Sutar

#### Tata Institute of Fundamental Research-B, Mumbai, India

S. Banerjee, S. Bhattacharya, S. Chatterjee, P. Das, M. Guchait, Sa. Jain, S. Kumar, M. Maity<sup>26</sup>, G. Majumder, K. Mazumdar, T. Sarkar<sup>26</sup>, N. Wickramage<sup>27</sup>

Indian Institute of Science Education and Research (IISER), Pune, India

S. Chauhan, S. Dube, V. Hegde, A. Kapoor, K. Kothekar, S. Pandey, A. Rane, S. Sharma

#### Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Chenarani28, E. Eskandari Tadavani, S.M. Etesami28, M. Khakzad, M. Mohammadi Najafabadi, M. Naseri, S. Paktinat Mehdiabadi<sup>29</sup>, F. Rezaei Hosseinabadi, B. Safarzadeh<sup>30</sup>, M. Zeinali

#### University College Dublin, Dublin, Ireland

M. Felcini, M. Grunewald

INFN Sezione di Bari <sup>a</sup>, Università di Bari  $^b$ , Politecnico di Bari  $^c$ , Bari, Italy M. Abbrescia<sup>a,b</sup>, C. Calabria<sup>a,b</sup>, A. Colaleo<sup>a</sup>, D. Creanza<sup>a,c</sup>, L. Cristella<sup>a,b</sup>, N. De Filippis<sup>a,c</sup>, M. De Palma<sup>a,b</sup>, F. Errico<sup>a,b</sup>, L. Fiore<sup>a</sup>, G. Iaselli<sup>a,c</sup>, S. Lezki<sup>a,b</sup>, G. Maggi<sup>a,c</sup>, M. Maggi<sup>a</sup>, G. Miniello<sup>a,b</sup>, S. My<sup>a,b</sup>, S. Nuzzo<sup>a,b</sup>, A. Pompili<sup>a,b</sup>, G. Pugliese<sup>a,c</sup>, R. Radogna<sup>a</sup>, A. Ranieri<sup>a</sup>, G. Selvaggi<sup>a,b</sup>, A. Sharma<sup>a</sup>, L. Silvestris<sup>a,16</sup>, R. Venditti<sup>a</sup>, P. Verwilligen<sup> $a$ </sup>

# INFN Sezione di Bologna <sup>a</sup>, Università di Bologna  $^b$ , Bologna, Italy

G. Abbiendi<sup>a</sup>, C. Battilana<sup>a,b</sup>, D. Bonacorsi<sup>a,b</sup>, L. Borgonovi<sup>a,b</sup>, S. Braibant-Giacomelli<sup>a,b</sup>, R. Campanini<sup>a,b</sup>, P. Capiluppi<sup>a,b</sup>, A. Castro<sup>a,b</sup>, F.R. Cavallo<sup>a</sup>, S.S. Chhibra<sup>a,b</sup>, G. Codispoti<sup>a,b</sup>, M. Cuffiani<sup>a,b</sup>, G.M. Dallavalle<sup>a</sup>, F. Fabbri<sup>a</sup>, A. Fanfani<sup>a,b</sup>, D. Fasanella<sup>a,b</sup>, P. Giacomelli<sup>a</sup>, C. Grandi<sup>a</sup>, L. Guiducci<sup>a,b</sup>, F. Iemmi, S. Marcellini<sup>a</sup>, G. Masetti<sup>a</sup>, A. Montanari<sup>a</sup>, F.L. Navarria<sup>a,b</sup>, A. Perrotta<sup>a</sup>, A.M. Rossi<sup>a,b</sup>, T. Rovelli<sup>a,b</sup>, G.P. Siroli<sup>a,b</sup>, N. Tosi $^a$ 

# INFN Sezione di Catania <sup>a</sup>, Università di Catania  $^b$ , Catania, Italy

S. Albergo<sup>a,b</sup>, S. Costa<sup>a,b</sup>, A. Di Mattia<sup>a</sup>, F. Giordano<sup>a,b</sup>, R. Potenza<sup>a,b</sup>, A. Tricomi<sup>a,b</sup>, C. Tuve<sup> $a,b$ </sup>

# INFN Sezione di Firenze  $^a,$  Università di Firenze  $^b, \,$  Firenze, Italy

G. Barbagli<sup>a</sup>, K. Chatterjee<sup>a,b</sup>, V. Ciulli<sup>a,b</sup>, C. Civinini<sup>a</sup>, R. D'Alessandro<sup>a,b</sup>, E. Focardi<sup>a,b</sup>, P. Lenzi<sup>a,b</sup>, M. Meschini<sup>a</sup>, S. Paoletti<sup>a</sup>, L. Russo<sup>a,31</sup>, G. Sguazzoni<sup>a</sup>, D. Strom<sup>a</sup>, L. Viliani<sup>a</sup>

# INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi, S. Bianco, F. Fabbri, D. Piccolo, F. Primavera<sup>16</sup>

INFN Sezione di Genova  $^a,$  Università di Genova  $^b, \,$  Genova, Italy V. Calvelli<sup>a,b</sup>, F. Ferro<sup>a</sup>, L. Panizzi<sup>a,b</sup>, F. Ravera<sup>a,b</sup>, E. Robutti<sup>a</sup>, S. Tosi<sup>a,b</sup>

# INFN Sezione di Milano-Bicocca <sup>a</sup>, Università di Milano-Bicocca <sup>b</sup>, Milano, Italy

A. Benaglia<sup>a</sup>, A. Beschi<sup>b</sup>, L. Brianza<sup>a,b</sup>, F. Brivio<sup>a,b</sup>, V. Ciriolo<sup>a,b,16</sup>, M.E. Dinardo<sup>a,b</sup>, S. Fiorendi<sup>a,b</sup>, S. Gennai<sup>a</sup>, A. Ghezzi<sup>a,b</sup>, P. Govoni<sup>a,b</sup>, M. Malberti<sup>a,b</sup>, S. Malvezzi<sup>a</sup>, R.A. Manzoni<sup>a,b</sup>, D. Menasce<sup>a</sup>, L. Moroni<sup>a</sup>, M. Paganoni<sup>a,b</sup>, K. Pauwels<sup>a,b</sup>, D. Pedrini<sup>a</sup>, S. Pigazzini<sup>a,b,32</sup>, S. Ragazzi<sup>a,b</sup>, T. Tabarelli de Fatis<sup>a,b</sup>

# INFN Sezione di Napoli <sup>a</sup>, Università di Napoli 'Federico II'  $^b$ , Napoli, Italy, Università della Basilicata  $^c$ , Potenza, Italy, Università G. Marconi  $^d$ , Roma, Italy

S. Buontempo<sup>a</sup>, N. Cavallo<sup>a,c</sup>, S. Di Guida<sup>a,d,16</sup>, F. Fabozzi<sup>a,c</sup>, F. Fienga<sup>a,b</sup>, A.O.M. Iorio<sup>a,b</sup>, W.A. Khan<sup>a</sup>, L. Lista<sup>a</sup>, S. Meola<sup>a,d,16</sup>, P. Paolucci<sup>a,16</sup>, C. Sciacca<sup>a,b</sup>, F. Thyssen<sup>a</sup>

# INFN Sezione di Padova <sup>a</sup>, Università di Padova  $^b$ , Padova, Italy, Università di  $Trento<sup>c</sup>$ , Trento, Italy

P. Azzi<sup>a</sup>, N. Bacchetta<sup>a</sup>, L. Benato<sup>a,b</sup>, A. Boletti<sup>a,b</sup>, R. Carlin<sup>a,b</sup>, A. Carvalho Antunes De Oliveira<sup>a,b</sup>, P. Checchia<sup>a</sup>, M. Dall'Osso<sup>a,b</sup>, P. De Castro Manzano<sup>a</sup>, T. Dorigo<sup>a</sup>, U. Dosselli<sup>a</sup>, F. Gasparini<sup>a,b</sup>, U. Gasparini<sup>a,b</sup>, A. Gozzelino<sup>a</sup>, S. Lacaprara<sup>a</sup>, P. Lujan, M. Margoni<sup>a,b</sup>, A.T. Meneguzzo<sup>a,b</sup>, N. Pozzobon<sup>a,b</sup>, P. Ronchese<sup>a,b</sup>, R. Rossin<sup>a,b</sup>, F. Simonetto<sup>a,b</sup>, A. Tiko, E. Torassa<sup>a</sup>, M. Zanetti<sup>a,b</sup>, P. Zotto<sup>a,b</sup>, G. Zumerle<sup>a,b</sup>

# INFN Sezione di Pavia <sup>a</sup>, Università di Pavia  $^b$ , Pavia, Italy

A. Braghieri<sup>a</sup>, A. Magnani<sup>a</sup>, P. Montagna<sup>a,b</sup>, S.P. Ratti<sup>a,b</sup>, V. Re<sup>a</sup>, M. Ressegotti<sup>a,b</sup>, C. Riccardi<sup>a,b</sup>, P. Salvini<sup>a</sup>, I. Vai<sup>a,b</sup>, P. Vitulo<sup>a,b</sup>

# INFN Sezione di Perugia  $^a,$  Università di Perugia  $^b, \,$  Perugia, Italy

L. Alunni Solestizi<sup>a,b</sup>, M. Biasini<sup>a,b</sup>, G.M. Bilei<sup>a</sup>, C. Cecchi<sup>a,b</sup>, D. Ciangottini<sup>a,b</sup>, L. Fanò<sup>a,b</sup>, P. Lariccia<sup>a,b</sup>, R. Leonardi<sup>a,b</sup>, E. Manoni<sup>a</sup>, G. Mantovani<sup>a,b</sup>, V. Mariani<sup>a,b</sup>, M. Menichelli<sup>a</sup>, A. Rossi<sup>a,b</sup>, A. Santocchia<sup>a,b</sup>, D. Spiga<sup>a</sup>

# INFN Sezione di Pisa <sup>a</sup>, Università di Pisa <sup>b</sup>, Scuola Normale Superiore di Pisa<sup>c</sup>, Pisa, Italy

K. Androsov<sup>a</sup>, P. Azzurri<sup>a, 16</sup>, G. Bagliesi<sup>a</sup>, L. Bianchini<sup>a</sup>, T. Boccali<sup>a</sup>, L. Borrello, R. Castaldi<sup>a</sup>, M.A. Ciocci<sup>a,b</sup>, R. Dell'Orso<sup>a</sup>, G. Fedi<sup>a</sup>, L. Giannini<sup>a,c</sup>, A. Giassi<sup>a</sup>, M.T. Grippo<sup>a,31</sup>, F. Ligabue<sup>a,c</sup>, T. Lomtadze<sup>a</sup>, E. Manca<sup>a,c</sup>, G. Mandorli<sup>a,c</sup>, A. Messineo<sup>a,b</sup>, F. Palla<sup>a</sup>, A. Rizzi<sup>a,b</sup>, P. Spagnolo<sup>a</sup>, R. Tenchini<sup>a</sup>, G. Tonelli<sup>a,b</sup>, A. Venturi<sup>a</sup>, P.G. Verdini $^a$ 

# INFN Sezione di Roma <sup>a</sup>, Sapienza Università di Roma  $^b$ , Rome, Italy

L. Barone<sup>a,b</sup>, F. Cavallari<sup>a</sup>, M. Cipriani<sup>a,b</sup>, N. Daci<sup>a</sup>, D. Del Re<sup>a,b</sup>, E. Di Marco<sup>a,b</sup>, M. Diemoz<sup>a</sup>, S. Gelli<sup>a,b</sup>, E. Longo<sup>a,b</sup>, F. Margaroli<sup>a,b</sup>, B. Marzocchi<sup>a,b</sup>, P. Meridiani<sup>a</sup>, G. Organtini<sup>a,b</sup>, R. Paramatti<sup>a,b</sup>, F. Preiato<sup>a,b</sup>, S. Rahatlou<sup>a,b</sup>, C. Rovelli<sup>a</sup>, F. Santanastasio<sup> $a,b$ </sup>

# INFN Sezione di Torino <sup>a</sup>, Università di Torino  $^b,$  Torino, Italy, Università del Piemonte Orientale  $^c$ , Novara, Italy

N. Amapane $^{a,b}$ , R. Arcidiacono $^{a,c}$ , S. Argiro $^{a,b}$ , M. Arneodo $^{a,c}$ , N. Bartosik $^a$ , R. Bellan $^{a,b}$ , C. Biino<sup>a</sup>, N. Cartiglia<sup>a</sup>, F. Cenna<sup>a,b</sup>, M. Costa<sup>a,b</sup>, R. Covarelli<sup>a,b</sup>, A. Degano<sup>a,b</sup>, N. Demaria<sup>a</sup>, B. Kiani<sup>a,b</sup>, C. Mariotti<sup>a</sup>, S. Maselli<sup>a</sup>, E. Migliore<sup>a,b</sup>, V. Monaco<sup>a,b</sup>, E. Monteil<sup>a,b</sup>, M. Monteno<sup>a</sup>, M.M. Obertino<sup>a,b</sup>, L. Pacher<sup>a,b</sup>, N. Pastrone<sup>a</sup>, M. Pelliccioni<sup>a</sup>, G.L. Pinna Angioni<sup>a, b</sup>, A. Romero<sup>a, b</sup>, M. Ruspa<sup>a, c</sup>, R. Sacchi<sup>a, b</sup>, K. Shchelina<sup>a, b</sup>, V. Sola<sup>a</sup>, A. Solano<sup>a,b</sup>, A. Staiano<sup>a</sup>, P. Traczyk<sup>a,b</sup>

# INFN Sezione di Trieste <sup>a</sup>, Università di Trieste  $^b$ , Trieste, Italy

S. Belforte<sup>a</sup>, M. Casarsa<sup>a</sup>, F. Cossutti<sup>a</sup>, G. Della Ricca<sup>a,b</sup>, A. Zanetti<sup>a</sup>

#### Kyungpook National University

D.H. Kim, G.N. Kim, M.S. Kim, J. Lee, S. Lee, S.W. Lee, C.S. Moon, Y.D. Oh, S. Sekmen, D.C. Son, Y.C. Yang

# Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

H. Kim, D.H. Moon, G. Oh

#### Hanyang University, Seoul, Korea

J.A. Brochero Cifuentes, J. Goh, T.J. Kim

#### Korea University, Seoul, Korea

S. Cho, S. Choi, Y. Go, D. Gyun, S. Ha, B. Hong, Y. Jo, Y. Kim, K. Lee, K.S. Lee, S. Lee, J. Lim, S.K. Park, Y. Roh

#### Seoul National University, Seoul, Korea

J. Almond, J. Kim, J.S. Kim, H. Lee, K. Lee, K. Nam, S.B. Oh, B.C. Radburn-Smith, S.h. Seo, U.K. Yang, H.D. Yoo, G.B. Yu

University of Seoul, Seoul, Korea

H. Kim, J.H. Kim, J.S.H. Lee, I.C. Park

#### Sungkyunkwan University, Suwon, Korea

Y. Choi, C. Hwang, J. Lee, I. Yu

#### Vilnius University, Vilnius, Lithuania

V. Dudenas, A. Juodagalvis, J. Vaitkus

# National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Ahmed, Z.A. Ibrahim, M.A.B. Md Ali<sup>33</sup>, F. Mohamad Idris<sup>34</sup>, W.A.T. Wan Abdullah, M.N. Yusli, Z. Zolkapli

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico Reyes-Almanza, R, Ramirez-Sanchez, G., Duran-Osuna, M. C., H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-De La Cruz<sup>35</sup>, Rabadan-Trejo, R. I., R. Lopez-Fernandez, J. Mejia Guisao, A. Sanchez-Hernandez

#### Universidad Iberoamericana, Mexico City, Mexico

S. Carrillo Moreno, C. Oropeza Barrera, F. Vazquez Valencia

#### Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

J. Eysermans, I. Pedraza, H.A. Salazar Ibarguen, C. Uribe Estrada

Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico A. Morelos Pineda

University of Auckland, Auckland, New Zealand D. Krofcheck

University of Canterbury, Christchurch, New Zealand P.H. Butler

#### National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad, M. Ahmad, Q. Hassan, H.R. Hoorani, A. Saddique, M.A. Shah, M. Shoaib, M. Waqas

#### National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska, M. Bluj, B. Boimska, T. Frueboes, M. G´orski, M. Kazana, K. Nawrocki, M. Szleper, P. Zalewski

# Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski, A. Byszuk36, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski, M. Misiura, M. Olszewski, A. Pyskir, M. Walczak

# Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

P. Bargassa, C. Beirão Da Cruz E Silva, A. Di Francesco, P. Faccioli, B. Galinhas, M. Gallinaro, J. Hollar, N. Leonardo, L. Lloret Iglesias, M.V. Nemallapudi, J. Seixas, G. Strong, O. Toldaiev, D. Vadruccio, J. Varela

# Joint Institute for Nuclear Research, Dubna, Russia

S. Afanasiev, V. Alexakhin, A. Golunov, I. Golutvin, N. Gorbounov, I. Gorbunov, A. Kamenev, V. Karjavin, A. Lanev, A. Malakhov, V. Matveev<sup>37,38</sup>, P. Moisenz, V. Palichik, V. Perelygin, S. Shmatov, S. Shulha, N. Skatchkov, V. Smirnov, A. Zarubin

#### Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia

Y. Ivanov, V. Kim<sup>39</sup>, E. Kuznetsova<sup>40</sup>, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, D. Sosnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev

#### Institute for Nuclear Research, Moscow, Russia

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, A. Karneyeu, M. Kirsanov, N. Krasnikov, A. Pashenkov, D. Tlisov, A. Toropin

#### Institute for Theoretical and Experimental Physics, Moscow, Russia

V. Epshteyn, V. Gavrilov, N. Lychkovskaya, V. Popov, I. Pozdnyakov, G. Safronov, A. Spiridonov, A. Stepennov, V. Stolin, M. Toms, E. Vlasov, A. Zhokin

Moscow Institute of Physics and Technology, Moscow, Russia

T. Aushev, A. Bylinkin<sup>38</sup>

# National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia

R. Chistov<sup>41</sup>, M. Danilov<sup>41</sup>, P. Parygin, D. Philippov, S. Polikarpov, E. Tarkovskii

P.N. Lebedev Physical Institute, Moscow, Russia V. Andreev, M. Azarkin38, I. Dremin38, M. Kirakosyan38, S.V. Rusakov, A. Terkulov

# Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia

A. Baskakov, A. Belyaev, E. Boos, V. Bunichev, M. Dubinin<sup>42</sup>, L. Dudko, A. Ershov, V. Klyukhin, O. Kodolova, I. Lokhtin, I. Miagkov, S. Obraztsov, M. Perfilov, S. Petrushanko, V. Savrin

Novosibirsk State University (NSU), Novosibirsk, Russia V. Blinov<sup>43</sup>, D. Shtol<sup>43</sup>, Y. Skovpen<sup>43</sup>

# State Research Center of Russian Federation, Institute for High Energy Physics of NRC " Kurchatov Institute ", Protvino, Russia

I. Azhgirey, I. Bayshev, S. Bitioukov, D. Elumakhov, A. Godizov, V. Kachanov, A. Kalinin, D. Konstantinov, P. Mandrik, V. Petrov, R. Ryutin, A. Sobol, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

National Research Tomsk Polytechnic University, Tomsk, Russia A. Babaev

# University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia

P. Adzic44, P. Cirkovic, D. Devetak, M. Dordevic, J. Milosevic

# Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J. Alcaraz Maestre, I. Bachiller, M. Barrio Luna, M. Cerrada, N. Colino, B. De La Cruz, A. Delgado Peris, C. Fernandez Bedoya, J.P. Fern´andez Ramos, J. Flix, M.C. Fouz, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, D. Moran, A. Pérez-Calero Yzquierdo, J. Puerta Pelayo, I. Redondo, L. Romero, M.S. Soares, A. Triossi, A. Alvarez ´ Fernández

#### Universidad Autónoma de Madrid, Madrid, Spain

C. Albajar, J.F. de Trocóniz

#### Universidad de Oviedo, Oviedo, Spain

J. Cuevas, C. Erice, J. Fernandez Menendez, I. Gonzalez Caballero, J.R. González Fern´andez, E. Palencia Cortezon, S. Sanchez Cruz, P. Vischia, J.M. Vizan Garcia

# Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

I.J. Cabrillo, A. Calderon, B. Chazin Quero, J. Duarte Campderros, M. Fernandez, P.J. Fernández Manteca, J. Garcia-Ferrero, A. García Alonso, G. Gomez, A. Lopez Virto, J. Marco, C. Martinez Rivero, P. Martinez Ruiz del Arbol, F. Matorras, J. Piedra Gomez, C. Prieels, T. Rodrigo, A. Ruiz-Jimeno, L. Scodellaro, N. Trevisani, I. Vila, R. Vilar Cortabitarte

#### CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo, B. Akgun, E. Auffray, P. Baillon, A.H. Ball, D. Barney, J. Bendavid, M. Bianco, A. Bocci, C. Botta, T. Camporesi, R. Castello, M. Cepeda, G. Cerminara, E. Chapon, Y. Chen, D. d'Enterria, A. Dabrowski, V. Daponte, A. David, M. De Gruttola, A. De Roeck, N. Deelen, M. Dobson, T. du Pree, M. Dünser, N. Dupont, A. Elliott-Peisert, P. Everaerts, F. Fallavollita, G. Franzoni, J. Fulcher, W. Funk, D. Gigi, A. Gilbert, K. Gill, F. Glege, D. Gulhan, J. Hegeman, V. Innocente, A. Jafari, P. Janot, O. Karacheban<sup>19</sup>, J. Kieseler, V. Knünz, A. Kornmayer, M.J. Kortelainen, M. Krammer<sup>1</sup>, C. Lange, P. Lecoq, C. Lourenço, M.T. Lucchini, L. Malgeri, M. Mannelli, A. Martelli, F. Meijers, J.A. Merlin, S. Mersi, E. Meschi, P. Milenovic<sup>45</sup>, F. Moortgat, M. Mulders, H. Neugebauer, J. Ngadiuba, S. Orfanelli, L. Orsini, L. Pape, E. Perez, M. Peruzzi, A. Petrilli, G. Petrucciani, A. Pfeiffer, M. Pierini, F.M. Pitters, D. Rabady, A. Racz, T. Reis, G. Rolandi<sup>46</sup>, M. Rovere, H. Sakulin, C. Schäfer, C. Schwick, M. Seidel, M. Selvaggi, A. Sharma, P. Silva, P. Sphicas<sup>47</sup>, A. Stakia, J. Steggemann, M. Stoye, M. Tosi, D. Treille, A. Tsirou, V. Veckalns<sup>48</sup>, M. Verweij, W.D. Zeuner

#### Paul Scherrer Institut, Villigen, Switzerland

W. Bertl<sup>†</sup>, L. Caminada<sup>49</sup>, K. Deiters, W. Erdmann, R. Horisberger, Q. Ingram, H.C. Kaestli, D. Kotlinski, U. Langenegger, T. Rohe, S.A. Wiederkehr

# ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

M. Backhaus, L. Bäni, P. Berger, B. Casal, G. Dissertori, M. Dittmar, M. Donegà, C. Dorfer, C. Grab, C. Heidegger, D. Hits, J. Hoss, G. Kasieczka, T. Klijnsma, W. Lustermann, B. Mangano, M. Marionneau, M.T. Meinhard, D. Meister, F. Micheli, P. Musella, F. Nessi-Tedaldi, F. Pandolfi, J. Pata, F. Pauss, G. Perrin, L. Perrozzi, M. Quittnat, M. Reichmann, D.A. Sanz Becerra, M. Schönenberger, L. Shchutska, V.R. Tavolaro, K. Theofilatos, M.L. Vesterbacka Olsson, R. Wallny, D.H. Zhu

#### Universität Zürich, Zurich, Switzerland

T.K. Aarrestad, C. Amsler<sup>50</sup>, M.F. Canelli, A. De Cosa, R. Del Burgo, S. Donato, C. Galloni, T. Hreus, B. Kilminster, D. Pinna, G. Rauco, P. Robmann, D. Salerno, K. Schweiger, C. Seitz, Y. Takahashi, A. Zucchetta

#### National Central University, Chung-Li, Taiwan

V. Candelise, Y.H. Chang, K.y. Cheng, T.H. Doan, Sh. Jain, R. Khurana, C.M. Kuo, W. Lin, A. Pozdnyakov, S.S. Yu

#### National Taiwan University (NTU), Taipei, Taiwan

Arun Kumar, P. Chang, Y. Chao, K.F. Chen, P.H. Chen, F. Fiori, W.-S. Hou, Y. Hsiung, Y.F. Liu, R.-S. Lu, E. Paganis, A. Psallidas, A. Steen, J.f. Tsai

# Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

B. Asavapibhop, K. Kovitanggoon, G. Singh, N. Srimanobhas

# Cukurova University, Physics Department, Science and Art Faculty, Adana, **Turkey**

A. Bat, F. Boran, S. Cerci<sup>51</sup>, S. Damarseckin, Z.S. Demiroglu, C. Dozen, I. Dumanoglu, S. Girgis, G. Gokbulut, Y. Guler, I. Hos<sup>52</sup>, E.E. Kangal<sup>53</sup>, O. Kara, A. Kayis Topaksu, U. Kiminsu, M. Oglakci, G. Onengut, K. Ozdemir<sup>54</sup>, D. Sunar Cerci<sup>51</sup>, U.G. Tok, H. Topakli55, S. Turkcapar, I.S. Zorbakir, C. Zorbilmez

# Middle East Technical University, Physics Department, Ankara, Turkey

G. Karapinar56, K. Ocalan57, M. Yalvac, M. Zeyrek

#### Bogazici University, Istanbul, Turkey

E. Gülmez, M. Kaya<sup>58</sup>, O. Kaya<sup>59</sup>, S. Tekten, E.A. Yetkin<sup>60</sup>

## Istanbul Technical University, Istanbul, Turkey

M.N. Agaras, S. Atay, A. Cakir, K. Cankocak, Y. Komurcu

# Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkov, Ukraine

B. Grynyov

# National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine

L. Levchuk

#### University of Bristol, Bristol, United Kingdom

F. Ball, L. Beck, J.J. Brooke, D. Burns, E. Clement, D. Cussans, O. Davignon, H. Flacher,

J. Goldstein, G.P. Heath, H.F. Heath, L. Kreczko, D.M. Newbold<sup>61</sup>, S. Paramesvaran,

T. Sakuma, S. Seif El Nasr-storey, D. Smith, V.J. Smith

#### Rutherford Appleton Laboratory, Didcot, United Kingdom

K.W. Bell, A. Belyaev<sup>62</sup>, C. Brew, R.M. Brown, L. Calligaris, D. Cieri, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, J. Linacre, E. Olaiya, D. Petyt, C.H. Shepherd-Themistocleous, A. Thea, I.R. Tomalin, T. Williams, W.J. Womersley

#### Imperial College, London, United Kingdom

G. Auzinger, R. Bainbridge, P. Bloch, J. Borg, S. Breeze, O. Buchmuller, A. Bundock, S. Casasso, M. Citron, D. Colling, L. Corpe, P. Dauncey, G. Davies, M. Della Negra, R. Di Maria, Y. Haddad, G. Hall, G. Iles, T. James, R. Lane, C. Laner, L. Lyons, A.-M. Magnan, S. Malik, L. Mastrolorenzo, T. Matsushita, J. Nash<sup>63</sup>, A. Nikitenko<sup>6</sup>, V. Palladino, M. Pesaresi, D.M. Raymond, A. Richards, A. Rose, E. Scott, C. Seez, A. Shtipliyski, S. Summers, A. Tapper, K. Uchida, M. Vazquez Acosta<sup>64</sup>, T. Virdee<sup>16</sup>, N. Wardle, D. Winterbottom, J. Wright, S.C. Zenz

#### Brunel University, Uxbridge, United Kingdom

J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, A. Morton, I.D. Reid, L. Teodorescu, S. Zahid

#### Baylor University, Waco, USA

A. Borzou, K. Call, J. Dittmann, K. Hatakeyama, H. Liu, N. Pastika, C. Smith

#### Catholic University of America, Washington DC, USA

R. Bartek, A. Dominguez

#### The University of Alabama, Tuscaloosa, USA

A. Buccilli, S.I. Cooper, C. Henderson, P. Rumerio, C. West

#### Boston University, Boston, USA

D. Arcaro, A. Avetisyan, T. Bose, D. Gastler, D. Rankin, C. Richardson, J. Rohlf, L. Sulak, D. Zou

#### Brown University, Providence, USA

G. Benelli, D. Cutts, M. Hadley, J. Hakala, U. Heintz, J.M. Hogan, K.H.M. Kwok, E. Laird, G. Landsberg, J. Lee, Z. Mao, M. Narain, J. Pazzini, S. Piperov, S. Sagir, R. Syarif, D. Yu

#### University of California, Davis, Davis, USA

R. Band, C. Brainerd, R. Breedon, D. Burns, M. Calderon De La Barca Sanchez, M. Chertok, J. Conway, R. Conway, P.T. Cox, R. Erbacher, C. Flores, G. Funk, W. Ko, R. Lander, C. Mclean, M. Mulhearn, D. Pellett, J. Pilot, S. Shalhout, M. Shi, J. Smith, D. Stolp, D. Taylor, K. Tos, M. Tripathi, Z. Wang

#### University of California, Los Angeles, USA

M. Bachtis, C. Bravo, R. Cousins, A. Dasgupta, A. Florent, J. Hauser, M. Ignatenko, N. Mccoll, S. Regnard, D. Saltzberg, C. Schnaible, V. Valuev

#### University of California, Riverside, Riverside, USA

E. Bouvier, K. Burt, R. Clare, J. Ellison, J.W. Gary, S.M.A. Ghiasi Shirazi, G. Hanson, G. Karapostoli, E. Kennedy, F. Lacroix, O.R. Long, M. Olmedo Negrete, M.I. Paneva, W. Si, L. Wang, H. Wei, S. Wimpenny, B. R. Yates

#### University of California, San Diego, La Jolla, USA

J.G. Branson, S. Cittolin, M. Derdzinski, R. Gerosa, D. Gilbert, B. Hashemi, A. Holzner, D. Klein, G. Kole, V. Krutelyov, J. Letts, M. Masciovecchio, D. Olivito, S. Padhi, M. Pieri, M. Sani, V. Sharma, S. Simon, M. Tadel, A. Vartak, S. Wasserbaech<sup>65</sup>, J. Wood, F. Würthwein, A. Yagil, G. Zevi Della Porta

# University of California, Santa Barbara - Department of Physics, Santa Barbara, USA

N. Amin, R. Bhandari, J. Bradmiller-Feld, C. Campagnari, A. Dishaw, V. Dutta, M. Franco Sevilla, L. Gouskos, R. Heller, J. Incandela, A. Ovcharova, H. Qu, J. Richman, D. Stuart, I. Suarez, J. Yoo

#### California Institute of Technology, Pasadena, USA

D. Anderson, A. Bornheim, J. Bunn, I. Dutta, J.M. Lawhorn, H.B. Newman, T. Q. Nguyen, C. Pena, M. Spiropulu, J.R. Vlimant, R. Wilkinson, S. Xie, Z. Zhang, R.Y. Zhu

#### Carnegie Mellon University, Pittsburgh, USA

M.B. Andrews, T. Ferguson, T. Mudholkar, M. Paulini, J. Russ, M. Sun, H. Vogel, I. Vorobiev, M. Weinberg

#### University of Colorado Boulder, Boulder, USA

J.P. Cumalat, W.T. Ford, F. Jensen, A. Johnson, M. Krohn, S. Leontsinis, E. Macdonald, T. Mulholland, K. Stenson, S.R. Wagner

#### Cornell University, Ithaca, USA

J. Alexander, J. Chaves, Y. Cheng, J. Chu, S. Dittmer, K. Mcdermott, N. Mirman, J.R. Patterson, D. Quach, A. Rinkevicius, A. Ryd, L. Skinnari, L. Soffi, S.M. Tan, Z. Tao, J. Thom, J. Tucker, P. Wittich, M. Zientek

#### Fermi National Accelerator Laboratory, Batavia, USA

S. Abdullin, M. Albrow, M. Alyari, G. Apollinari, A. Apresyan, A. Apyan, S. Banerjee, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, G. Bolla† , K. Burkett, J.N. Butler, A. Canepa, G.B. Cerati, H.W.K. Cheung, F. Chlebana, M. Cremonesi, J. Duarte, V.D. Elvira, J. Freeman, Z. Gecse, E. Gottschalk, L. Gray, D. Green, S. Grünendahl, O. Gutsche, J. Hanlon, R.M. Harris, S. Hasegawa, J. Hirschauer, Z. Hu, B. Jayatilaka, S. Jindariani, M. Johnson, U. Joshi, B. Klima, B. Kreis, S. Lammel, D. Lincoln, R. Lipton, M. Liu, T. Liu, R. Lopes De S´a, J. Lykken, K. Maeshima, N. Magini, J.M. Marraffino, D. Mason, P. McBride, P. Merkel, S. Mrenna, S. Nahn, V. O'Dell, K. Pedro, O. Prokofyev, G. Rakness, L. Ristori, A. Savoy-Navarro<sup>66</sup>, B. Schneider, E. Sexton-Kennedy, A. Soha, W.J. Spalding, L. Spiegel, S. Stoynev, J. Strait, N. Strobbe, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, C. Vernieri, M. Verzocchi, R. Vidal, M. Wang, H.A. Weber, A. Whitbeck, W. Wu

#### University of Florida, Gainesville, USA

D. Acosta, P. Avery, P. Bortignon, D. Bourilkov, A. Brinkerhoff, A. Carnes, M. Carver, D. Curry, R.D. Field, I.K. Furic, S.V. Gleyzer, B.M. Joshi, J. Konigsberg, A. Korytov, K. Kotov, P. Ma, K. Matchev, H. Mei, G. Mitselmakher, K. Shi, D. Sperka, N. Terentyev, L. Thomas, J. Wang, S. Wang, J. Yelton

#### Florida International University, Miami, USA

Y.R. Joshi, S. Linn, P. Markowitz, J.L. Rodriguez

#### Florida State University, Tallahassee, USA

A. Ackert, T. Adams, A. Askew, S. Hagopian, V. Hagopian, K.F. Johnson, T. Kolberg, G. Martinez, T. Perry, H. Prosper, A. Saha, A. Santra, V. Sharma, R. Yohay

#### Florida Institute of Technology, Melbourne, USA

M.M. Baarmand, V. Bhopatkar, S. Colafranceschi, M. Hohlmann, D. Noonan, T. Roy, F. Yumiceva

#### University of Illinois at Chicago (UIC), Chicago, USA

M.R. Adams, L. Apanasevich, D. Berry, R.R. Betts, R. Cavanaugh, X. Chen, O. Evdokimov, C.E. Gerber, D.A. Hangal, D.J. Hofman, K. Jung, J. Kamin, I.D. Sandoval Gonzalez, M.B. Tonjes, H. Trauger, N. Varelas, H. Wang, Z. Wu, J. Zhang

#### The University of Iowa, Iowa City, USA

B. Bilki<sup>67</sup>, W. Clarida, K. Dilsiz<sup>68</sup>, S. Durgut, R.P. Gandrajula, M. Haytmyradov, V. Khristenko, J.-P. Merlo, H. Mermerkaya<sup>69</sup>, A. Mestvirishvili, A. Moeller, J. Nachtman, H. Ogul70, Y. Onel, F. Ozok71, A. Penzo, C. Snyder, E. Tiras, J. Wetzel, K. Yi

## Johns Hopkins University, Baltimore, USA

B. Blumenfeld, A. Cocoros, N. Eminizer, D. Fehling, L. Feng, A.V. Gritsan, P. Maksimovic, J. Roskes, U. Sarica, M. Swartz, M. Xiao, C. You

#### The University of Kansas, Lawrence, USA

A. Al-bataineh, P. Baringer, A. Bean, S. Boren, J. Bowen, J. Castle, S. Khalil, A. Kropivnitskaya, D. Majumder, W. Mcbrayer, M. Murray, C. Rogan, C. Royon, S. Sanders, E. Schmitz, J.D. Tapia Takaki, Q. Wang

#### Kansas State University, Manhattan, USA

A. Ivanov, K. Kaadze, Y. Maravin, A. Mohammadi, L.K. Saini, N. Skhirtladze

#### Lawrence Livermore National Laboratory, Livermore, USA

F. Rebassoo, D. Wright

#### University of Maryland, College Park, USA

A. Baden, O. Baron, A. Belloni, S.C. Eno, Y. Feng, C. Ferraioli, N.J. Hadley, S. Jabeen, G.Y. Jeng, R.G. Kellogg, J. Kunkle, A.C. Mignerey, F. Ricci-Tam, Y.H. Shin, A. Skuja, S.C. Tonwar

#### Massachusetts Institute of Technology, Cambridge, USA

D. Abercrombie, B. Allen, V. Azzolini, R. Barbieri, A. Baty, G. Bauer, R. Bi, S. Brandt, W. Busza, I.A. Cali, M. D'Alfonso, Z. Demiragli, G. Gomez Ceballos, M. Goncharov, P. Harris, D. Hsu, M. Hu, Y. Iiyama, G.M. Innocenti, M. Klute, D. Kovalskyi, Y.-J. Lee, A. Levin, P.D. Luckey, B. Maier, A.C. Marini, C. Mcginn, C. Mironov, S. Narayanan, X. Niu, C. Paus, C. Roland, G. Roland, J. Salfeld-Nebgen, G.S.F. Stephans, K. Sumorok, K. Tatar, D. Velicanu, J. Wang, T.W. Wang, B. Wyslouch

#### University of Minnesota, Minneapolis, USA

A.C. Benvenuti, R.M. Chatterjee, A. Evans, P. Hansen, J. Hiltbrand, S. Kalafut, Y. Kubota, Z. Lesko, J. Mans, S. Nourbakhsh, N. Ruckstuhl, R. Rusack, J. Turkewitz, M.A. Wadud

#### University of Mississippi, Oxford, USA

J.G. Acosta, S. Oliveros

#### University of Nebraska-Lincoln, Lincoln, USA

E. Avdeeva, K. Bloom, D.R. Claes, C. Fangmeier, F. Golf, R. Gonzalez Suarez, R. Kamalieddin, I. Kravchenko, J. Monroy, J.E. Siado, G.R. Snow, B. Stieger

#### State University of New York at Buffalo, Buffalo, USA

J. Dolen, A. Godshalk, C. Harrington, I. Iashvili, D. Nguyen, A. Parker, S. Rappoccio, B. Roozbahani

#### Northeastern University, Boston, USA

G. Alverson, E. Barberis, C. Freer, A. Hortiangtham, A. Massironi, D.M. Morse, T. Orimoto, R. Teixeira De Lima, T. Wamorkar, B. Wang, A. Wisecarver, D. Wood

#### Northwestern University, Evanston, USA

S. Bhattacharya, O. Charaf, K.A. Hahn, N. Mucia, N. Odell, M.H. Schmitt, K. Sung, M. Trovato, M. Velasco

#### University of Notre Dame, Notre Dame, USA

R. Bucci, N. Dev, M. Hildreth, K. Hurtado Anampa, C. Jessop, D.J. Karmgard, N. Kellams, K. Lannon, W. Li, N. Loukas, N. Marinelli, F. Meng, C. Mueller, Y. Musienko<sup>37</sup>, M. Planer, A. Reinsvold, R. Ruchti, P. Siddireddy, G. Smith, S. Taroni, M. Wayne, A. Wightman, M. Wolf, A. Woodard

#### The Ohio State University, Columbus, USA

J. Alimena, L. Antonelli, B. Bylsma, L.S. Durkin, S. Flowers, B. Francis, A. Hart, C. Hill, W. Ji, T.Y. Ling, B. Liu, W. Luo, B.L. Winer, H.W. Wulsin

#### Princeton University, Princeton, USA

S. Cooperstein, O. Driga, P. Elmer, J. Hardenbrook, P. Hebda, S. Higginbotham, A. Kalogeropoulos, D. Lange, J. Luo, D. Marlow, K. Mei, I. Ojalvo, J. Olsen, C. Palmer, P. Piroué, D. Stickland, C. Tully

#### University of Puerto Rico, Mayaguez, USA

S. Malik, S. Norberg

#### Purdue University, West Lafayette, USA

A. Barker, V.E. Barnes, S. Das, S. Folgueras, L. Gutay, M. Jones, A.W. Jung, A. Khatiwada, D.H. Miller, N. Neumeister, C.C. Peng, H. Qiu, J.F. Schulte, J. Sun, F. Wang, R. Xiao, W. Xie

#### Purdue University Northwest, Hammond, USA

T. Cheng, N. Parashar, J. Stupak

#### Rice University, Houston, USA

Z. Chen, K.M. Ecklund, S. Freed, F.J.M. Geurts, M. Guilbaud, M. Kilpatrick, W. Li, B. Michlin, B.P. Padley, J. Roberts, J. Rorie, W. Shi, Z. Tu, J. Zabel, A. Zhang

#### University of Rochester, Rochester, USA

A. Bodek, P. de Barbaro, R. Demina, Y.t. Duh, T. Ferbel, M. Galanti, A. Garcia-Bellido, J. Han, O. Hindrichs, A. Khukhunaishvili, K.H. Lo, P. Tan, M. Verzetti

#### The Rockefeller University, New York, USA

R. Ciesielski, K. Goulianos, C. Mesropian

#### Rutgers, The State University of New Jersey, Piscataway, USA

A. Agapitos, J.P. Chou, Y. Gershtein, T.A. G´omez Espinosa, E. Halkiadakis, M. Heindl, E. Hughes, S. Kaplan, R. Kunnawalkam Elayavalli, S. Kyriacou, A. Lath, R. Montalvo, K. Nash, M. Osherson, H. Saka, S. Salur, S. Schnetzer, D. Sheffield, S. Somalwar, R. Stone, S. Thomas, P. Thomassen, M. Walker

#### University of Tennessee, Knoxville, USA

A.G. Delannoy, J. Heideman, G. Riley, K. Rose, S. Spanier, K. Thapa

#### Texas A&M University, College Station, USA

O. Bouhali<sup>72</sup>, A. Castaneda Hernandez<sup>72</sup>, A. Celik, M. Dalchenko, M. De Mattia, A. Delgado, S. Dildick, R. Eusebi, J. Gilmore, T. Huang, T. Kamon<sup>73</sup>, R. Mueller, Y. Pakhotin, R. Patel, A. Perloff, L. Perniè, D. Rathjens, A. Safonov, A. Tatarinov, K.A. Ulmer

#### Texas Tech University, Lubbock, USA

N. Akchurin, J. Damgov, F. De Guio, P.R. Dudero, J. Faulkner, E. Gurpinar, S. Kunori, K. Lamichhane, S.W. Lee, T. Mengke, S. Muthumuni, T. Peltola, S. Undleeb, I. Volobouev, Z. Wang

#### Vanderbilt University, Nashville, USA

S. Greene, A. Gurrola, R. Janjam, W. Johns, C. Maguire, A. Melo, H. Ni, K. Padeken, P. Sheldon, S. Tuo, J. Velkovska, Q. Xu

#### University of Virginia, Charlottesville, USA

M.W. Arenton, P. Barria, B. Cox, R. Hirosky, M. Joyce, A. Ledovskoy, H. Li, C. Neu, T. Sinthuprasith, Y. Wang, E. Wolfe, F. Xia

#### Wayne State University, Detroit, USA

R. Harr, P.E. Karchin, N. Poudyal, J. Sturdy, P. Thapa, S. Zaleski

#### University of Wisconsin - Madison, Madison, WI, USA

M. Brodski, J. Buchanan, C. Caillol, D. Carlsmith, S. Dasu, L. Dodd, S. Duric, B. Gomber,

M. Grothe, M. Herndon, A. Hervé, U. Hussain, P. Klabbers, A. Lanaro, A. Levine, K. Long,

R. Loveless, V. Rekovic, T. Ruggles, A. Savin, N. Smith, W.H. Smith, N. Woods

- 1: Also at Vienna University of Technology, Vienna, Austria
- 2: Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France
- 3: Also at Universidade Estadual de Campinas, Campinas, Brazil
- 4: Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil
- 5: Also at Université Libre de Bruxelles, Bruxelles, Belgium
- 6: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia
- 7: Also at Joint Institute for Nuclear Research, Dubna, Russia
- 8: Also at Cairo University, Cairo, Egypt
- 9: Also at Suez University, Suez, Egypt
- 10: Now at British University in Egypt, Cairo, Egypt
- 11: Now at Ain Shams University, Cairo, Egypt
- 12: Also at Department of Physics, King Abdulaziz University, Jeddah, Saudi Arabia
- 13: Also at Université de Haute Alsace, Mulhouse, France
- 14: Also at Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia
- 15: Also at Tbilisi State University, Tbilisi, Georgia
- 16: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- 17: Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- 18: Also at University of Hamburg, Hamburg, Germany
- 19: Also at Brandenburg University of Technology, Cottbus, Germany
- 20: Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- 21: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- 22: Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
- 23: Also at Indian Institute of Technology Bhubaneswar, Bhubaneswar, India
- 24: Also at Institute of Physics, Bhubaneswar, India
- 25: Also at Shoolini University, Solan, India
- 26: Also at University of Visva-Bharati, Santiniketan, India
- 27: Also at University of Ruhuna, Matara, Sri Lanka
- 28: Also at Isfahan University of Technology, Isfahan, Iran
- 29: Also at Yazd University, Yazd, Iran
- 30: Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran
- 31: Also at Università degli Studi di Siena, Siena, Italy
- 32: Also at INFN Sezione di Milano-Bicocca; Università di Milano-Bicocca, Milano, Italy
- 33: Also at International Islamic University of Malaysia, Kuala Lumpur, Malaysia
- 34: Also at Malaysian Nuclear Agency, MOSTI, Kajang, Malaysia
- 35: Also at Consejo Nacional de Ciencia y Tecnología, Mexico city, Mexico
- 36: Also at Warsaw University of Technology, Institute of Electronic Systems, Warsaw, Poland
- 37: Also at Institute for Nuclear Research, Moscow, Russia
- 38: Now at National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia

<sup>†</sup>: Deceased

- 39: Also at St. Petersburg State Polytechnical University, St. Petersburg, Russia
- 40: Also at University of Florida, Gainesville, USA
- 41: Also at P.N. Lebedev Physical Institute, Moscow, Russia
- 42: Also at California Institute of Technology, Pasadena, USA
- 43: Also at Budker Institute of Nuclear Physics, Novosibirsk, Russia
- 44: Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia
- 45: Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia
- 46: Also at Scuola Normale e Sezione dell'INFN, Pisa, Italy
- 47: Also at National and Kapodistrian University of Athens, Athens, Greece
- 48: Also at Riga Technical University, Riga, Latvia
- 49: Also at Universität Zürich, Zurich, Switzerland
- 50: Also at Stefan Meyer Institute for Subatomic Physics (SMI), Vienna, Austria
- 51: Also at Adiyaman University, Adiyaman, Turkey
- 52: Also at Istanbul Aydin University, Istanbul, Turkey
- 53: Also at Mersin University, Mersin, Turkey
- 54: Also at Piri Reis University, Istanbul, Turkey
- 55: Also at Gaziosmanpasa University, Tokat, Turkey
- 56: Also at Izmir Institute of Technology, Izmir, Turkey
- 57: Also at Necmettin Erbakan University, Konya, Turkey
- 58: Also at Marmara University, Istanbul, Turkey
- 59: Also at Kafkas University, Kars, Turkey
- 60: Also at Istanbul Bilgi University, Istanbul, Turkey
- 61: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
- 62: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- 63: Also at Monash University, Faculty of Science, Clayton, Australia
- 64: Also at Instituto de Astrofísica de Canarias, La Laguna, Spain
- 65: Also at Utah Valley University, Orem, USA
- 66: Also at Purdue University, West Lafayette, USA
- 67: Also at Beykent University, Istanbul, Turkey
- 68: Also at Bingol University, Bingol, Turkey
- 69: Also at Erzincan University, Erzincan, Turkey
- 70: Also at Sinop University, Sinop, Turkey
- 71: Also at Mimar Sinan University, Istanbul, Istanbul, Turkey
- 72: Also at Texas A&M University at Qatar, Doha, Qatar
- 73: Also at Kyungpook National University, Daegu, Korea