

Reduced tillage, but not organic matter input, increased nematode diversity and food web stability in European long-term field experiments

Journal Article**Author(s):**

Bongiorno, Giulia; Bodenhausen, Natacha; Bünemann, Else K.; Brussaard, Lijbert; Geisen, Stefan; Mäder, Paul; Quist, Casper W.; Walser, Jean-Claude; de Goede, Ron G. M.

Publication date:

2019-11

Permanent link:

<https://doi.org/10.3929/ethz-b-000377318>

Rights / license:




[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#)

Originally published in:

Molecular Ecology 28(22), <https://doi.org/10.1111/mec.15270>

ORIGINAL ARTICLE

Reduced tillage, but not organic matter input, increased nematode diversity and food web stability in European long-term field experiments

Giulia Bongiorno^{1,2}  | Natacha Bodenhausen² | Else K. Bünemann² |
Lijbert Brussaard¹ | Stefan Geisen³  | Paul Mäder² | Casper W. Quist^{4,5}  |
Jean-Claude Walser⁶ | Ron G. M. de Goede¹

¹Soil Biology Group, Wageningen University & Research, Wageningen, The Netherlands

²Department of Soil Sciences, Research Institute of Organic Agriculture (FiBL), Frick, Switzerland

³Department of Terrestrial Ecology, Netherland Institute of Ecology, Wageningen, The Netherlands

⁴Laboratory of Nematology, Wageningen University & Research, Wageningen, The Netherlands

⁵Biosystematics Group, Wageningen University & Research, Wageningen, The Netherlands

⁶Genetic Diversity Centre, ETH Zürich, Zürich, Switzerland

Correspondence

Giulia Bongiorno, Soil Biology Group, Wageningen University & Research, Wageningen, The Netherlands and Department of Soil Science, Research Institute of Organic Agriculture (FiBL), Frick, Switzerland.

Emails: giulia.bongiorno@wur.nl; giulia.bongiorno@fiBL.org

Funding information

Netherlands Organisation for Scientific Research, Grant/Award Number: 016.Veni.181.078; European Union Horizon 2020, Grant/Award Number: 635750

Abstract

Soil nematode communities and food web indices can inform about the complexity, nutrient flows and decomposition pathways of soil food webs, reflecting soil quality. Relative abundance of nematode feeding and life-history groups are used for calculating food web indices, i.e., maturity index (MI), enrichment index (EI), structure index (SI) and channel index (CI). Molecular methods to study nematode communities potentially offer advantages compared to traditional methods in terms of resolution, throughput, cost and time. In spite of such advantages, molecular data have not often been adopted so far to assess the effects of soil management on nematode communities and to calculate these food web indices. Here, we used high-throughput amplicon sequencing to investigate the effects of tillage (conventional vs. reduced) and organic matter addition (low vs. high) on nematode communities and food web indices in 10 European long-term field experiments and we assessed the relationship between nematode communities and soil parameters. We found that nematode communities were more strongly affected by tillage than by organic matter addition. Compared to conventional tillage, reduced tillage increased nematode diversity (23% higher Shannon diversity index), nematode community stability (12% higher MI), structure (24% higher SI), and the fungal decomposition channel (59% higher CI), and also the number of herbivorous nematodes (70% higher). Total and labile organic carbon, available K and microbial parameters explained nematode community structure. Our findings show that nematode communities are sensitive indicators of soil quality and that molecular profiling of nematode communities has the potential to reveal the effects of soil management on soil quality.

KEYWORDS

amplicon sequencing, food web indices, long-term field experiments, nematode communities, organic matter addition, tillage

1 | INTRODUCTION

The capacity of soils to perform multiple processes defines and determines soil quality (Bünemann et al., 2018). Soil management can negatively affect soil processes exerting threats (e.g., soil erosion, compaction, acidification and organic matter losses) on chemical, physical and biological properties (Toth, Montanarella, & Rusco, 2008). Tillage and fertilization are widespread soil management measures which can have a substantial influence on these soil threats, ultimately affecting soil processes and soil quality.

Soil nematodes are abundant and ubiquitous organisms that have an important role in various processes such as nutrient cycling, decomposition, pest and pathogen population regulation (Ekschmitt et al., 2001; Neher, Weicht, & Barbercheck, 2012). In soils, nematodes are present at all trophic levels, and can therefore be divided into functional groups based on their feeding preferences (Yeates, Bongers, Goede, Freckman, & Georgieva, 1993). Nematodes can also be differentiated according to their life-history strategies reflected in their position on a colonizer-persister (c-p) scale, which goes from group 1 (colonizers = r selected species) to group 5 (persisters = K selected species; Bongers, 1990). Colonizers thrive in nutrient-rich habitats, are generally bacterivores, tolerant to stress and pollutants, with short generation times, while persisters poorly react to conditions of high food availability, are bigger omnivorous and/or predatory nematodes sensitive to stress, have longer generation times and generally live in temporally stable habitat. Many species have intermediate characteristics. Relative abundance of nematode feeding and life-history groups are used for calculating food web indices, i.e., the maturity index (MI: measure of environmental disturbance), enrichment index (EI: measure of resource availability), structure index (SI: measure of degree of trophic links and capacity to recover from stress) and channel index (CI: indication of predominantly fungal or bacterial decomposition pathway; Bongers, 1990; Ferris, Bongers, & Goude, 2001), which are used to determine soil processes affecting soil quality.

Due to interactions with other soil biota and the influence of chemical and physical abiotic factors (Bongers & Ferris, 1999), changes induced by soil management affect nematode communities (Ferris & Bongers, 2006; Sánchez-Moreno, Nicola, Ferris, & Zalom, 2009). These changes in the nematode community can be due to modifications in food resources such as plant residues, nutrients, and environmental properties such as pH, oxygen content, porosity and temperature (Mekonen, Petros, & Hailemariam, 2017; Yeates & Bongers, 1999). Thus, data on nematode communities integrate information from soil chemical, physical and biological properties (Mekonen et al., 2017; Neher, 2001). This can increase our understanding of the impact of soil management on soil processes and, indeed, on soil quality in general.


Nematode diversity and specific nematode groups (i.e., based on feeding and/or life-history strategies) or taxa (i.e., family, genus, or species) have been shown to respond differently to soil management such as tillage and fertilization (Moura & Franzener, 2017; Yeates & Bongers, 1999). More in detail, previous studies found higher

nematode diversity and higher percentages of fungal feeders, omnivores and predators (slow-growing nematodes of c-p groups 4 and 5) in less disturbed conditions such as systems under reduced tillage or with perennial crops (Liu et al., 2016; Niles & Freckman, 1998; Yeates & Bongers, 1999). In contrast, fast-growing bacterivorous nematodes (c-p groups 1 and 2) have been associated with eutrophic and mineral fertilized, disturbed systems (Darby, Todd, & Herman, 2013; De Goede, Bongers, & Ettema, 1993; Quist et al., 2016; Zhao & Neher, 2013). Also the application of different organic materials such as manure, compost and cattle slurry has been shown to increase the abundance of bacterivorous nematodes (Forge, Bittman, & Kowalenko, 2005; Leroy, Bommele, Reheul, Moens, & De Neve, 2007), and, in some cases, to decrease the abundance of plant parasitic nematodes (Leroy et al., 2007).

In most publications so far, the response of nematode communities to tillage and fertilization was studied in single field experiments (Ito, Araki, Komatsuzaki, Kaneko, & Ohta, 2015; Quist et al., 2016; Zhao & Neher, 2013), sometimes yielding contradictory results (Ferris et al., 2012; Leroy et al., 2007; Treonis et al., 2018). One factor hampering the study of management effects across multiple study sites is that traditional microscopy is the most common method to study nematodes, which is time-consuming, requires specialists and is expensive (Ritz, Black, Campbell, Harris, & Wood, 2009). Molecular methods to assess nematode absolute abundances (qPCR) and diversity (high-throughput amplicon sequencing, DGGE, T-RFLP) are faster, cheaper, and allow higher throughput than visual methods (Ahmed, Sapp, Prior, Karssen, & Back, 2016; Geisen et al., 2018). Amplicon sequencing may allow identification of taxa that cannot be distinguished morphologically. One limitation of PCR-based molecular methods is that not actual abundances of the specimen but rather their relative number of DNA copies are assessed (Porazinska et al., 2009; Waite et al., 2003). However, there is recent evidence that molecular methods might give similar ecological patterns as traditional methods (Geisen et al., 2018; George & Lindo, 2015; Hamilton, Strickland, Wickings, Bradford, & Fierer, 2009; Porazinska, Sung, Giblin-Davis, & Thomas, 2010; Quist et al., 2016). Hence, amplicon sequencing has high potential to assess soil management effects on nematode communities across multiple field experiments.

The goal of the present study was to: (a) Assess the effect of tillage and organic matter addition on nematode qPCR counts, alpha and beta-diversity, and food web indices as measured by amplicon sequencing of the 18S rRNA gene; (b) investigate the relationships between nematode community characteristics and other soil parameters related to soil processes; and (c) identify taxa that could serve as indicator organisms for soil management. We expected that molecular techniques would be sensitive, efficient tool to reveal general patterns of soil management effects on nematode communities in 10 long-term field experiments across Europe. We hypothesized that (a) reduced tillage would increase nematode qPCR counts, alpha diversity, MI, SI and CI, and decrease levels of bacterivorous nematodes with short life-cycles compared to conventional tillage, and that (b) high organic matter addition would increase qPCR counts, alpha diversity, EI, and alter the

FIGURE 1 Main pedoclimatic characteristics and soil management (tillage, organic matter input, or a combination of the two) of 10 long-term field experiments analysed in the current study. CH1, Frick trial; CH2, Aesch trial; CH3, DOK trial; ES4, Pago trial; HU1, Keszthely trial; HU4, Keszthely trial; NL1, Basis trial; NL2, De Peel trial; PT1, Vitichar trial; SL1 Tillorg trial. For detailed information about the experiments see Table S1



nematode communities towards higher populations of bacterivorous nematodes compared to low organic matter input. We also hypothesized that (c) the positive effect of reduced tillage and organic matter addition on total and labile organic matter, available nutrients, water stable aggregates, and microbial biomass and activity would result in a positive relationship between these soil parameters and the nematode communities and that (d) nematode taxa with long life cycles and sensitive to management (such as predatory and omnivorous nematodes in c-p groups 4 and 5) would be more associated with less disturbed systems, and as such would be sensitive indicator taxa for soil disturbance.

2 | MATERIALS AND METHODS

2.1 | Long-term field experiments and management

We selected 10 European long-term field experiments with either arable or permanent crops and a minimum duration of 5 years and a maximum duration of 44 years (Figure 1, Table S1). Throughout the paper we will refer to these long-term field experiments as “LTEs”.

This selection covered five different European climatic zones (Köppen, 1918; Figure 1, Table S1) and six soil textural classes (Table S1; WRB, 2014).

Each LTE had unique management characteristics and a different experimental design, with three or four replicates per treatment (Table S1). However, LTEs were comparable because the main soil management types were tillage (T) and organic matter addition (OM) as described in Bongiorno, Bünemann, et al.

(2019). The contrast in tillage was classified as conventional tillage (ploughing at 20–25 cm depth, CT) versus reduced tillage (no-tillage or noninversion tillage at 0–10 cm with different light machinery, RT). The contrast in organic matter addition was classified as low organic matter addition (LOW, no organic matter additions or only mineral fertilization) versus high organic matter addition (HIGH, organic matter additions without or with mineral fertilizer). At some LTEs, both treatment factors (i.e., tillage and organic matter addition) were applied and at others only one of these was present (Figure 1).

2.2 | Sampling procedure and sample handling

A total of 167 soil samples were collected in spring 2016 before any major soil or crop management was started in the LTEs. Each sample consisted of a composite sample of 20 soil cores randomly collected in the central area of the plot, to avoid border effects, and mixed. In the tilled LTEs, samples were taken from two depths: 0–10 cm and 10–20 cm. In the LTEs with organic matter addition as the only management factor (no tillage factor), samples were taken from the 0–20 cm layer because we did not expect to find a stratification effect due to tillage. After soil sampling, 400 g of the samples were air-dried (40°C) for subsequent chemical analysis. Fresh soil samples were sent to Wageningen University (The Netherlands), Research Institute of Organic Agriculture (Frick, Switzerland), University of Trier (Germany) and University Miguel Hernandez (Alicante, Spain), and air-dried samples were sent to University of Ljubljana (Slovenia).

Upon arrival, the samples were sieved at 5 mm and, when fresh, stored at 3°C until further processing.

2.3 | Chemical, physical and biological soil properties

The following soil properties were measured for this study: total organic carbon (TOC: %), pH (CaCl₂), total nitrogen (TN: %), cation exchange capacity (CEC: mmol 100 g soil), plant available phosphorus (P: mg/kg soil), plant available potassium (K: mg/kg soil), exchangeable magnesium, calcium, and sodium (Mg²⁺, Ca²⁺, Na⁺; mg/kg soil), water-stable aggregates (WSA: mg/kg soil), water holding capacity (WHC: %), bulk density (BD: g/cm³), percentages of silt, clay, and sand, microbial biomass carbon (MBC: mg/kg soil), microbial biomass nitrogen (MBN: mg/kg soil), soil respiration (SR: μg CO₂-C hr⁻¹ g⁻¹ soil), number and biomass of earthworms (number and g/m²), decomposition through tea bag index (% mass loss) and soil suppressiveness to *Pythium ultimum* (%; Bongiorno, Postma, et al., 2019). Microbial quotient (qMic) and metabolic quotient (qCO₂) were calculated as the microbial biomass carbon divided by the total organic carbon, and the soil respiration divided by the microbial biomass carbon, respectively. Besides chemical, physical and biological parameters, five different labile carbon fractions were measured: hydrophilic dissolved organic carbon (Hy-DOC: mg/kg soil), dissolved organic carbon (DOC: mg/kg soil), permanganate oxidizable carbon (POXC: mg/kg soil), hot water extractable carbon (HWEC: mg/kg soil), and particulate organic matter carbon (POMC: mg/kg soil; Bongiorno, Bünemann, et al., 2019). In addition, the specific ultraviolet absorbance of Hy (Hy SUVA: L g C⁻¹ cm⁻¹) and DOC (DOC SUVA: L g C⁻¹ cm⁻¹) was measured to assess the recalcitrance of these labile carbon fractions. All analyses were performed within 6 months after sampling and the details of the methodology and locations where the analyses took place are presented in Table S2 (modified from Bongiorno, Postma, et al., 2019).

2.4 | Nematode analysis

2.4.1 | Nematode extraction, DNA extraction and DNA purification

Within 2 weeks after sampling nematodes were extracted from 100 g field moist subsamples using a modified elutriator (Oostenbrink, 1960). Thereafter nematodes were incubated for 72 hr on a double cotton-wool filter (Hygia milac). A subset of samples from each LTE (a total of 97 samples) was counted microscopically, with 1/10 of each sample counted in duplicate under a dissecting microscope. The number of nematodes was expressed per 100 g of field moist soil. The nematode suspensions were subsequently concentrated and lysed with a lysis buffer containing proteinase K, β-mercaptoethanol and an internal mammalian standard in order to correct for the loss of DNA during lysis and DNA purification (Holterman et al., 2006; Vervoort et al., 2012). Thereafter, DNA extracts were purified using a glass fibre

column-based procedure (Ivanova, Dewaard, & Hebert, 2006) and stored at -20°C until further use.

2.4.2 | Quantitative PCR (qPCR) analysis of total nematode DNA

The purified DNA extracts were used as templates in qPCR using two primer sets to assess total nematode densities (Quist et al., 2017; Vervoort et al., 2012). The first primer set targeted DNA across the phylum Nematoda and the second targeted the mammalian internal standard. After the qPCR reactions, the C_t-values obtained were related to the microscopic counts to obtain a calibration curve at the ¹⁰Log scale (see Vervoort et al., 2012). Thereafter, C_t-values were converted into nematode densities using this linear relationship between the C_t values and the ¹⁰Log (number of target nematodes; Figure S1a). The maxima of the negative, first mathematic derivative of the melting curves were checked to confirm the correct nature of the amplicons. The internal control was used to monitor and correct for loss of DNA during the sampling handling. Throughout the manuscript qPCR-based quantification of nematode densities is referred to as "nematode qPCR counts".

2.4.3 | 18S rRNA gene amplification and sequencing

Nematode DNA was quantified with Nanodrop (NanoDrop 2000 Spectrophotometer, Thermo Fischer Scientific) and subsequently sent on dry ice to GenomeQuebec for 18S rRNA gene amplification and sequencing on the Illumina MiSeq platform. In a first step a targeted PCR amplification with tagged primers for the hypervariable eukaryotic V4 region of the 18S rRNA gene was performed (Table S3). We used the universal eukaryotic primers 3NDf (5'-GGCAAGTCTGGTGCCAG-3') in combination with 1132rmod (5'-TCCGTC AATTYCTTTAAGT-3') as used in Geisen et al. (2018). In a next step, Illumina adapters with barcodes sequences were added by PCR to each sample (barcoding step; Table S3). For each sample, the barcoding step was verified with gel electrophoresis. The DNA concentration was quantified with Quant-iT PicoGreen dsDNA Assay kit (Life Technologies) and for each sample, an equal amount of DNA was pooled for a sequencing library. After purification with AMPure beads (Beckman Coulter), the pooled DNA library was quantified using the Quant-iT PicoGreen dsDNA Assay kit (Life Technologies) and the Kapa Illumina GA Library Quantification kit with revised primers (KAPA SYBR FAST qPCR Universal kit, Kapa Biosystems). Average fragment size was determined using a LabChip GX (PerkinElmer) instrument. Sequencing was performed with MiSeq Reagent kit v3 (600 cycles) from Illumina. After sequencing, the sequences were demultiplexed by GenomeQuebec using the Illumina bcl2fastq Conversion Software version 2.17.1.14.

2.4.4 | Bioinformatic analysis

The amplicon sequencing data was analyzed by the Genetic Diversity Centre (GDC), ETH Zurich, using the HPC Euler of ETH Zurich. The merging efficiency of the forward (R1) and reverse (R2) reads was

relatively low (<11%). For this reason, we restricted the analysis to the forward read only. In a first step, the primer sites were trimmed off the R1 reads and all the reads were trimmed to an equal length of 280 nt using USEARCH (Edgar, 2010). Subsequently, the reads were quality filtered (parameter: GC range 20–80, minimum quality mean 20, no ambiguous nucleotides, and a low complexity filter, dust with threshold 30) using PRINSEQ-LITE (version 0.20.4). About 10% of the total sequencing data was lost during primer trimming (7.5%), trimming (<1%), and quality filtering (2.6%). In a next step, UPARSE (Edgar, 2013) was used to cluster the sequences and create a count table. For the annotation of the OTUs SINTAX (Edgar, 2016) and the protist ribosomal reference database (PR2) were used. The OTUs which could not be assigned to a taxonomic group were verified with manual BLAST searches with NCBI nt based references databases (see Figure S2).

2.4.5 | Nematode alpha diversity, trophic groups and food web indices

Alpha diversity is defined as the diversity of organisms within groups (in our case calculated within plots), while beta diversity is defined as the diversity of organisms between groups (Jost, 2010). Alpha diversity is measured through indices of richness, diversity and evenness (Jost, 2010). Nematode OTU or genus richness was calculated as the sum of the OTUs or genera, respectively. Nematode OTU and genus diversity was calculated as the exponential of the Shannon Index (Magurran, 1988):

$$\exp^H = \exp^{\sum_{i=1}^S - (P_i \cdot \ln P_i)} \quad (1)$$

where H is the Shannon diversity index, P_i is the fraction of the entire population made of OTU or genus i , S is the number of OTUs or genera encountered, and \sum is the sum of OTU or genus 1 to OTU or genus S . Nematode OTU and genus evenness (Sheldon evenness) was calculated as the exponential of the Shannon diversity divided by the number of OTUs or genera (Heip, 1974).

We calculated the percentages of five trophic groups (bacterivorous, fungivorous, herbivorous, predators and omnivorous nematodes), maturity index (MI), enrichment index (EI), structure index (SI), and channel index (CI), according to the classification of nematode OTUs into functional groups, uploading the count table based on OTU observed abundance with taxonomic information obtained after the bioinformatic analysis of the nematode sequencing data in the nematode indicator joint analysis (NINJA) program (Sieriebriennikov, Ferris, & de Goede, 2014; <http://sieriebriennikov.shinyapps.io/ninja/> consulted on 9 January 2019). NINJA was used also to assign nematodes to the colonizer-persister (c-p) scale (from 1 to 5; Bongers, 1990; Ferris et al., 2001). The absolute abundance of trophic groups and c-p groups was calculated by multiplying the total qPCR counts by the trophic and the c-p groups percentages calculated with NINJA.

2.5 | Statistical analysis

All statistical calculations were carried out using R version 3.5.1 and RSTUDIO version 1.1.456 (R Development Core Team, 2013; RStudio

Team, 2016). The R script is provided as Appendix S2, and a workflow of the data analysis steps is given in Figure S3. The nematode OTU counts and taxonomy tables were filtered before the analysis to exclude OTUs which were classified as non-nematodes, or whose kingdom or phylum was unassigned. All test results, except for the indicator species analysis, were considered statistically significant at $p \leq .05$.

2.5.1 | Nematode qPCR counts, alpha and beta diversity per LTE

Nematode OTU richness and diversity were calculated after rarefaction (500×) to 10,537 seq/sample (the minimum sample sequencing depth; Bodenhause, Horton, & Bergelson, 2013).

A general beta diversity analysis was conducted on the nematode communities of all the sites. For this analysis, we filtered the OTU sequence counts retaining only OTUs with a minimum of five counts in at least eight samples. After normalization using the total sum scaling (TSS) with the decostand (method = "total") function in the VEGAN package (Oksanen et al., 2018), we computed Bray-Curtis dissimilarity matrices on the squared rooted transformed data (Leff et al., 2015). Canonical analysis of proximities (CAP) with vegan function capscale was performed to visualize and test the relationships between the nematode community and the most important soil chemical, physical and biological parameters measured in the LTEs (Anderson & Willis, 2003). The function vif.cca (threshold used $vif \leq 10$) was used to retain variables which were not highly correlated ($\rho > 0.80$). The effect of the environmental variables on the nematode communities was assessed with permutation analysis (using the anova function in VEGAN by "margin") with 10^4 permutations and correlations between the environmental variables and the first two axes of the CAP to assign their relative importance.

2.5.2 | Effects of tillage and organic matter additions on nematode qPCR counts, alpha and beta diversity

To test the effects of tillage and organic matter addition on soil nematode communities, two groups of LTEs were created because we expected stratification effects in LTEs with reduced tillage only, as shown in previous analyses (Bongiorno, Bünemann, et al., 2019). The following two groups were studied separately in the subsequent analyses:

Group A

The LTEs in which the layers 0–10 cm and 10–20 cm were sampled separately in space: CH1, CH2, NL1, NL2, SL1, HU4 and ES4. In this group, we assessed the effect of tillage, organic matter addition and soil layer.

Group B

The LTEs where the layer 0–20 was sampled: CH3, PT1 and HU1. In this group we only assessed the effect of organic matter addition, since these LTEs were under conventional tillage.

The effect of tillage and/or organic matter addition and, if present, layer on total nematode qPCR counts, OTU and genus richness and diversity, and OTU evenness were assessed in group A and B (using overall models merging the LTEs in the same group) by performing an analysis of variance (standard function anova) on fitted linear mixed effect models. Mixed models were used to take into account the possible correlations introduced by the multisite field experiments and to generalize the effect of the soil management practices across the different LTEs (Bongiorno, Bünemann, et al., 2019). The tillage and/or the soil organic matter addition and, if present, the layer, their two-way and possibly three-way interactions were used as fixed factors. Random effects for LTEs, blocks, main plots and subplots were introduced in the models to represent the experimental designs of the different LTEs. The effect of the pedoclimatic zone was not included in the fixed part of the model because we were interested in management effects across pedoclimatic zones. The model assumptions of normality and homogeneity of variances of the residuals were checked both visually and with the Shapiro-Wilk and Levene's tests (Zuur, 2009). Total nematode qPCR counts and OTU richness, diversity and evenness were square-root-transformed in order to meet the assumption of normality. All tests were considered statistically significant at $p \leq .05$. For the linear mixed effects model, the packages nlme, and emmeans were used (Pinheiro, Bates, DebRoy, & Sarkar, 2018). The same linear mixed effect models were used to assess differences in relative and absolute abundances of trophic and c-p groups, and in food web indices between soil management.

We then performed multivariate analysis of nematode communities on Bray-Curtis dissimilarities as outlined by Anderson and Willis (2003) using squared-root TSS normalized data. Using a permutational multivariate analysis of variance (PERMANOVA) with 10^4 permutations we tested the effect of tillage and/or organic matter and, if present, the layer on the community dissimilarity. In this analysis, the LTE was specified as random factor in the strata argument which restricts permutations to within LTEs (Anderson, 2001). The function betadisp was used to perform permutational analysis of multivariate dispersion (BETADISP) with 10^4 permutations.

We then visualized the effect of soil management with canonical analysis of proximities (CAP) constrained ordination (Anderson & Willis, 2003) using the function capscale in the vegan package with the LTE as a conditional factor in order to control for the effect of the pedoclimatic zone on the nematode communities. Statistical significance of the CAP was assessed using the permutest function in the VEGAN package.

2.5.3 | Relationships between nematodes and soil parameters

Partial correlations, correcting for the variation caused by the intrinsic differences between the LTEs (pedoclimatic zones), were used to test the relationships between nematode qPCR counts, OTU richness, diversity and evenness and the soil chemical, physical and

biological parameters. For the correlation analyses the packages CAR, STATS and PPCOR were used (Kim, 2015).

The relationships between nematode communities and environmental variables shaped by the effect of the soil management practices was visualised using canonical analysis of proximities (CAP) and tested using the envfit function in the package VEGAN. The effect of the soil parameters was assessed with permutation analysis with 10^4 permutations.

2.5.4 | Identification of putative indicator OTUs

Determination of nematode OTUs associated with specific management combinations was done using correlation-based indicator analysis with the function multipatt of the R package indicpecies (De Caceres, 2016) to calculate the point-biserial correlation coefficient (r) of an OTU's positive association to a soil management factor or a combination of factors. The analysis was done with 10^4 permutations and considered a more stringent significance level at $p \leq .01$, in order to limit the indicator species to a subgroup of highly sensitive OTUs associated with soil management. In the analysis we restricted the permutation within the blocks and within the LTEs to take into account the nested structure of the design.

3 | RESULTS

3.1 | Nematode beta diversity across the long-term field experiments

In the CAP, the community composition showed a clustering of samples according to the long-term field experiments (LTEs; Figure 2), and PERMANOVA confirmed that the nematode communities were affected by the LTE ($R^2 = .64$; $p = .001$). A total of 50% of variation in the nematode beta diversity among the different LTEs was explained by the constraining variables used in the CAP.

According to ANOVA of the constraining variables, all the soil parameters were significantly related to the nematode beta diversity in the LTEs (Table S4). The soil parameters that were most important in explaining the variation between the different LTEs (i.e., significant relationship and Pearson correlation coefficient (r) with the canonical axes greater than +0.50 or smaller than -0.50) were for CAP1: sand content, pH, microbial biomass carbon (MBC), cation exchange capacity (CEC), and total nitrogen (TN); for CAP2: permanganate oxidizable carbon (POXC), water stable aggregates (WSA), and total nitrogen (TN).

3.2 | Effect of soil management on total nematode qPCR counts and alpha diversity

In group A (i.e., LTEs with tillage and organic matter addition as treatments, sampled at two soil depths), nematode qPCR counts were higher in the first layer (0–10 cm) than in the second layer (10–20 cm; Table 1). We found higher nematode OTU richness, diversity, and evenness and genus diversity and evenness in reduced tillage compared to conventional tillage across the LTEs of group A. In this




FIGURE 2 Constrained analysis of proximities (CAP) of the nematode communities in the long-term field experiments and the relation with soil parameters. The first axis, CAP1 explains 16.7% and the second axis explains 10.6% of the variation in the beta diversity between the nematode communities in the different sites. BD, bulk density; CEC, cation exchange capacity; HWEC, hot water extractable carbon; K, available potassium; MBC, microbial biomass carbon; Mg, magnesium; pH, potential of hydrogen; POXC, permanganate oxidizable carbon; Sand, sand; TN, total nitrogen; WSA, water stable aggregates. CH1, Frick trial; CH2, Aesch trial; CH3, DOK trial; ES4, Pago trial; HU1, Keszthely trial; HU4, Keszthely trial; NL1, Basis trial; NL2, De Peel trial; PT1, Vitichar trial; SL1, Tillorg trial [Colour figure can be viewed at wileyonlinelibrary.com]

analysis, OTU richness and diversity, and genus richness had higher values in the upper than in the lower layer, regardless of the tillage treatment (OTU richness and diversity 11% and 18% higher, respectively, and genus richness 9% higher). OTU and genus diversity and evenness were lower (16% and 22% for the OTU and 28% and 28% for genus, respectively) in the high organic matter addition plots.

In group B (i.e., LTEs with organic matter addition only, sampled between 0–20 cm soil depth), we found no significant effects of organic matter addition on total nematode qPCR counts, OTU and genus richness and diversity (Table 1).

3.3 | Effect of soil management on beta diversity

PERMANOVA of group A revealed that the largest proportion of the variation in nematode beta diversity was explained by the LTEs ($R^2 = .628$, $p = .0001$). Despite this, tillage ($R^2 = .012$, $p = .0001$), organic matter addition ($R^2 = .006$, $p = .006$), layer ($R^2 = .014$, $p = .0001$) and the interaction between tillage and layer ($R^2 = .006$, $p = .002$) had significant effects on the nematode beta diversity (Figure 3a, Table S5). The significant interaction between tillage and layer indicates that under reduced tillage a significant effect of the layer was found, but not under conventional tillage.

The CAP model of group A explained in total 8% of the variation in beta diversity related to soil management (tillage, organic matter addition), and the first two axes explained 2.6% and 2.3% of variation, respectively. CAP1 axis separated the samples belonging to the lower layer of reduced tillage from the rest, while CAP2 axis, from top to bottom, separated the different tillage treatments.

In group B, PERMANOVA did not reveal effects of organic matter addition ($R^2 = .013$, $p = .186$) on the nematode beta diversity (Table S5).

The dispersion tests were not significant, suggesting that differences between management were driven primarily by true biological differences and not by an artefact of the differences of the within-group dispersion (Table S6).

3.4 | Effect of soil management on nematode trophic groups and food web indices

Bacterivorous nematodes were the most abundant trophic group, followed by herbivorous, fungivorous, omnivorous and predatory nematodes (Table 2, Table S7). For group A, we found a stratification effect of reduced tillage on relative abundance of bacterivorous nematodes, with lower values in the lower than in the upper layer (24% lower, $p = .0005$; Figure S4). The proportion of herbivorous nematodes was higher in the lower layer of reduced tillage (44%) compared to the upper layer of reduced tillage (19%) and both layers of conventional tillage (16% and 19% for higher and lower layer, respectively; $p = .0004$; Figure S4). Its absolute abundance was 70% higher in reduced tillage compared to conventional tilled treatment ($p = .007$), both in the 0–10 and 10–20 cm soil layer and regardless of organic matter. There was a 44% higher proportion of fungivorous nematodes in the upper layer of reduced tillage combined with low organic matter addition compared to the lower layer of the same treatment ($p = .009$). No effect of soil management was found for relative abundances of omnivorous and predatory nematodes, but the relative abundance of omnivorous nematodes was 68% higher in the upper than in the lower layer across tillage and organic matter treatments.

The food web indices, MI, SI and CI were significantly higher in plots where reduced tillage was applied (MI = 1.8, SI = 37.0, CI = 8.0) than in conventional tillage plots (MI = 1.6, SI = 29.8, CI = 5.0), while the EI was significantly higher under conventional tillage (EI = 81.1) than under reduced tillage (EI = 75.1; Table 3, Figure 4). We found significantly higher values of MI in the upper (MI = 1.7) than in the lower layer (MI = 1.6), and significantly higher values of EI in the lower (EI = 79.4) than in the upper layer (EI = 76.8; Table 3). Accordingly, we found a 13% higher proportion of c-p 1 (colonizers) and a 32% lower proportion of c-p 4 (persisters) in the lower layer than in the upper layer (Table S8), but in terms of absolute abundance the c-p 1 nematodes were 29% higher in the upper layer (2,286 nematodes 100 g field moist soil⁻¹) compared to the lower one (1,812 nematodes 100 g field moist soil⁻¹; Table S9).

TABLE 1 Results of the mixed linear models testing the effect of soil management on total nematode qPCR counts, OTU richness, diversity, and evenness, and genus richness, diversity and evenness

Group A	qPCR counts	OTU richness (total OTU number)	OTU diversity (exp ^H)	OTU evenness (exp ^H /OTU number)	Genus richness (total genus number)	Genus diversity (exp ^H)	Genus evenness (exp ^H /genus number)
0–10 cm							
CT–LOW	6,373 (4,428–8,671)	112 (101–122)	15.8 (12.6–19.4)	0.14 (0.11–0.17)	41 (38–45)	8.3 (6.9–9.6)	0.20 (0.17–0.24)
RT–LOW	6,640 (4,596–9,059)	118 (107–130)	19.2 (15.6–23.3)	0.16 (0.13–0.20)	42 (38–45)	9.8 (8.4–11.2)	0.24 (0.20–0.28)
CT–HIGH	6,725 (4,574–9,289)	117 (105–129)	13.7 (10.5–17.2)	0.11 (0.09–0.15)	42 (38–45)	6.3 (4.9–7.8)	0.15 (0.11–0.19)
RT–HIGH	6,999 (4,870–9,512)	124 (112–136)	16.8 (13.4–20.6)	0.14 (0.11–0.17)	42 (38–46)	7.9 (6.5–9.3)	0.19 (0.15–0.23)
10–20 cm							
CT–LOW	4,832 (3,162–6,856)	100 (90–110)	13.4 (10.5–16.7)	0.13 (0.11–0.16)	39 (36–43)	7.5 (6.2–8.8)	0.19 (0.16–0.23)
RT–LOW	5,065 (3,304–7,201)	106 (96–118)	16.6 (13.2–20.3)	0.16 (0.12–0.19)	40 (36–43)	9.1 (7.7–10.5)	0.23 (0.19–0.27)
CT–HIGH	5,139 (3,285–7,407)	105 (94–116)	11.4 (8.5–14.7)	0.11 (0.08–0.14)	40 (36–43)	5.6 (4.1–7.1)	0.15 (0.11–0.19)
RT–HIGH	5,379 (3,537–7,606)	112 (101–123)	14.3 (11.2–17.9)	0.13 (0.10–0.16)	40 (36–44)	7.2 (5.8–8.6)	0.18 (0.14–0.22)
Tillage (T)							
F	0.22	6.56	10.26	6.45	0.39	7.31	5.89
p	.64	.02	.004	.02	.54	.01	.02
OM							
F	0.18	2.70	3.78	8.69	0.21	12.49	11.05
p	.67	.11	.05	.007	.65	.002	.003
Layer (L)							
F	8.40	38.73	7.75	0.75	10.60	2.47	0.53
p	.005	<.0001	.007	.40	.002	.12	.47
Group B							
LOW–CT	4,353 (617–11,473)	110 (83–140)	17.6 (10.7–26.3)	0.16 (0.10–0.25)	41 (36–46)	9.0 (5.0–14.0)	0.22 (0.12–0.35)
HIGH–CT	5,898 (1,393–13,521)	117 (91–147)	16.6 (10.9–23.4)	0.14 (0.08–0.21)	43 (38–47)	8.9 (5.4–13.2)	0.21 (0.12–0.32)
OM							
F	3.65	2.29	0.22	1.29	2.47	0.01	0.17
p	.08	.16	.65	.28	.14	.94	.69

Note: We tested: for group A (CH1, CH2, NL1, NL2, SL1, HU4 and ES4) the effect of tillage, organic matter addition and layer, and for group B (CH3, PT1 and CH3) only the effect of organic matter addition. For each group, in the upper part of the table the estimated means and 95% confidence intervals (in parentheses) are reported. In the lower part of the table, F statistics and p-values (values $\leq .05$ in bold) for the main factors and their interactions are reported. The interactions are not reported because they were all not significant.

Abbreviations: CT, conventional tillage; exp^H, exponential of the Shannon diversity index; OM, organic matter; RT, reduced tillage.




FIGURE 3 Constrained analysis of principal coordinates (CAP) showing in (a) the effect of management and layer on the nematode beta diversity in group A (CH1, CH2, NL1, NL2, SL1, ES4 and HU4). The CAP model explained in total 8% of the variation in beta diversity related to soil management (tillage, organic matter addition), and the first two axes explained 2.6% and 2.3% of the total variation, respectively. (b) Shows the relationship between the nematode communities (displayed as centroids) and the soil parameters. Only the significant variables at $p < .01$ are shown. The long-term field experiment (LTE) was used as a random effect (conditioned), and the blocking structure plus tillage, organic matter addition and layer were used as fixed effects. The different colours show the soil management and the different shapes show the different layers [Colour figure can be viewed at wileyonlinelibrary.com]

For the LTEs belonging to group B, the proportion of bacterivorous nematodes was significantly increased with high compared to low organic matter addition, while herbivorous nematodes showed the opposite pattern (Table 2). However, in absolute abundance the herbivorous nematodes did not differ between the two treatments (Table S7). We found no effect of organic matter addition on most food web indices. Only the CI was significantly higher in the low than in the high organic matter treatment (Table 3).

3.5 | Relationships between soil parameters and nematode communities

Partial correlations between total nematode qPCR counts and soil chemical, physical, and biological parameters are reported in Table 4. In group A, qPCR counts were positively correlated with many chemical (TN, TOC, available K, Mg), physical (WSA) and biological (SR, MBC, MBN, qMic, soil suppressiveness) parameters, and with four of the labile carbon fractions (Hy-DOC, POXC, HWEC, and POMC). Negative correlations were found with the soil C to N ratio, BD, tea bag decomposition, and Hy- and DOC SUVA (Table 4).

Correlations between OTU richness and soil parameters were similar to those of nematode qPCR counts and soil parameters, although the correlation coefficients were weaker for all the variables except K (Table 4). In contrast, correlations between OTU diversity or evenness and soil parameters were fewer, and, with the exception of CEC, explained less or the same amount of the variance (Table 4). For group

B we found very few and not very strong significant relationships between soil parameters and nematode communities (Table 4).

TOC, available K, BD, MBC, MBN, SR, HWEC, POXC, and POMC were significantly associated with nematode community composition (Table S10). Of these variables, only the ones with a significance level <0.01 are reported in Figure 4b (BD, available K, MBN, POMC, HWEC, SR). With the exception of BD, these parameters, plus TN and Mg, were positively correlated with CAP1 and negatively correlated with CAP2 (Table S11), being higher in the upper compared to lower layers (Figure 3b). The contrary was true for the BD, which was higher in the lower layer, in particular under reduced tillage (Figure 3b, Table S11). In addition, qMic and DOC SUVA were positively and negatively related, respectively, only with CAP1, CEC and WSA were negatively correlated only with CAP2, and C to N ratio was positively correlated only with CAP2 (Table S11).

3.6 | Indicator OTUs for tillage and organic matter addition

Out of 349 OTUs finally used for analysis, 12 OTUs were significantly associated with specific management combinations in the upper layer, and 10 OTUs were significantly associated with the lower layer (group A only, as no differences in nematode communities were found in group B, Table 5). The indicator OTUs were herbivorous (OTUs assigned as *Pratylenchus*, *Neopsilenchus*, Merlinidae), fungivorous (OTUs assigned as *Aphelenchoides*, *Nothotylenchus*) and bacterivorous (OTUs assigned as *Acrobeloides*, *Panagrolaimus*,

TABLE 2 Results of the mixed linear models testing the effect of soil management on the percentage of nematode trophic groups (bacterivores, fungivores, herbivores, omnivores and predators)

	Bacterivores	Fungivores	Herbivores	Omnivores	Predators
	Relative abundance (%)				
Group A					
0–10 cm					
CT–LOW	52 (35–68)	12 (6–22) ^{b, c}	17 (6–39)	1.3 (0.3–4.6)	0.6 (0.2–2.4)
RT–LOW	53 (35–70)	13 (6–25) ^c	18 (7–41)	2.2 (0.5–8.3)	0.9 (0.2–3.50)
CT–HIGH	65 (46–80)	9 (4–18) ^{a, b, c}	16 (5–38)	1.1 (0.2–4.8)	0.4 (0.1–1.5)
RT–HIGH	56 (38–73)	7 (3–15) ^{a, b, c}	20 (7–44)	1.4 (0.3–5.5)	0.8 (0.2–3.0)
10–20 cm					
CT–LOW	58 (40–73)	10 (5–19) ^{a, b, c}	21 (8–45)	0.7 (0.2–2.9)	0.9 (0.2–3.1)
RT–LOW	40 (25–58)	7 (3–14) ^a	45 (21–72)	0.5 (0.1–2.3)	0.8 (0.2–3.0)
CT–HIGH	67 (49–81)	6 (3–13) ^{a, b}	17 (6–40)	0.3 (0.1–1.5)	0.4 (0.1–1.6)
RT–HIGH	43 (26–61)	8 (4–17) ^{a, b, c}	43 (19–70)	0.4 (0.1–1.6)	0.6 (0.1–2.1)
Tillage					
<i>F</i>	12.2	0.97	20.15	0.09	1.52
<i>p</i>	.002	.33	.0001	.76	.23
OM					
<i>F</i>	3.7	5.98	0.20	1.27	3.45
<i>p</i>	.067	.02	.65	.27	.07
Layer					
<i>F</i>	3.64	10.27	27.43	25.35	0.02
<i>p</i>	.06	.002	<.0001	<.0001	.88
T × OM					
<i>F</i>	2.14	0.83	0.52	0.01	1.01
<i>p</i>	.15	.37	.47	.92	.32
T × L					
<i>F</i>	13.55	0.17	14.49	1.82	1.60
<i>p</i>	.0005	.68	.0004	.18	.21
OM × L					
<i>F</i>	0.13	3.92	0.39	0.43	0.25
<i>p</i>	.71	.05	.53	.51	.62
T × OM × L					
<i>F</i>	0.06	7.22	0.006	0.90	0.005
<i>p</i>	.79	.009	.94	.35	.94
Group B					
LOW–CT	47 (11–86)	9 (4–21)	29 (6–72)	0.7 (0.01–33)	1.7 (0.06–32)
HIGH–CT	62 (20–92)	11 (4–26)	18 (6–72)	0.9 (0.01–0.36)	1.8 (0.07–32)
OM					
<i>F</i>	9.82	1.55	6.65	0.33	0.05
<i>p</i>	.009	.24	.02	.58	.82

Note: We assessed for group A (CH1, CH2, NL1, NL2, SL1, HU4 and ES4) the effect of tillage, organic matter addition and layer, and for group B (CH3, PT1 and CH3) the effect of organic matter addition. For each group, in the upper part of the table the estimated means and 95% confidence intervals (in parentheses) are reported. In the lower part of the table, *F* statistics and *p*-values (values ≤ .05 in bold) for the main factors and their interactions are reported. Different superscript letters (a, b, c) following means (to be read per column) show treatments which are significantly different (*p* ≤ .05) according to Tukey post-hoc tests for the three way interactions.

Abbreviations: CT, conventional tillage; L, layer; OM, organic matter; RT, reduced tillage; T, tillage.

TABLE 3 Results of the mixed linear model testing the effect of soil management on the maturity index, enrichment index, structure index and channel index

	Maturity index	Enrichment index	Structure index	Channel index
Group A				
0–10 cm				
CT–LOW	1.64 (1.44–1.85)	79.4 (65.9–92.9)	32.9 (15.6–50.4)	6.5 (2.0–21.3)
RT–LOW	1.84 (1.63–2.04)	73.4 (59.9–86.9)	40.3 (22.8–57.8)	10.3 (3.1–33.8)
CT–HIGH	1.56 (1.35–1.77)	80.1 (66.4–93.7)	29.2 (11.5–47.0)	4.9 (1.5–16.4)
RT–HIGH	1.75 (1.54–1.96)	74.1 (60.5–87.6)	36.5 (18.9–54.1)	7.8 (2.4–25.7)
10–20 cm				
CT–LOW	1.56 (1.36–1.76)	82.1 (68.6–95.5)	30.2 (12.8–47.6)	5.2 (1.6–16.9)
RT–LOW	1.75 (1.54–1.96)	76.1 (62.5–89.6)	37.5 (19.9–55.0)	8.1 (2.5–26.7)
CT–HIGH	1.48 (1.26–1.69)	82.7 (69.1–96.3)	26.4 (8.7–44.2)	3.9 (1.2–12.9)
RT–HIGH	1.67 (1.46–1.88)	76.7 (63.1–90.2)	33.7 (16.1–51.3)	6.1 (1.9–20.3)
Tillage				
<i>F</i>	13.13	12.56	8.16	8.28
<i>p</i>	.001	.001	.008	.008
OM				
<i>F</i>	2.40	0.12	1.64	2.65
<i>p</i>	0.13	0.72	0.21	0.11
Layer				
<i>F</i>	4.92	4.45	1.56	3.58
<i>p</i>	.03	.04	.22	.06
Group B				
LOW–CT	2.1 (1.1–3.1)	67.2 (42.9–91.5)	49.0 (–0.24.8–122.9)	20.8 (–1.2–42.9)
HIGH–CT	1.9 (1.0–2.9)	74.4 (51.6–97.2)	47.5 (–25.9–121.0)	11.8 (–9.3–33.0)
OM				
<i>F</i>	1.85	3.10	0.17	8.8
<i>p</i>	.20	.10	.69	.01

Note: We assessed for group A (CH1, CH2, NL1, NL2, SL1, HU4 and ES4) the effect of tillage, organic matter addition and layer, and for group B (CH3, PT1 and CH3) the effect of organic matter addition. In the table *F* statistics and *p*-values (significance at $p \leq .05$ in bold) for the main factors are reported. The interactions are not reported because they were all not significant.

Abbreviations: CT, conventional tillage; OM, organic matter; RT, reduced tillage.

Rhabditis). Indicator OTUs belonged mainly to c-p groups 1 and 2 and were all present in relative abundance <0.1%, apart from OTU_2 (OTU assigned as *Rhabditis*) which was an indicator OTU for conventional tillage in the lower layer. This OTU comprised more than 20% of the relative abundance of all nematode reads.

4 | DISCUSSION

4.1 | Largest proportion of variation in nematode communities is explained by site

Measured abiotic and biotic (MBC) differences between the LTEs explained most of the variation in nematode communities, in line with results from Neher, Peck, Rawlings, and Campbell (1995) and

Thomson et al. (2015). This result is plausible, since the LTEs were selected to maximize intersite variation and to test if, in spite of large differences in sites across pedoclimatic conditions, effects of agricultural management were yet significant. Indeed, nematode communities were significantly related to all other measured soil parameters when LTE was not used as a random factor.

4.2 | Reduced tillage increases nematode alpha diversity and alters beta diversity compared to conventional tillage

In accordance with our first hypothesis, nematode OTU richness and, to a larger extent, OTU (and genus) diversity and evenness were increased in reduced compared to conventional tillage across




FIGURE 4 Enrichment (y axis)—structure (x axis) diagram for the long-term field experiments (LTEs) of group A (CH1, CH2, NL1, NL2, SL1, ES4, HU4). The points and the triangles represent the estimated means from the linear effect mixed models for the respective combination of factors (tillage, organic matter addition) for the first layer and the second layer, respectively. The bars represent the estimated standard errors for the group averages. In the corner of each of the four quadrants we report information relative to structure of the food web and nutrient enrichment, respectively, according to Ferris et al. (2001) [Colour figure can be viewed at wileyonlinelibrary.com]

the LTEs of group A, i.e., in LTEs where the 0–10 and 10–20 cm layers were sampled (LTEs: CH1, CH2, SL1, NL1, NL2, ES4, HU4). Previous studies reported positive effects of reduced tillage on nematode abundance, richness, and diversity (Fu, Coleman, Hendrix, & Crossley, 2000; Okada & Harada, 2007; Zhang et al., 2015). Reduced soil disturbance (here very shallow or noninversion cultivation in the 0–10 cm layer) can exert a positive effect on nematodes through the increase of total organic carbon, soil aggregation and microbial biomass, and a lower physical pressure (Kladivko, 2001). The lower nematode qPCR counts, richness and diversity in the lower soil layer under reduced tillage, where disturbance is lower, could be due to decreased resources present in this layer. Under reduced tillage, soil parameters related to soil organic matter and nutrients have lower values below the plough layer (Franzluebbers, 2002), which can be explained by the retention of crop residues on the soil surface, and the lack of mechanical mixing of soil layers.

In group A, reduced tillage led to a shift in nematode community structures, in agreement with previous studies (Brmež, Ivezić, & Raspudić, 2006; Griffiths, Daniell, Donn, & Neilson, 2012; Okada & Harada, 2007). In this group of LTEs, nematode beta diversity was affected by the organic matter additions, and OTU diversity was lower in the plots with high organic matter additions, which might suggest positive effects of the organic matter added on a few opportunistic nematodes. However, in disagreement with our second hypothesis, we did not find an effect of organic matter additions on nematodes qPCR counts, and alpha and beta diversity in group B, i.e., in LTEs where the 0–20 cm layer was sampled as a whole (LTEs: CH3, PT1, HU1). Also in the literature contradictory results were found, reporting negative (Wang, McSorley, & Gallaher, 2004), neutral (Ito et al., 2015; Li et al., 2018; Quist et al., 2016) and positive effects of organic matter on nematode

numbers (Nahar et al., 2006; Sánchez-Moreno et al., 2009; Ugarte, Zaborski, & Wander, 2013), richness (Sánchez-Moreno et al., 2009) and alpha diversity (van Diepeningen, de Vos, Korthals, & van Bruggen, 2006; Okada & Harada, 2007) in systems where organic matter was added.

Organic matter is a food source for microorganisms which in turn are a food source for bacterivorous, fungivorous and omnivorous nematodes; therefore, organic matter, similarly to reduced tillage, can change soil properties favourable to nematodes (food availability, but also water retention and soil aggregation; Bongers & Ferris, 1999). In the LTEs of group B, we found higher concentrations of total (TOC) and labile (POXC) organic matter ($p = .03$ and $p < .0001$, respectively) in the high compared to the low organic matter input treatments, but we did not find differences in microbial biomass, cation exchange capacity and water stable aggregates ($p = .06$, $p = .12$ and $p = .51$, respectively). Our contradicting results on the effect of organic matter additions on nematodes could be related to the different types of organic matter used in our LTEs (e.g., compost, biochar, farmyard manure, etc). The composition and the amount of organic matter applied to the soil is an important factor for its effect on nematodes (Ito et al., 2015; Li et al., 2018; Liu et al., 2016). Also, it is possible that the conventional tillage applied to the LTEs of group B neutralized the effect of organic matter additions (Briar, Grewal, Somasekhar, Stinner, & Miller, 2007). This weak effect of organic matter addition supports previous studies that suggested that tillage has a stronger effect on nematode communities than organic matter addition or other agricultural practices such as organic versus conventional management, irrigation, and cover crops (Ito et al., 2015; Neher, 1999; du Preez, Daneel, Wepener, & Fourie, 2018; Zhong, Zeng, & Jin, 2017).

TABLE 4 Partial correlation coefficients between total nematode qPCR counts, OU richness, diversity, and evenness and chemical, physical and biological indicators for the samples belonging to group A ($n = 132$) and group B ($n = 35$)

	Group A				Group B			
	qPCR counts	OTU richness (total OTUs number)	OTU diversity (\exp^H)	OTU evenness (\exp^H /OTU number)	qPCR counts	OTU richness (total OTUs number)	OTU diversity (\exp^H)	OTU evenness (\exp^H /OTU number)
Chemical parameters								
TOC	0.31**	0.36***	0.14	-0.0002	0.12	0.003	0.15	0.14
pH	-0.02	0.06	-0.02	-0.04	0.37	0.06	-0.05	-0.07
TN	0.34***	0.34***	0.18*	0.04	0.02	-0.22	0.05	0.15
C/N	-0.35**	-0.28*	-0.25*	-0.15	0.12	0.29	0.17	0.06
CEC	0.10	0.14	0.33***	0.30**	-0.14	-0.39*	0.19	0.37
Ca	-0.02	0.04	-0.09	-0.10	0.03	0.07	0.29	0.28
Mg	0.13	0.18*	0.24*	0.18*	-0.15	-0.27	0.04	0.15
K	0.21*	0.39***	0.25*	0.14	0.25	0.17	0.11	0.03
Na	-0.19*	-0.20*	-0.10	-0.04	-0.05	0.10	0.11	0.06
P	0.14	0.25*	0.08	-0.01	-0.10	0.21	0.15	0.07
Physical parameters								
WSA	0.24*	0.30***	0.17*	0.06	0.10	-0.14	-0.24	-0.17
WHC	0.06	0.04	0.03	0.01	0.007	-0.03	0.10	0.11
BD	-0.38***	-0.38***	-0.17	-0.03	-0.20	-0.06	0.15	0.17
Sand	0.04	-0.009	-0.08	-0.08	-0.11	0.48*	0.28	0.07
Silt	0.07	0.10	-0.06	-0.11	0.27	0.23	0.02	-0.08
Clay	-0.05	-0.20*	-0.04	0.03	-0.51*	-0.37	0.19	0.36
Biological parameters								
MBC	0.43***	0.41***	0.16	0.0007	-0.08	-0.23	-0.06	0.04
MBN	0.44***	0.21*	0.05	-0.04	-0.24	0.13	0.19	0.14
SR	0.45***	0.33***	0.24*	0.09	0.10	-0.05	-0.15	-0.12
qMic	0.22*	0.22*	0.09	0.009	0.009	-0.22	-0.15	-0.05
qCO ₂	-0.02	-0.02	0.13	0.19*	0.20*	0.21	-0.02	-0.11
Earthworm number	-0.10	-0.09	-0.17	-0.02	0.08	-0.16	-0.10	-0.03
Earthworm biomass	0.05	-0.04	-0.12	-0.05	0.09	-0.24	-0.16	-0.06
Tea bag decomposition	-0.49*	-0.31*	-0.35*	-0.27*	0.002	0.22	-0.12	-0.20
Soil suppressiveness	0.37*	0.20	0.13	0.07	-0.16	0.09	0.0008	-0.04
Labile carbon fractions								
Hy SUVA	-0.20*	0.06	0.07	0.05	-0.10	0.17	-0.11	-0.19
DOC SUVA	-0.26*	-0.06	0.07	0.10	0.009	-0.02	0.03	0.04
Hy-DOC	0.27*	0.14	-0.06	-0.13	0.08	-0.02	0.05	0.06
DOC	0.06	0.09	0.08	0.03	0.10	-0.13	-0.09	-0.03
HWEC	0.48***	0.35***	0.19*	0.05	0.06	-0.16	0.08	0.14
POXC	0.46***	0.36***	0.18*	0.04	0.17	0.003	0.10	0.09
POMC	0.49***	0.46***	0.12	-0.07	0.24	0.16	0.08	0.01

Abbreviations: BD, bulk density; C/N, carbon to nitrogen ratio; CEC, cation exchange capacity; DOC SUVA, specific ultraviolet absorbance of dissolved organic carbon; DOC, dissolved organic carbon; HWEC, hot water extractable carbon; Hy SUVA, specific ultraviolet absorbance of hydrophilic carbon; Hy, hydrophilic carbon; MBC, microbial biomass carbon; MBN, microbial biomass nitrogen; POMC, particulate organic matter carbon; POXC, permanganate oxidizable carbon; qCO₂, metabolic quotient; qMic, microbial quotient; TOC, total organic carbon; TON, total nitrogen; WHC, water holding capacity; WSA, water stable aggregates.

* $p \leq .05$.

** $p \leq .001$.

*** $p \leq .0001$.

TABLE 5 Indicator species for the combination of tillage, organic matter addition and layer for group A (CH1, CH2, NL1, NL2, SL1, ES4 and HU4)

OTU	Genus	Family	Feeding group	c-p	Correlation	p-Value	Relative abundance (%)				
							CT-Low	CT-High	RT-Low	RT-High	
							N = 20	N = 13	N = 17	N = 16	
Layer 0–10 cm											
CT–LOW	OTU_329	Aphelenchoiodes	Aphelenchoiidae	F	2	0.34	0.007	0.68 (12)	0.06 (6)	0.01 (6)	0.142 (9)
CT–HIGH	OTU_436	NA	Tylenchidae	H (1e)	NA	0.38	0.005	0.0001 (2)	0.02 (4)	0.0003 (1)	0 (0)
	OTU_75	Aphelenchoiodes	Aphelenchoiidae	F	2	0.26	0.007	0.35 (15)	0.68 (10)	0.15 (15)	0.035 (9)
RT–LOW	OTU_310	Dicelis	Drilonematidae	B	2	0.39	0.006	0.02 (8)	0.004 (9)	0.02 (14)	0.01 (9)
	OTU_44	Oscheius	Rhabditidae	B	1	0.39	0.0001	0.07 (14)	0.21 (13)	0.71 (15)	0.25 (14)
	OTU_84	Aphelenchoiodes	Aphelenchoiidae	F	2	0.37	0.004	0.07 (9)	0.04 (10)	0.28 (11)	0.05 (6)
RT–HIGH	OTU_120	Pratylenchus	Pratylenchidae	H (1b)	3	0.39	0.003	0.002 (2)	0.0001 (1)	0.0001 (1)	0.20 (6)
	OTU_48	NA	NA	NA	NA	0.33	0.0007	0.25 (11)	0.0005 (3)	0.003 (7)	1.41 (12)
	OTU_79	Acroboloides	Cephalobidae	B	2	0.35	0.005	0.113 (9)	0.047 (3)	0.01 (6)	1.29 (7)
CT–HIGH, RT–LOW, RT–HIGH	OTU_256	NA	NA	NA	NA	0.32	0.004	0.002 (2)	0.01 (5)	0.02 (10)	0.03 (10)
	OTU_82	Aphelenchoiodes	Aphelenchoiidae	F	2	0.43	0.0008	0.06 (16)	0.16 (12)	0.25 (17)	0.23 (16)
	OTU_728	Oscheius	Rhabditidae	B	1	0.33	0.002	0.02 (11)	0.12 (9)	0.21 (14)	0.15 (9)
Layer 10–20 cm											
CT–LOW	OTU_329	Aphelenchoiodes	Aphelenchoiidae	F	2	0.31	0.005	0.71 (11)	0.02 (7)	0.03 (8)	0.06 (7)
CT–HIGH	OTU_75	Aphelenchoiodes	Aphelenchoiidae	F	2	0.37	0.0007	0.38 (12)	1.06 (10)	0.04 (8)	0.02 (12)
RT–HIGH	OTU_218	Nothotylenchus	Anguimidae	F	2	0.24	0.006	0.01 (5)	0.004 (4)	0.01 (3)	0.08 (6)
	OTU_257	Panagrolaimus	Panagrolaimidae	B	1	0.37	0.009	0.00 (2)	0.0002 (1)	0.0003 (2)	0.01 (6)
	OTU_30	Pratylenchus	Pratylenchidae	H (1b)	3	0.38	0.003	0.04 (12)	0.003 (9)	0.007 (11)	4.26 (15)
High	OTU_712	Neopsilenchus	Tylenchidae	H (1e)	2	0.51	0.005	0.06 (13)	0.43 (12)	0.07 (12)	0.82 (13)
RT	OTU_15	NA	Merlinidae	H (1d)	NA	0.25	0.008	0.70 (20)	0.07 (13)	2.91 (17)	1.04 (16)
	OTU_70	NA	Tylenchidae	H (1e)	2	0.28	0.001	0.14 (11)	0.04 (7)	0.47 (14)	0.48 (10)
CT	OTU_2	Rhabditis	Rhabditidae	B	1	0.40	0.002	24.82 (20)	30.31 (13)	14.14 (17)	7.85 (16)
	OTU_39	NA	NA	NA	NA	0.32	0.008	0.88 (17)	2.73 (10)	0.02 (13)	0.11 (11)

Note: In the table we report the OTU number, taxonomic information at the level of nematode family and genus, feeding group, c-p colonizer-persister class, the correlation coefficient and the *p*-value from the analysis, and the relative abundances of the taxa for the combinations of soil management (tillage and organic matter addition). Above the columns with the OTU relative abundance the number of samples belonging to each group is indicated, and in parentheses the number of samples in which that specific OTU was found is given. The analysis has been done for the two layers separately, and only OTUs that had a level of significance $p \leq .01$ are reported.

Feeding group: B, bacterivorous nematode; F, fungivorous nematode; H, herbivorous nematode; 1b, herbivorous migratory endoparasitic nematodes; 1d, herbivorous ectoparasitic nematodes; 1e, herbivorous epidermal and root hair feeders; NA, not assigned to taxon. Abbreviations: CT, conventional tillage; RT, reduced tillage.

4.3 | Reduced tillage increases stability and structure of the nematode community compared to conventional tillage

Agricultural management did not have strong effects on the relative abundance of the trophic groups, but it affected the food web indices, indicating effects on rates rather than on structural changes in the food web. This observation supports the suggestion by Neher (1999) that food web indices are less variable and more likely to detect effects of management practices on soil processes than measures based on individual trophic groups.

In accordance with our first hypothesis and in line with previous reports (Habig & Swanepoel, 2015; Zhang et al., 2015; Zhong et al., 2017), reduced tillage resulted in a less disturbed environment than conventional tillage, increasing the stability and the number of food web interactions of the nematode communities (higher MI and SI) in the LTEs of group A. Despite the decreasing level of disturbance in the lower soil layer of reduced tillage, a lower MI and reduced proportions of omnivorous and stress-tolerant c-p 4 nematodes compared to the upper layer seems to indicate a more stressed environment where opportunistic nematodes can prevail. In our study, reduced tillage increased the channel index (CI), i.e., among the opportunistic microbivorous nematodes there was an increase in the proportion of fungal feeders, confirming previous findings (Minoshima et al., 2007; Okada & Harada, 2007; Sánchez-Moreno, Minoshima, Ferris, & Jackson, 2006). Reduced tillage is known to favour the fungal decomposition pathway (Six, Frey, Thiet, & Batten, 2006), due to less or no disruption of the hyphal network (Minoshima et al., 2007). Since lower values of CI are associated with faster rates of decomposition and nutrient turnover, our results suggest that changes in nematode communities under reduced tillage may contribute to the increased capability of the system to retain nutrients and store carbon (Griffiths et al., 2012). The higher relative and absolute abundance of herbivorous nematodes in reduced tillage compared to conventional tillage is in line with previous studies (Brmež et al., 2006; Freckman & Ettema, 1993; Fu et al., 2000; Treonis et al., 2010, 2018), and can be explained by a higher incidence of roots in the field, stimulating this nematode group (Minton, 1986; You et al., 2017). Our results indicate a possible trade-off in reduced tillage systems in terms of soil processes, and that in these types of systems care must be taken regarding the assessment and control of herbivorous nematodes. However, the higher alpha diversity, MI and SI found in reduced tillage could indicate that the activity of herbivorous populations might be controlled by a more stable and structured food web.

In agreement with our second hypothesis, high organic matter addition plots resulted in higher percentages of bacterivorous nematodes than low organic matter addition plots, and they showed a statistically lower CI and a tendency towards lower SI, MI, and higher EI. High EI (Berkelmans, Ferris, Tenuta, & van Bruggen, 2003; Forge et al., 2005; Sánchez-Moreno et al., 2009), low MI (Forge et al., 2005; Neher & Olson, 1999; Wang, McSorley, Marshall, & Gallaher, 2006) and low SI (Pan et al., 2015; Villenave

et al., 2010) have been previously reported in systems with organic matter addition. Such changes in MI and CI can be explained by an increase in opportunistic bacterivores (Ferris & Bongers, 2006), and a stimulation of the bacterivore decomposition channel (Pan et al., 2010; Wang et al., 2004). Altogether, these results indicate higher nutrient cycling, N mineralization and fertility in soils with high organic matter additions (Ferris & Matute, 2003). By contrast, the addition of organic matter decreased the proportion of herbivorous nematodes, but this did not coincide with an absolute decrease as this relative decrease resulted from the absolute increase of bacterivorous nematodes.

4.4 | Nematode communities are mainly related to soil organic carbon and biological parameters

Total and labile organic carbon and microbial parameters were most strongly and positively related to nematode qPCR counts and richness, partly confirming our third hypothesis. Abundance (Sánchez-Moreno et al., 2006), richness (van Diepeningen et al., 2006), but also diversity (Zhong et al., 2017) of soil nematodes have previously been positively linked with the levels of total and labile organic carbon fractions. Higher total and labile carbon are linked to higher microbial biomass, soil respiration, water retention, soil structure and lower bulk density (Bongiorno, Bünemann, et al., 2019). Increased levels in these soil parameters can optimize the habitat conditions for nematodes, and facilitate their movement through the soil pore water (Nielsen et al., 2014).

Some of the properties that correlated most with nematode qPCR counts and richness (total organic and labile carbon, available K, bulk density, microbial biomass and activity) proved important in explaining differences between nematode communities caused by reduced versus conventional tillage. This suggests that reduced tillage affects nematode communities through its positive effects on these soil properties, either directly through absence of soil inversion, i.e. lower soil disturbance, or indirectly through retention of crop residues at the soil surface, which can increase water retention and infiltration, soil organic carbon, and organism biomass and activity (Mloza-Banda, Makwiza, & Mloza-Banda, 2016; Ranaivoson et al., 2017).

4.5 | Only *r* selected taxa were found to be indicator OTUs for tillage and organic matter addition

Indicator OTU analysis based on group A revealed OTUs that were significantly associated with tillage and organic matter management. Most of the indicator OTUs had a very low relative abundance. These taxa belonged mainly to the c-p 2 group, and to bacterivorous, fungivorous and herbivorous nematode trophic groups. Therefore, contrary to our fourth hypothesis none of the predatory and omnivorous nematodes, or nematodes belonging to c-p groups 4 and 5 were detected as indicator taxa. This can be due to the fact that in these intensively managed European arable systems, relative and absolute abundances of highly sensitive nematode taxa were underrepresented and too variable (i.e., not present in all samples).

4.6 | Advantages and limitations of studying nematode communities with amplicon sequencing

Our molecular analyses revealed that, despite the big influence of the pedoclimatic characteristics, agricultural soil management resulted in changes in nematode communities and nematode food web structure in line with previous findings from microscopic analysis and general knowledge of agricultural systems. In addition, nematode molecular analyses provided advantages in terms of costs and number of samples analyzed at the same time, and did not require expert skills for morphological characterisation.

A limitation of current amplicon sequences approaches is that previous studies found that the relative read abundance obtained do not perfectly match absolute abundance data determined microscopically. Possibly, the number of ribosomal DNA copies differ depending on the taxon, the organism's body size, the developmental stage, and PCR primer bias (Darby et al., 2013; Geisen et al., 2018). This has to be considered and standardized in future efforts to allow direct comparisons between morphological and molecular approaches in determining nematode communities.

In our study, a relatively large group of OTUs could not be classified at all. This underlines the problems in reliably assigning OTUs to their correct taxonomic group. Such taxa could belong to not yet studied nematode species, but most likely could indicate lack of information in the data bases. In addition, our methodology used to assign taxonomy, using only forward reads, could have had negative consequences for annotation (resolution power) and error correction which can be applied during read merging.

All in all, future studies should work towards an optimization of molecular methods for assessing relative and total nematode abundance, nematode taxonomy and the definition of standardized protocols and the amelioration of data bases in order to guarantee a more confident application of nematode communities studied with molecular methods in soil quality assessments.

In conclusion, molecular nematode community analyses effectively differentiate soil management across 10 different European long-term field experiments. In particular, reduced tillage had a stronger effect on nematode communities than organic matter addition, increasing nematode taxon richness, diversity and evenness. Reduced tillage also affected the nematode food web indices, stimulating more mature and fungal-based nematode communities, indicating a more stable food web with higher nutrient retention capability, but also increasing the number of herbivorous nematodes. These results are in line with previous findings based on microscopic analysis and general knowledge on nematode community dynamics in agricultural systems.

The relationships found between soil nematode communities and total and labile organic carbon, total nitrogen, available K, and microbial biomass and activity, underline the relationship between nematode communities and biological soil quality achieved by reduced tillage, and indicate that nematode communities are equally sensitive indicators of soil quality as these parameters.

Our findings indicate that molecular methods are promising in the assessment of biological soil quality based on nematode community

structure and indices, especially if future research will work toward an optimization and standardization of the methods.

ACKNOWLEDGEMENTS

This work was supported by the EU Horizon 2020 project Interactive Soil Quality Assessment in Europe and China for agricultural productivity and environmental resilience (iSQAPER) [grant number 635750] for GB, NB, EKB, LB, PM, and RGMG (mediated through the Swiss State Secretariat for Education, Research and Innovation in the case of NB, EKB, PM and, partly, GB). SG was supported by a VENI grant from the Netherlands Organisation for Scientific Research (016.Veni.181.078). We acknowledge the University of Ljubljana (SL), Trier University (DE), University Miguel Hernandez (ES) and the long-term field experiment owners for the data provided. We thank Tamas Salanki for helping in the nematode extractions and counting.

AUTHOR CONTRIBUTIONS

G.B., R.G.M.G., E.K.B., L.B., and P.M. agreed on the study design with options given by the iSQAPER consortium. C.W.Q., and S.G. contributed analytical tools and primers. G.B., and C.W.Q. collected the data. J.-C.W. performed the bioinformatic analysis. G.B., N.B., and R.G.M.G. analyzed and interpreted the data. G.B. wrote the manuscript and N.B., C.W.Q., S.G., J.-C.W., R.G.M.G., E.K.B., L.B., and P.M. reviewed it.

ORCID

Giulia Bongiorno  <https://orcid.org/0000-0001-9148-1206>

Stefan Geisen  <https://orcid.org/0000-0003-0734-727X>

Casper W. Quist  <https://orcid.org/0000-0002-9706-6684>

DATA AVAILABILITY STATEMENT

Sequences reads have been submitted to the European Nucleotide Archive (ENA) and are available under accession number ERP114920. The script used for data analyses is provided as part of the Appendix S2. The data on soil chemical, physical and biological indicators others than nematodes that support part of the findings of this study are available from iSQAPER work package 3.3. Restrictions apply to the availability of these data, which were used under license for this study. Data are available from the authors with the permission of iSQAPER work package 3.3 consortium, and they will be submitted in a public database after their publication in a separate paper. All the other data have been published in the DRYAD database (<https://doi.org/10.5061/dryad.q21d0b9>; Bongiorno et al., 2016a, 2016b).

REFERENCES

- Ahmed, M., Sapp, M., Prior, T., Karssen, G., & Back, M. A. (2016). Technological advancements and their importance for nematode identification. *Soil*, 2(2), 257–270. <https://doi.org/10.5194/soil-2-257-2016>

- Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, 26(1), 32–46. <https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x>
- Anderson, M. J., & Willis, T. J. (2003). Canonical analysis of principal coordinates: A useful method of constrained ordination for ecology. *Ecology*, 84(2), 511–525. [https://doi.org/10.1890/0012-9658\(2003\)084\[0511:CAOPCA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[0511:CAOPCA]2.0.CO;2)
- Berkelmans, R., Ferris, H., Tenuta, M., & van Bruggen, A. H. C. (2003). Effects of long-term crop management on nematode trophic levels other than plant feeders disappear after 1 year of disruptive soil management. *Applied Soil Ecology*, 23(3), 223–235. [https://doi.org/10.1016/s0929-1393\(03\)00047-7](https://doi.org/10.1016/s0929-1393(03)00047-7)
- Bodenhausen, N., Horton, M. W., & Bergelson, J. (2013). Bacterial communities associated with the leaves and the roots of *Arabidopsis thaliana*. *PLoS ONE*, 8(2), e56329. <https://doi.org/10.1371/journal.pone.0056329>
- Bongers, T. (1990). The maturity index: An ecological measure of environmental disturbance based on nematode species composition. *Oecologia*, 83, 14–19. <https://doi.org/10.1007/bf00324627>
- Bongers, T., & Ferris, H. (1999). Nematode community structure as a bioindicator in environmental monitoring. *Trends in Ecology and Evolution*, 14(6), 224–228. [https://doi.org/10.1016/S0169-5347\(98\)01583-3](https://doi.org/10.1016/S0169-5347(98)01583-3)
- Bongiorno, G., Bodenhausen, N., Bunemann, E. K., Brussaard, L., Geisen, S., & Mäder, P., ... de Goede, R. G. M. (2016a). Nematodes sequence reads. European Nucleotide Archive (ENA); Accession number ERP114920.
- Bongiorno, G., Bodenhausen, N., Bunemann, E. K., Brussaard, L., Geisen, S., Mäder, P., ... de Goede, R. G. M. (2016b). Data from: Reduced tillage, but not organic matter input, increased nematode diversity and food web stability in European long-term field experiment. Dryad Digital Repository, <https://doi.org/10.5061/dryad.q21d0b9>
- Bongiorno, G., Bünemann, E. K., Oguejiofor, C. U., Meier, J., Gort, G., Comans, R., ... de Goede, R. (2019). Sensitivity of labile carbon fractions to tillage and organic matter management and their potential as comprehensive soil quality indicators across pedoclimatic conditions in Europe. *Ecological Indicators*, 99, 38–50. <https://doi.org/10.1016/j.ecolind.2018.12.008>
- Bongiorno, G., Postma, J., Bünemann, E. K., Brussaard, L., de Goede, R. G. M., Mäder, P., ... Thuerig, B. (2019). Soil suppressiveness to *Pythium ultimum* in ten European long-term field experiments and its relation with soil parameters. *Soil Biology and Biochemistry*, 133, 174–187. <https://doi.org/10.1016/j.soilbio.2019.03.012>
- Briar, S. S., Grewal, P. S., Somasekhar, N., Stinner, D., & Miller, S. A. (2007). Soil nematode community, organic matter, microbial biomass and nitrogen dynamics in field plots transitioning from conventional to organic management. *Applied Soil Ecology*, 37(3), 256–266. <https://doi.org/10.1016/j.apsoil.2007.08.004>
- Brmež, M., Ivezić, M., & Raspudić, E. (2006). Effect of mechanical disturbances on nematode communities in arable land. *Helminthologia*, 43(2), 117–121. <https://doi.org/10.2478/s11687-006-0022-0>
- Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R. E., De Deyn, G., de Goede, R., ... Brussaard, L. (2018). Soil quality – A critical review. *Soil Biology and Biochemistry*, 120, 105–125. <https://doi.org/10.1016/j.soilbio.2018.01.030>
- Darby, B. J., Todd, T. C., & Herman, M. A. (2013). High-throughput amplicon sequencing of rRNA genes requires a copy number correction to accurately reflect the effects of management practices on soil nematode community structure. *Molecular Ecology*, 22(21), 5456–5471. <https://doi.org/10.1111/mec.12480>
- De Caceres, M. (2016). *Package 'indicspecies' – Relationship between species and groups of sites*.
- De Goede, R. G. M., Bongers, T., & Ettema, C. H. (1993). Graphical presentation and interpretation of nematode community structure: c-p triangles. *Mededelingen Van De Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent*, 58(2b), 743–750.
- du Preez, G. C., Daneel, M. S., Wepener, V., & Fourie, H. (2018). Beneficial nematodes as bioindicators of ecosystem health in irrigated soils. *Applied Soil Ecology*, 132, 155–168. <https://doi.org/10.1016/j.apsoil.2018.08.008>
- Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, 26(19), 2460–2461. <https://doi.org/10.1093/bioinformatics/btq461>
- Edgar, R. (2013). UPARSE: Highly accurate OTU sequences from microbial amplicon reads. *Nature Methods*, 10, 996–998.
- Edgar, R. C. (2016). SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. *bioRxiv*, 074161. <https://doi.org/10.1101/074161>
- Ekschmitt, K., Bakonyi, G., Bongers, M., Bongers, T., Boström, S., Dogan, H., ... Wolters, V. (2001). Nematode community structure as an indicator of soil functioning in European grassland soils. *European Journal of Soil Biology*, 37(4), 263–268. [https://doi.org/10.1016/S1164-5563\(01\)01095-0](https://doi.org/10.1016/S1164-5563(01)01095-0)
- Ferris, H., & Bongers, T. (2006). Nematode indicators of organic enrichment. *Journal of Nematology*, 38(1), 3–12.
- Ferris, H., Bongers, T., & Goude, R. M. (2001). A framework for soil food web diagnostics: Extension of the nematode faunal analysis concept. *Applied Soil Ecology*, 18, 13–29. [https://doi.org/10.1016/s0929-1393\(01\)00152-4](https://doi.org/10.1016/s0929-1393(01)00152-4)
- Ferris, H., Griffiths, B. S., Porazinska, D. L., Powers, T. O., Wang, K.-H., & Tenuta, M. (2012). Reflections on plant and soil nematode ecology: Past, present and future. *Journal of Nematology*, 44(2), 115–126.
- Ferris, H., & Matute, M. M. (2003). Structural and functional succession in the nematode fauna of a soil food web. *Applied Soil Ecology*, 23(2), 93–110.
- Forge, T. A., Bittman, S., & Kowalenko, C. G. (2005). Responses of grassland soil nematodes and protozoa to multi-year and single-year applications of dairy manure slurry and fertilizer. *Soil Biology and Biochemistry*, 37(10), 1751–1762. <https://doi.org/10.1016/j.soilbio.2004.11.013>
- Franzuebbers, A. J. (2002). Soil organic matter stratification ratio as an indicator of soil quality. *Soil and Tillage Research*, 66(2), 95–106. [https://doi.org/10.1016/S0167-1987\(02\)00018-1](https://doi.org/10.1016/S0167-1987(02)00018-1)
- Freckman, D. W., & Ettema, C. H. (1993). Assessing nematode communities in agroecosystems of varying human intervention. *Agriculture, Ecosystems & Environment*, 45(3), 239–261. [https://doi.org/10.1016/0167-8809\(93\)90074-Y](https://doi.org/10.1016/0167-8809(93)90074-Y)
- Fu, S., Coleman, D. C., Hendrix, P. F., & Crossley, D. A. (2000). Responses of trophic groups of soil nematodes to residue application under conventional tillage and no-till regimes. *Soil Biology and Biochemistry*, 32(11), 1731–1741. [https://doi.org/10.1016/S0038-0717\(00\)00091-2](https://doi.org/10.1016/S0038-0717(00)00091-2)
- Geisen, S., Snoek, L. B., ten Hooven, F. C., Duyts, H., Kostenko, O., Bloem, J., ... van der Putten, W. H. (2018). Integrating quantitative morphological and qualitative molecular methods to analyse soil nematode community responses to plant range expansion. *Methods in Ecology and Evolution*, 9, 1366–1378. <https://doi.org/10.1111/2041-210X.12999>
- George, P. B. L., & Lindo, Z. (2015). Congruence of community structure between taxonomic identification and T-RFLP analyses in free-living soil nematodes. *Pedobiologia*, 58(2–3), 113–117. <https://doi.org/10.1016/j.pedobi.2015.04.003>
- Griffiths, B. S., Daniell, T. J., Donn, S., & Neilson, R. (2012). Bioindication potential of using molecular characterisation of the nematode community: Response to soil tillage. *European Journal of Soil Biology*, 49, 92–97. <https://doi.org/10.1016/j.ejsobi.2011.09.002>
- Habig, J., & Swanepoel, C. (2015). Effects of conservation agriculture and fertilization on soil microbial diversity and activity. *Environments*, 2(3), 358–384. <https://doi.org/10.3390/environments2030358>
- Hamilton, H. C., Strickland, M. S., Wicks, K., Bradford, M. A., & Fierer, N. (2009). Surveying soil faunal communities using a direct molecular

- approach. *Soil Biology and Biochemistry*, 41(6), 1311–1314. <https://doi.org/10.1016/j.soilbio.2009.03.021>
- Heip, C. (1974). A new index measuring evenness. *Journal of the Marine Biological Association of the United Kingdom*, 54(3), 555–557. <https://doi.org/10.1017/s0025315400022736>
- Holterman, M., van der Wurff, A., van den Elsen, S., van Megen, H., Bongers, T., Holovachov, O., ... Helder, J. (2006). Phylum-wide analysis of SSU rDNA reveals deep phylogenetic relationships among nematodes and accelerated evolution toward crown clades. *Molecular Biology and Evolution*, 23(9), 1792–1800. <https://doi.org/10.1093/molbev/msl044>
- Ito, T., Araki, M., Komatsuzaki, M., Kaneko, N., & Ohta, H. (2015). Soil nematode community structure affected by tillage systems and cover crop managements in organic soybean production. *Applied Soil Ecology*, 86, 137–147. <https://doi.org/10.1016/j.apsoil.2014.10.003>
- Ivanova, N. V., Dewaard, J. R., & Hebert, P. D. N. (2006). An inexpensive, automation-friendly protocol for recovering high-quality DNA. *Molecular Ecology Notes*, 6(4), 998–1002. <https://doi.org/10.1111/j.1471-8286.2006.01428.x>
- Jost, L. (2010). Independence of alpha and beta diversities. *Ecology*, 91(7), 1969–1974. <https://doi.org/10.1890/09-0368.1>
- Kim, S. (2015). `PPCOR`: An R package for a fast calculation to semi-partial correlation coefficients. *Communications for Statistical Application and Methods*, 22(6), 665–674. <https://doi.org/10.5351/CSAM.2015.22.6.665>
- Kladivko, E. J. (2001). Tillage systems and soil ecology. *Soil and Tillage Research*, 61(1), 61–76. [https://doi.org/10.1016/S0167-1987\(01\)00179-9](https://doi.org/10.1016/S0167-1987(01)00179-9)
- Köppen, W. (1918). Klassifikation der Klimate nach Temperatur. *Niederschlag und Jahresablauf Petermans Geographische Mitteil*, 64, 193–203, 243–248.
- Leff, J. W., Jones, S. E., Prober, S. M., Barberán, A., Borer, E. T., Firn, J. L., ... Fierer, N. (2015). Consistent responses of soil microbial communities to elevated nutrient inputs in grasslands across the globe. *Proceedings of the National Academy of Sciences of the United States of America*, 112(35), 10967–10972. <https://doi.org/10.1073/pnas.1508382112>
- Leroy, B. L. M. M., Bommele, L., Reheul, D., Moens, M., & De Neve, S. (2007). The application of vegetable, fruit and garden waste (VFG) compost in addition to cattle slurry in a silage maize monoculture: Effects on soil fauna and yield. *European Journal of Soil Biology*, 43(2), 91–100. <https://doi.org/10.1016/j.ejsobi.2006.10.005>
- Li, J., Wang, D., Fan, W., He, R., Yao, Y., Sun, L., ... Wu, J. (2018). Comparative effects of different organic materials on nematode community in continuous soybean monoculture soil. *Applied Soil Ecology*, 125, 12–17. <https://doi.org/10.1016/j.apsoil.2017.12.013>
- Liu, T., Chen, X., Hu, F., Ran, W., Shen, Q., Li, H., & Whalen, J. K. (2016). Carbon-rich organic fertilizers to increase soil biodiversity: Evidence from a meta-analysis of nematode communities. *Agriculture, Ecosystems & Environment*, 232, 199–207. <https://doi.org/10.1016/j.agee.2016.07.015>
- Magurran, A. E. (1988). *Ecological diversity and its measurements*. Princeton, NJ: Princeton University Press.
- Mekonen, S., Petros, I., & Hailemariam, M. (2017). The role of nematodes in the processes of soil ecology and their use as bioindicators. *Agriculture and Biology Journal of North America*, 8(4), 132–140.
- Minoshima, H., Jackson, L. E., Cavagnaro, T. R., Sánchez-Moreno, S., Ferris, H., Temple, S. R., ... Mitchell, J. P. (2007). Soil food webs and carbon dynamics in response to conservation tillage in California. *Soil Science Society of America Journal*, 71(3), 952–963. <https://doi.org/10.2136/sssaj2006.0174>
- Minton, N. A. (1986). Impact of conservation tillage on nematode populations. *Journal of Nematology*, 18(2), 135–140.
- Mloza-Banda, H. R., Makwiza, C. N., & Mloza-Banda, M. L. (2016). Soil properties after conversion to conservation agriculture from ridge tillage in Southern Malawi. *Journal of Arid Environments*, 127, 7–16. <https://doi.org/10.1016/j.jaridenv.2015.11.001>
- Moura, G. S., & Franzener, G. (2017). Biodiversity of nematodes biological indicators of soil quality in the agroecosystems. *Arquivos do Instituto Biológico*, 84, 1–8. <https://doi.org/10.1590/1808-1657000142015>
- Nahar, M. S., Grewal, P. S., Miller, S. A., Stinner, D., Stinner, B. R., Kleinhenz, M. D., ... Doohan, D. (2006). Differential effects of raw and composted manure on nematode community, and its indicative value for soil microbial, physical and chemical properties. *Applied Soil Ecology*, 34(2), 140–151. <https://doi.org/10.1016/j.apsoil.2006.03.011>
- Neher, D. A. (1999). Nematode communities in organically and conventionally managed agricultural soils. *Journal of Nematology*, 31(2), 142–154.
- Neher, D. A. (2001). Role of nematodes in soil health and their use as indicators. *Journal of Nematology*, 33(4), 161–168.
- Neher, D. A., & Olson, R. K. (1999). Nematode communities in soils of four farm cropping management systems. *Pedobiologia*, 43, 430–438.
- Neher, D. A., Peck, S. L., Rawlings, J. O., & Campbell, C. L. (1995). Measures of nematode community structure and sources of variability among and within agricultural fields. *Plant and Soil*, 170(1), 167–181. <https://doi.org/10.1007/bf02183065>
- Neher, D. A., Weicht, T. R., & Barbercheck, M. E. (2012). Linking invertebrate communities to decomposition rate and nitrogen availability in pine forest soils. *Applied Soil Ecology*, 54, 14–23. <https://doi.org/10.1016/j.apsoil.2011.12.001>
- Nielsen, U. N., Ayres, E., Wall, D. H., Li, G., Bardgett, R. D., Wu, T., & Garey, J. R. (2014). Global-scale patterns of assemblage structure of soil nematodes in relation to climate and ecosystem properties. *Global Ecology and Biogeography*, 23(9), 968–978. <https://doi.org/10.1111/geb.12177>
- Niles, R. K., & Freckman, D. W. (1998). From the ground up: Nematode ecology in bioassessment and ecosystem health. In *Plant and nematode interactions*. Agronomy monograph (vol. 36, pp. 65–85). Madison, WI: Soil Science Society of America.
- Okada, H., & Harada, H. (2007). Effects of tillage and fertilizer on nematode communities in a Japanese soybean field. *Applied Soil Ecology*, 35(3), 582–598. <https://doi.org/10.1016/j.apsoil.2006.09.008>
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., & Wagner, H. (2018). *VEGAN: Community ecology package*.
- Oostenbrink, M. (1960). Estimating nematode populations by some selected methods. *Nematology*, 6, 85–102.
- Pan, F., Han, X., McLaughlin, N. B., Han, X., Li, C., Zhao, D., ... Xu, Y. (2015). Effect of long-term fertilization on free-living nematode community structure in Mollisols. *Journal of Soil Science and Plant Nutrition*, 15, 129–141. <https://doi.org/10.4067/S0718-95162015005000011>
- Pan, F., McLaughlin, N. B., Yu, Q., Xue, A. G., Xu, Y., Han, X., ... Zhao, D. (2010). Responses of soil nematode community structure to different long-term fertilizer strategies in the soybean phase of a soybean-wheat-corn rotation. *European Journal of Soil Biology*, 46(2), 105–111. <https://doi.org/10.1016/j.ejsobi.2010.01.004>
- Pinheiro, J., Bates, D., DebRoy, S., & Sarkar, D. (2018). *nlme: Linear and nonlinear mixed effect models*. Retrieved from <https://CRAN.R-project.org/package=nlme>
- Porazinska, D. L., Giblin-davis, R. M., Faller, L., Farmerie, W., Kanzaki, N., Morris, K., ... Thomas, W. K. (2009). Evaluating high-throughput sequencing as a method for metagenomic analysis of nematode diversity. *Molecular Ecology Resources*, 9(6), 1439–1450. <https://doi.org/10.1111/j.1755-0998.2009.02611.x>
- Porazinska, D. L., Sung, W., Giblin-Davis, R. M., & Thomas, W. K. (2010). Reproducibility of read numbers in high-throughput sequencing analysis of nematode community composition and structure. *Molecular Ecology Resources*, 10(4), 666–676. <https://doi.org/10.1111/j.1755-0998.2009.02819.x>
- Quist, C. W., Gort, G., Mulder, C., Wilbers, R. H. P., Termorshuizen, A. J., Bakker, J., & Helder, J. (2017). Feeding preference as a main determinant of microscale patchiness among terrestrial nematodes. *Molecular Ecology Resources*, 17(6), 1257–1270. <https://doi.org/10.1111/1755-0998.12672>

- Quist, C. W., Schrama, M., de Haan, J. J., Smant, G., Bakker, J., van der Putten, W. H., & Helder, J. (2016). Organic farming practices result in compositional shifts in nematode communities that exceed crop-related changes. *Applied Soil Ecology*, *98*, 254–260. <https://doi.org/10.1016/j.apsoil.2015.10.022>
- R Development Core Team (2013). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ranaivoson, L., Naudin, K., Ripoche, A., Affholder, F., Rabeharisoa, L., & Corbeels, M. (2017). Agro-ecological functions of crop residues under conservation agriculture. A review. *Agronomy for Sustainable Development*, *37*(4), 26. <https://doi.org/10.1007/s13593-017-0432-z>
- Ritz, K., Black, H. I. J., Campbell, C. D., Harris, J. A., & Wood, C. W. (2009). Selecting biological indicators for monitoring soils: A framework for balancing scientific and technical opinion to assist policy development. *Ecological Indicators*, *9*(6), 1212–1221. <https://doi.org/10.1016/j.ecolind.2009.02.009>
- RStudio Team (2016). *RStudio: Integrated development for R*. Boston, MA: RStudio Team. Retrieved from <http://www.rstudio.com/>
- Sánchez-Moreno, S., Minoshima, H., Ferris, H., & Jackson, L. (2006). Linking soil properties and nematode community composition: Effects of soil management on soil food webs. *Nematology*, *8*(5), 703–715. <https://doi.org/10.1163/156854106778877857>
- Sánchez-Moreno, S., Nicola, N. L., Ferris, H., & Zalom, F. G. (2009). Effects of agricultural management on nematode-mite assemblages: Soil food web indices as predictors of mite community composition. *Applied Soil Ecology*, *41*(1), 107–117. <https://doi.org/10.1016/j.apsoil.2008.09.004>
- Sieriebriennikov, B., Ferris, H., & de Goede, R. G. M. (2014). NINJA: An automated calculation system for nematode-based biological monitoring. *European Journal of Soil Biology*, *61*, 90–93. <https://doi.org/10.1016/j.ejsobi.2014.02.004>
- Six, J., Frey, S. D., Thiet, R. K., & Batten, K. M. (2006). Bacterial and fungal contributions to carbon sequestration in agroecosystems. *Soil Science Society of America Journal*, *70*(2), 555. <https://doi.org/10.2136/sssaj2004.0347>
- Thomson, B. C., Tisserant, E., Plassart, P., Uroz, S., Griffiths, R. I., Hannula, S. E., ...Lemanceau, P. (2015). Soil conditions and land use intensification effects on soil microbial communities across a range of European field sites. *Soil Biology and Biochemistry*, *88*, 403–413. <https://doi.org/10.1016/j.soilbio.2015.06.012>
- Toth, G., Montanarella, L., & Rusco, E. (2008). *Threats to soil quality in Europe EUR 23438. Scientific and technical research series* (pp. 61–74). Luxembourg: Office for Official Publications of the European Communities.
- Treonis, A. M., Austin, E. E., Buyer, J. S., Maul, J. E., Spicer, L., & Zasada, I. A. (2010). Effects of organic amendment and tillage on soil microorganisms and microfauna. *Applied Soil Ecology*, *46*(1), 103–110. <https://doi.org/10.1016/j.apsoil.2010.06.017>
- Treonis, A. M., Unangst, S. K., Kepler, R. M., Buyer, J. S., Cavigelli, M. A., Mirsky, S. B., & Maul, J. E. (2018). Characterization of soil nematode communities in three cropping systems through morphological and DNA metabarcoding approaches. *Scientific Reports*, *8*(1), 2004. <https://doi.org/10.1038/s41598-018-20366-5>
- Ugarte, C. M., Zaborski, E. R., & Wander, M. M. (2013). Nematode indicators as integrative measures of soil condition in organic cropping systems. *Soil Biology and Biochemistry*, *64*, 103–113. <https://doi.org/10.1016/j.soilbio.2013.03.035>
- van Diepeningen, A. D., de Vos, O. J., Korthals, G. W., & van Bruggen, A. H. C. (2006). Effects of organic versus conventional management on chemical and biological parameters in agricultural soils. *Applied Soil Ecology*, *31*(1), 120–135. <https://doi.org/10.1016/j.apsoil.2005.03.003>
- Vervoort, M. T. W., Vonk, J. A., Mooijman, P. J. W., Van den Elsen, S. J. J., Van Megen, H. H. B., Veenhuizen, P., ...Helder, J. (2012). SSU ribosomal DNA-based monitoring of nematode assemblages reveals distinct seasonal fluctuations within evolutionary heterogeneous feeding guilds. *PLoS ONE*, *7*(10), e47555. <https://doi.org/10.1371/journal.pone.0047555>
- Villenave, C., Saj, S., Pablo, A.-L., Sall, S., Djigal, D., Chotte, J.-L., & Bonzi, M. (2010). Influence of long-term organic and mineral fertilization on soil nematofauna when growing *Sorghum bicolor* in Burkina Faso. *Biology and Fertility of Soils*, *46*(7), 659–670. <https://doi.org/10.1007/s00374-010-0471-y>
- Waite, I. S., O'Donnell, A. G., Harrison, A., Davies, J. T., Colvan, S. R., Ekschmitt, K., ...Boström, S. (2003). Design and evaluation of nematode 18S rDNA primers for PCR and denaturing gradient gel electrophoresis (DGGE) of soil community DNA. *Soil Biology and Biochemistry*, *35*(9), 1165–1173. [https://doi.org/10.1016/S0038-0717\(03\)00177-9](https://doi.org/10.1016/S0038-0717(03)00177-9)
- Wang, K.-H., McSorley, R., & Gallaher, R. (2004). Relationship of soil management history and nutrient status to nematode community structure. *Nematropica*, *34*(1), 83–95.
- Wang, K. H., McSorley, R., Marshall, A., & Gallaher, R. N. (2006). Influence of organic *Crotalaria juncea* hay and ammonium nitrate fertilizers on soil nematode communities. *Applied Soil Ecology*, *31*(3), 186–198. <https://doi.org/10.1016/j.apsoil.2005.06.006>
- WRB, I. W. G. (2014). *World Reference Base for Soil Resources 2014. International soil classification system for naming soils and creating legends for soil maps*. World soil resources reports no. 106. Rome, Italy: FAO.
- Yeates, G. W., & Bongers, T. (1999). Nematode diversity in agroecosystems. *Agriculture, Ecosystems & Environment*, *74*(1–3), 113–135. [https://doi.org/10.1016/S0167-8809\(99\)00033-X](https://doi.org/10.1016/S0167-8809(99)00033-X)
- Yeates, G. W., Bongers, T., Goede, R. G. M., Freckman, D. W., & Georgieva, S. S. (1993). Feeding habitats in soil families and genera—An outline for soil ecologist. *Journal of Nematology*, *25*, 315–331.
- You, D., Tian, P., Sui, P., Zhang, W., Yang, B., & Qi, H. (2017). Short-term effects of tillage and residue on spring maize yield through regulating root-shoot ratio in Northeast China. *Scientific Reports*, *7*(1), 13314. <https://doi.org/10.1038/s41598-017-13624-5>
- Zhang, S., Li, Q., Lü, Y., Sun, X., Jia, S., Zhang, X., & Liang, W. (2015). Conservation tillage positively influences the microflora and microfauna in the black soil of Northeast China. *Soil and Tillage Research*, *149*, 46–52. <https://doi.org/10.1016/j.still.2015.01.001>
- Zhao, J., & Neher, D. A. (2013). Soil nematode genera that predict specific types of disturbance. *Applied Soil Ecology*, *64*, 135–141. <https://doi.org/10.1016/j.apsoil.2012.11.008>
- Zhong, S., Zeng, H.-C., & Jin, Z.-Q. (2017). Influences of different tillage and residue management systems on soil nematode community composition and diversity in the tropics. *Soil Biology and Biochemistry*, *107*, 234–243. <https://doi.org/10.1016/j.soilbio.2017.01.007>
- Zuur, A. F. (2009). *Mixed effects models and extensions in ecology with R*. New York, NY: Springer.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Bongiorno G, Bodenhausen N, Bünemann EK, et al. Reduced tillage, but not organic matter input, increased nematode diversity and food web stability in European long-term field experiments. *Mol Ecol*. 2019;28:4987–5005. <https://doi.org/10.1111/mec.15270>