## **ETH** zürich

# Is There a Scientific Rationale for the Refixation of Delaminated Chondral Flaps in Femoroacetabular Impingement? A Laboratory Study

**Journal Article** 

Author(s): Levinson, Clara; Naal, Florian D.; Salzmann, Gian M.; Zenobi-Wong, Marcy; Leunig, Michael

Publication date: 2020-04

Permanent link: https://doi.org/10.3929/ethz-b-000407130

Rights / license: Creative Commons Attribution-NonCommercial 4.0 International

**Originally published in:** Clinical Orthopaedics and Related Research 478(4), <u>https://doi.org/10.1097/CORR.000000000001135</u>

**Funding acknowledgement:** 159783 - A Bio-inspired Cartilage Mimetic with Enhanced Chondrogenic and Anti-inflammatory Properties (SNF)

This page was generated automatically upon download from the <u>ETH Zurich Research Collection</u>. For more information, please consult the <u>Terms of use</u>.

### Is There a Scientific Rationale for the Refixation of Delaminated Chondral Flaps in Femoroacetabular Impingement? A Laboratory Study

Running Title: Refixation of Delaminated Chondral Flaps

Clara Levinson PharmD, Florian D. Naal MD, Gian M. Salzmann MD, Marcy Zenobi-Wong Prof Dr, Michael Leunig MD

C. Levinson, M. Zenobi-Wong Tissue Engineering + Biofabrication, Swiss Federal Institute of Technology (ETH), Zürich, Switzerland

F. D. Naal, G. M. Salzman, M. Leunig Department of Orthopaedic Surgery, Hip and Knee Department, Schulthess Clinic, Zürich, Switzerland

The institution of one or more of the authors (MZ-W) has received, during the study period, lov

Each author certifies that neither he or she, nor any member of his or her immediate family, has funding or commercial associations (consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted article.

All ICMJE Conflict of Interest Forms for authors and *Clinical Orthopaedics and Related Research*<sup>®</sup> editors and board members are on file with the publication and can be viewed on request.

*Clinical Orthopaedics and Related Research*<sup>®</sup> neither advocates nor endorses the use of any treatment, drug, or device. Readers are encouraged to always seek additional information, including FDA approval status, of any drug or device before clinical use.

Each author certifies that his or her institution approved the human protocol for this investigation and that all investigations were conducted in conformity with ethical principles of research.

This work was performed at Schulthess Klinik and ETH Zürich, Zürich, Switzerland.

M. Zenobi-Wong ⊠ Tissue Engineering + Biofabrication Laboratory ETH Zürich Otto-Stern-Weg 7, HPL J22 8093 Zürich, Switzerland Email: marcy.zenobi@hest.ethz.ch

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

#### 1 Abstract

2 Background Debonding of the acetabular cartilage is a characteristic type of hip damage found in cam type femoroacetabular impingement, which remains a treatment challenge. In 3 addition to resection, refixation of these flaps using fibrin sealants has been recently 4 suggested. However, there is only limited evidence available that the proposed refixation 5 6 method results in sufficient viable cartilage formation to ensure long-term grafting of the flap 7 and restored tissue function. 8 Questions/purposes To determine flap tissue characteristics that would justify refixation of delaminated chondral flaps with a fibrin sealant, we characterized (1) the extracellular matrix 9 10 (ECM) of chondral flaps in terms of chondrocyte viability and distribution of ECM components and (2) the chondrogenic potential of resident cells to migrate into fibrin and 11 12 produce a cartilaginous matrix. 13 Methods Ten acetabular chondral flaps and three non-delaminated control cartilage samples were resected during surgery. Chondrocyte viability was quantified using a live-dead assay. 14 15 To assess the ECM, histological staining of glycosaminoglycans, collagen II and collagen I allowed the qualitative study of their distribution. Chondrocyte ability to migrate out of the 16 17 ECM was tested by encapsulating minced flap cartilage in fibrin gels and semi-quantitatively assessing the projected area of the gel covered with migrating cells. The potential of 18 19 chondrocytes to produce a cartilaginous matrix was studied with a pellet assay, a standard 3-20 D culture system to test chondrogenesis. Positive controls were pellets of knee chondrocytes 21 of age-matched donors, which we found in a previous study to have a good capacity to 22 produce cartilage matrix. Statistical significance of controlled quantitative assays was

- 23 determined by the Student t test with Welch's correction.
- 24 *Results* The proportion of viable chondrocytes in flaps was lower than in non-delaminated AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

25	cartilage (50% $\pm$ 19% vs 76 $\pm$ 6%; p = 0.02). Histology showed a disrupted ECM in flaps as					
26	compared to non-delaminated controls, with the presence of fibrillation, a loss of					
27	glycosaminoglycan at the delaminated edge, collagen II throughout the whole thickness of					
28	the flap and some collagen I-positive area in two samples. The resident chondrocytes were					
29	able to migrate out of this disrupted ECM in all tested samples. However in pellet culture,					
30	cells isolated from the flaps showed a qualitatively lower chondrogenic potential compared					
31	with positive controls, with a clearly inhomogeneous cell and matrix distribution and an					
32	overall smaller projected area (0.4 versus 0.7 mm <sup>2</sup> ; $p = 0.038$ ).					
33	Conclusion Despite the presence of viable chondrocytes with migration potential, the cells					
34	reside in a structurally altered ECM and have limited capacity to deposit ECM, leading us to					
35	question their capacity to produce sufficient ECM within the fibrin sealant for stable long-					
36	term attachment of such flaps.					

- 37 Clinical Relevance The characterization of delaminated cartilage in cam FAI patients
- 38 suggests that the refixation strategy might be adversely influenced by the low level of ECM
- 39 produced by the residing cells.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

#### 40 Introduction

41 Cartilage delamination (also referred to as a "cartilage flap") is a frequent finding of 42 acetabular cartilage damage secondary to cam type femoroacetabular impingement (FAI) and is classified as Stage 4 damage according to Beck et al. [5]. The presence of such flaps has 43 been documented in 37% to 85% of patients undergoing surgery [2, 4, 10, 20, 21, 31], with an 44 45 increased incidence in male patients [2, 31]. Despite an increased failure rate of open surgery 46 for patients whose hips demonstrate cartilage damage, there is no consensus on how to best treat chondral flaps. Débridement (removal) of cartilage is used to resolve symptoms and 47 allow for regrowth of repair cartilage. Conversely, reattachment (refixation) of cartilage with 48 a fibrin sealant would theoretically prevent migration of the femoral head in the defect and 49 50 thus joint space loss. Refixation could also potentially augment the reintegration of the 51 cartilage if viable chondrocytes secreted sufficient extracellular matrix (ECM) to strengthen the interface between the flap and the subchondral bone. 52 53 Recently, viable cells in flap tissues were reported, although the reported viability values vary 54 widely (from 39% to 95%) [11, 18, 35, 43]. Additional efforts have been made to characterize 55 the matrix composition of the chondral flaps with histologic [23, 35] and biochemical [18] 56 studies. These results have consistently highlighted the fibrocartilaginous nature of the ECM 57 within the flaps with fibrillations and loss of glycosaminoglycan (GAG) staining. Still, none 58 of these studies have had optimal controls, nor did they aim to demonstrate whether the 59 residual cells could actively produce cartilaginous ECM. The production of ECM would be important to support refixation and chondral flap repair. 60

- 61 Therefore, we investigated flap tissue characteristics in patients with FAI that would support
- 62 the refixation of delaminated chondral flaps with fibrin sealants, namely (1) the quality of the
- 63 ECM of chondral flaps as measured by chondrocyte viability and distribution of ECM

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

Kommentiert [TjG1]: AU: Fair to add this qualifier? ED AU response: yes components compared with controls and (2) the chondrogenic potential of resident cells interms of their ability to migrate into fibrin and produce a cartilaginous matrix.

66 Materials and Methods

67 Experimental Overview

68 This study had two main goals: First, to assess the quality of the ECM of chondral flaps <u>(the</u>

69 anatomical position of such flaps is schematized in Fig. 1A ; the orientation of resected flaps

70 with a suture is illustrated in Fig. 1B), and second, to evaluate the chondrogenic potential of

71 the cells residing in the flaps. For the first part, to assess the quality of the flap ECM, we

72 focused on (1) cell viability, as healthy tissue presumably contains a majority of viable cells,

73 and (2) cartilage ECM distribution, as the function of the tissue is linked to its structure. Both

tests were performed on sectioned tissue samples (Fig. 1C), either a slice of fresh tissue (for

viability) or a 5µm-thick slice of tissue embedded in paraffin (for histological staining of

76 ECM). In the second part, we investigated the chondrogenic potential of the residing cells by

studying (1) their migration capacity and (2) their ability to produce cartilaginous matrix.

78 Considering the poor stability of fibrin glue [13], attachment of the flap to the subchondral

79 bone relies on the premise that cells residing in flaps can migrate into the fibrin and

subsequently deposit matrix. The capacity of the cells to migrate out of the ECM into fibrin

81 glue was tested on minced flap pieces embedded in fibrin gels (Fig. 1D). The capacity of the

82 cells to produce cartilaginous matrix was assessed on cells enzymatically isolated from the

flaps, expanded and cultured in pellets (Fig. 1E). Of note, the small size of the flap tissues did

84 not allow us to perform all four assays with sections (viability, ECM composition), pieces

85 (migration) and cells (ECM deposition) of the same piece of tissue (Table 1). This explains

86 why the sample size is not 10 for all assays.

87 Sample Collection

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

Kommentiert [CO2]: AU: Figures must appear in proper alphanumeric order, that is, Fig. 1C cannot appear in the text before Fig. 1A. Please reorder your figures.

Author response: Fig. 1A and Fig. 1B were not mentioned before, this was corrected by the addition lines 69-70.

Acetabular chondral flaps, resected during surgical procedures, were obtained from 10 88 patients (nine male and one female) who had a primary diagnosis of cam-type FAI with 89 90 Grade IV cartilage lesions according to Beck et al. [5]. Patients had no history of childhood 91 hip disease, no previous hip surgery, and were between 20 and 50 years old. The mean period 92 of symptoms was 3.0 +/- 1.9 years, the mean Tegner activity score [38] was 5.3 +/- 1.8 and 93 the mean calculated Tonnis grade was 0.3 +/- 0.5 (Table 2). All male patients underwent 94 surgical hip dislocation; the female patient was treated with hip arthroscopy (Table 2). Inclusion criteria were cam-type FAI without signs of instability (lateral central edge angle < 95 96 25°, acetabular index > 10°, neck shaft angle < 135°, antetorsion < 15°, see Table 3), as 97 measured from radiographs (Fig. 2). Patient consent was received and approved by the 98 Ethics Commission of the Canton of Zurich (Kantonale Ethikkommission, Kanton Zürich, approval number KEK-ZH 2013-0097). The highly fibrous parts appearing to be the labrum 99 (when the cartilage separation included the chondrolabral junction) were removed, and the 100 101 remaining cartilage was minced for further tests (Fig. 1). Control, non-delaminated acetabular 102 cartilage samples used for viability tests were obtained from patients undergoing either 103 arthrodesis or THA for avascular necrosis (AVN) with laterally adherent cartilage. The AVN 104 was due to a procedure performed for the diagnosis of slipped capital femoral epiphysis. 105 Although AVN can potentially lead to secondary cartilage alterations, the intraoperative 106 visual aspect showed the presence of areas with no obvious cartilage alterations, and this 107 patient was therefore included in the study. Adult knee chondrocytes (used as controls for the 108 pellet assays) were obtained from healthy-appearing (white-blueish color, no fibrillation, 109 glistening aspect) articular cartilage pieces obtained from TKAs (approval number KEK-ZH 110 2017-00381), for instance when the medial condyle is affected by varus arthritis but not the

111 lateral condyle.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

hat gelöscht: All clinical information about the enrolled patients, including the duration of symptoms, the activity level ...

#### hat gelöscht:

hat gelöscht: osteoarthritis degree,

**hat gelöscht:** no history of childhood hip disease, no previous hip surgery, and age between 20 and 50 years

#### hat gelöscht:

**hat gelöscht:** the type of surgeries and the acetabulum and femur radiographic measurements are provided in Tables 2 and 3 as well as Fig. 2

hat gelöscht:

#### 124 Chemicals

125 All chemicals were ordered from Sigma-Aldrich (Buchs, Switzerland) unless mentioned126 otherwise.

- 127 Chondrocyte Viability Quantitative Measurement with Live-dead Staining
- 128 Chondrocyte viability was investigated using a standard live-dead assay according to the
- 129 manufacturer's instructions, as previously described [27]. Viability was quantified with Fiji
- 130 software, calculated as the number of living cells divided by the total number of cells (live
- 131 and dead) and averaging three values for each patient's sample.
- 132 Quality of ECM Compared with Non-delaminated Controls Qualitative Assessment of
- 133 Matrix Components with Histology on Paraffin Sections
- 134 Oriented native tissue samples were fixed overnight in 4% formaldehyde, washed in
- 135 phosphate-buffered saline, dehydrated, and embedded in paraffin. Five-microns thick paraffin
- 136 sections were deparaffinized in xylene and rehydrated in a series of ethanol baths with
- 137 decreasing concentrations of ethanol. Immunostaining of type I and II collagens were
- 138 performed as previously described [27], using products from Abcam (Cambridge, United
- 139 Kingdom) unless specified otherwise. Briefly, after antigen retrieval with hyaluronidase,
- sections were blocked with 5% normal goat serum and primary antibodies, diluted in 1%
- 141 normal goat serum, were applied overnight at 4° C (rabbit anti-human Type II collagen
- 142 antibody, 600-401-104, 1:200/Type I collagen antibody, #ab138492, 1:1500). For negative
- 143 controls, slides were incubated without a primary antibody. Endogenous peroxidase activity
- 144 was quenched with 1% H<sub>2</sub>O<sub>2</sub> for 20 minutes before applying the secondary antibody (goat
- 145 anti-rabbit IgG-HPR ab6721, 1:1500 dilution). Antibodies were detected with an Abcam
- 146 DAB kit, as per the manufacturer's protocol. Staining with hematoxylin and eosin (H&E),

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

- 147 Alcian blue (pH 2.5) and Safranin O was performed in accordance with standard protocols.
- 148 All stainings were followed by dehydration and two baths of xylene before the slides were
- 149 mounted with a resin mounting media.
- 150 All samples were stained at the same time to avoid slight changes of intensity, which could
- 151 result from sequential staining. The qualitative assessment was a yes/no answer to the
- 152 following questions: (1) Are there fibrillations in the H&E sections? (2) Is a clear red staining
- 153 visible in the Safranin O sections? (3) Is there a gradient in Safranin O with increasing
- intensity from surface to deeper layer, as expected in healthy cartilage? (4) Is there a positive
- 155 type I and II collagen staining? All stainings were observed and described by three
- researchers (two blinded, one not blinded) to ensure interobserver repeatability.
- 157 Chondrogenic Potential of Flap Chondrocytes Semi-quantitative Assessment of
- 158 Chondrocyte Migration into Fibrin Gels
- 159 The capacity of the chondrocytes to migrate into a fibrin gel was investigated as described
- 160 previously [27]. Briefly, a piece of cartilage was placed in a 24-well, non-tissue culture plate
- 161 in a 4-mm-diameter polydimethylsiloxane mold that was subsequently filled with 25  $\mu$ L of
- 162 fibrin glue (2 mM CaCl2, 1 U/mL thrombin, 20 mg/ml fibrinogen. After crosslinking for 20
- 163 minutes in the incubator (37° C, 5% CO<sub>2</sub>, 95% humidity), 1 mL of culture medium was added
- 164 onto the scaffolds, which were cultured in free-floating conditions for 4 weeks and
- subsequently fixed in 4% paraformaldehyde for 1 hour. After cell permeabilization with
- 166 Triton X-100, the gels were stained with 4', 6-diamidino-2-phenylindole (DAPI) and
- 167 phalloidin-rhodamine for 1 hour in the dark at room temperature. Whole gels were imaged
- 168 with a Zeiss Axio Explorer using the ApoTome.2 (Carl Zeiss Microscopy, Munich, Germany)
- 169 at 10 x magnification. The projected area of the gel covered with outgrown cells was
- 170 quantified manually with Fiji software, and the migration area was calculated as the area

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

Kommentiert [TjG4]: AU: Did you mean INTERobserver reliability here? Did you perform a kappa/ICC analysis? ED

AU response: thank you for noticing the word error, we indeed meant "interobserver". No kappa/ICC analysis was perfomed since the observations were entirely descriptive and did not rely on a scoring system.

hat gelöscht: ra

**Kommentiert [CO5]:** AU: Please parenthetically provide the manufacturer name and location (city, state, and country).

Author response: this information was added

172	covered with outgrown cells divided by the total area available to the cells to outgrow (total				
173	area of the gel minus the area of the cartilage piece). Although the contrast of the cells on				
174	background is clear enough to delineate the migration area, the manual aspect of this				
175	quantification can lead to slight variability in the results. However, the goal of this				
176	measurement is to give an idea of the migration capacity of these cells within the pool of				
177	replicates and does not provide an exact threshold for a clinical recommendation.				
178	Semi-quantitative and Qualitative Assessment of Flap Chondrocytes Ability to Produce				
179	Cartilaginous Matrix - Chondrocyte Isolation, Expansion, and Differentiation in Pellets				
180	The chondrocytes pellet assay is a 3-D culture system, originally developed to test the				
181	chondrogenic potential of mesenchymal progenitor cells [22]. The assay mimics cell				
182	condensation during limb development and is widely used to test the capacity of				
183	chondrocytes to deposit cartilaginous matrix in a 3-D environment. Chondrocytes were				
184	isolated by digesting minced cartilage samples with collagenase overnight [14]. Chondrocyte				
185	viability in the resulting cell suspension was determined using a standard trypan blue				
186	exclusion assay, before plating the cells in a T25 flask and passaging them once at $80\%$				
187	confluency. Cells were then trypsinized once more (cells in pellets were thus at Passage 2)				
188	and 250,000 cells were pipetted and spun in conical bottom 96-well plates to form pellets,				
189	subsequently cultured for 4 weeks in chondrogenic medium containing 10 ng/mL				
190	transforming growth factor $\beta$ 3. Of note, control samples were healthy knee chondrocytes				
191	isolated from samples of patients of the same age range. Although these cells originated from				
192	a different joint, they have been shown in a previous study to produce cartilage in pellet				
193	culture at passage 2 [14], thus constituting good positive controls. Lastly, pellets were				
194	embedded in an optimal cutting temperature compound, snap frozen and cut with a cryostat				
195	into 5-µm-thick sections before staining as described above in the histology section.				
	AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.				

#### 196 Statistical Analysis

197 All experiments were performed with material from a minimum of five patients with FAI 198 (due to limited amount of tissue, a sample could most often not be used for all four assays), with at least two (outgrowth) and three (viability) technical replicates for quantitative assays. 199 200 Due to the exploratory nature of this study, the effect size could not be estimated, a 201 requirement for sample size calculation. Considering the limited availability of chondral 202 flaps, the study was performed on a relatively small cohort of patients. For viability 203 quantification, histology, and pellet assays, three controls were used due to the restricted 204 availability of healthy-looking tissue. The results of the quantitative assays are reported as the 205 mean  $\pm$  SD of the biological replicates (each single point is the average of the technical 206 replicates of each patient's sample). Statistical tests were performed with Prism (GraphPad 207 Software Inc, version 7.0.3). Normality was verified with the Shapiro-Wilk test. Because sample sizes were unequal, Student t-tests with Welch's correction were used to compare two 208 209 groups. To compare the results of the two viability assays (live-dead versus trypan blue), a paired, two-tailed t-test was performed. The correlation between viability and age was 210 211 evaluated with a non-parametric Pearson rank correlation test. Statistical significance was set at p < 0.05. 212

#### 213 Results

- 214 Cartilage Flaps Contain Fewer Viable Chondrocytes than Non-delaminated Hip Cartilage
- 215 Samples
- 216 In delaminated flaps, chondrocyte viability was decreased when compared with non-
- 217 delaminated hip cartilage samples  $(50\% \pm 19\% \text{ versus } 76 \pm 6\%; \text{ means difference } -26 [95\%]$
- 218 CI -46 to -6]; p = 0.02) (Fig. 3). There was no correlation between viability and the age of the

**Kommentiert [CO6]:** AU: Please see the query in the Fig. 3 legend.

219 patients (r = -0.19; p = 0.7). Viability quantification with a trypan blue exclusion assay on a AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

 $(84\% \pm 9\% \text{ versus } 54\% \pm 18\% \text{ in the same five chondral samples; mean difference } 30 [95\%]$ 221 222 CI 4 to 56]; p = 0.03) (Supplemental Fig. 1; supplemental materials are available with the 223 online version of CORR®). 224 Chondrocytes in Chondral Flaps Reside in a Disrupted ECM 225 In all seven flap samples tested for histology, the qualitative assessment of the tissue 226 organization with H&E staining showed the presence of a fibrillated ECM, which was present 227 either throughout the sample or located at the surface of the cartilage towards the delaminated edge of the flap, suggesting a gradient of degeneration (Fig. 4). Fibrillation was associated 228 229 with a loss of normal tissue stratification; that is, neither flattened cells in the superficial layer 230 nor columnar cells in the deeper layers were visible. In addition, this fibrillated area 231 correlated with a loss of GAG (no Safranin O staining). The gradient of GAG in healthy 232 cartilage, characterized by a progressively increasing amount of GAG toward deeper layers, 233 was consistently lost in all flat samples. Some GAG was retained along the attached side in 234 some samples, whereas in some samples, GAG could not be detected (Fig. 4). The loss of 235 GAG was not accompanied by a loss of collagen, as suggested by the strong intensity of the 236 picrosirius red staining, an anionic dye that associates to the cationic collagen fibrils (data not 237 shown). Patches of Type I collagen-positive staining were visible in two of the seven flap 238 samples used for histology, but only in fibrillated areas (Fig. 4). All samples had positive 239 staining results for Type II collagen, which was mostly present in the superficial zone and detected only pericellularly in the middle and deep zones of healthy cartilage. This 240 localization was lost in the flap samples, where Type II collagen was present throughout the 241 242 thickness, with a higher staining intensity than in healthy controls.

suspension of freshly isolated chondrocytes led to higher values than with live-dead staining

220

243 Chondrocytes Are Able to Migrate Out of Chondral Flaps into Fibrin Gels

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

**Kommentiert [CO7]:** AU: Please see the query in the Supplemental Fig. 1 legend.

Kommentiert [CO8]: AU: Please see the query in the Fig. 4 legend.

244	Chondrocyte migration out of the chondral ECM was evaluated semi-quantitatively upon
245	encapsulation in fibrin gels. The presence of cells migrating and spreading within the fibrin
246	gel was observed in all five flap samples tested. The capacity of the cells to "colonize" the
247	fibrin gel was further quantitatively assessed by measuring the proportion of the projected
248	area of the gel covered with migrating cells. This quantification showed that the mean
249	proportion of the gel where migrating chondrocytes were present was 71 $\pm$ 25%. In other
250	terms, the fibrin gel supported the outgrowth and the proliferation of flap chondrocytes,
251	resulting in an extended coverage of the gel with cells (Fig. 5A-B). These observations
252	indicate that the ECM permitted chondrocyte migration into fibrin glue, possibly because of
253	the above-described degraded state of the ECM [30]. There was variability among patients
254	and among the technical triplicates for each patient's sample (two or three independent fibrin
255	gels were made for each flap sample) (Fig. 5C).
256	Cells Isolated from Flaps have a Lower Capacity to Deposit Cartilaginous Matrix than
257	Controls
258	To semi-quantitatively describe the capacity of flap chondrocytes to produce cartilaginous
259	matrix, which would be important for the long-term fixation of the flap (considering the low
260	stability of fibrin sealants in vivo) and for the flap tissue repair once refixated, we compared
261	pellets of chondrocytes isolated from either flaps or from controls. Of the five biological
262	replicates for pellets isolated from flaps, four led to the formation of pellets with ECM
263	production, while one showed almost no matrix production (Fig. 6A). The qualitative
264	assessment of cell distribution show that cells were less homogeneous in pellets isolated from
265	flaps (presence of "cell patches") compared with control samples, as shown by the
266	hematoxylin counterstain. We also observed that the ECM produced in pellets from controls

267 was homogeneously distributed (Fig. 6B), in contrast to pellets from flaps, which displayed

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

**Kommentiert [CO9]:** AU: Please see the query in the Fig. 5 legend.

**Kommentiert [CO10]:** AU: Please see the query in the Fig. 6 legend.

268	"holes" of fainter staining. Type I collagen was present in all pellets. In addition, the pellets in
269	flaps were smaller than those in controls (Fig. 6D), as shown by their lower average surface
270	area compared to pellets from control samples (0.4 $mm^2\pm0.09versus~0.7\pm0.1~mm^2$
271	respectively, mean difference 0.3 [95% CI, 0.03 to 0.6]; $p = 0.038$ ). This observation,
272	together with the inhomogeneous patches of cells, suggests that flap chondrocytes
273	proliferated less compared to "healthy" knee chondrocytes, previously found to have a good
274	chondrogenic potential.

#### 275 Discussion

276 Recent clinical and experimental work has suggested that cam-type FAI frequently causes 277 cartilage delamination and the subsequent development of osteoarthritis of the hip. To avoid 278 such cartilage damage in severe cases of cam FAI, surgery may be advised [1]. During such 279 procedures, delamination of the acetabular cartilage is often encountered and the damaged 280 cartilage is generally resected. Based on the reported presence of various amounts of viable cells in chondral flaps [11, 18, 35, 43], refixation of these flaps with a fibrin sealant has also 281 282 been suggested as an alternative to resection. However, there is no well-defined scientific 283 rationale supporting this technique. Legitimate questions regarding the tissue characteristics 284 remain, notably whether there are enough viable cells in the flaps, whether the structure, and therefore the function, of the flap tissue is preserved, or whether the residing cells are able to 285 286 provide solid attachment to the subchondral bone. The present study augments cell viability 287 and histology data with a study of chondrocytes' capacity to outgrow from the ECM into 288 fibrin gels and produce a cartilaginous ECM. Our main findings are (1) a lower percentage of 289 viable chondrocytes were present in flap cartilage compared with controls; (2) the ECM in 290 flaps displayed an altered structure at the delaminated edge of the flap or throughout the 291 whole flap in some cases, probably induced by repeated mechanical insult; (3) the

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

292	chondrocytes could migrate out of the flap tissue into fibrin glue; and (4) flap chondrocyte			
293	have a limited capacity to produce ECM. These findings challenge the refixation practice and			
294	suggest the need for further mechanistic large animals and clinical studies focusing on the			
295	long-term outcome of flap refixation based on follow-up MRI.			

296 Limitations

297 We are aware that our study has several limitations. First, although the sample size is similar 298 to previous studies investigating chondral flaps [11, 18, 35], it remains too small to generalize 299 the results. This in vitro study is meant to be exploratory and to stimulate re-evaluation of this 300 clinical practice, rather than to provide a definitive surgical recommendation. Large animal 301 studies would provide insights into the repair mechanism and the long-term efficiency. 302 Additionally, the samples were small, which limited the number of assays that could be done 303 per sample. For instance, the cell yield after chondrocyte isolation and expansion in vitro was 304 up to 720,000 cells at Passage 2, which illustrates the small amount of living cells initially 305 present in the flaps and the need for a large amount of tissue to perform pellet assays 306 (250,000 cells per pellet). Despite this limitation, the sample size for each experiment was at 307 least 5 biological replicates, and the consistent results between these replicates can give 308 confidence into the presented results. 309 Another limitation is the use of two different types of controls. For the viability and ECM 310 composition assays, the goal was to assess whether the flap displayed a lower quality than 311 non-delaminated cartilage. Three non-delaminated acetabular cartilage samples could be 312 obtained. This low number of controls as compared with flaps is directly linked to the limited 313 availability of healthy-looking cartilage in these patients, and does not represent an obstacle 314 to performing statistical tests. These samples originated from hips of patients undergoing 315 surgery (arthrodesis, derotational femoral osteotomy and THA) with acetabular cartilage AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

316	appearing well fixed while the femoral head cartilage was affected (AVN of femoral head and				
317	anterior and lateral hip instability with suprafoveolar head damage). Histological staining of				
318	these controls displayed all the known characteristics of healthy acetabular cartilage at the				
319	tissue level and we, therefore, considered these as a fair control tissue. The question of flap				
320	chondrocytes' ability to deposit ECM was a yes/no question. Therefore, we included a				
321	positive control, namely knee chondrocytes known to effectively deposit matrix, to describe				
322	the features of pellets made of cells displaying a good chondrogenic potential. In a previous				
323	study, our laboratory has shown that the knee chondrocytes from donors in the same age-				
324	range as the FAI patients included in this study, could deposit a good quality of cartilaginous				
325	matrix at Passage 2 [14]. Consequently, although the mechanical environment of knee and				
326	hip cartilage is different, these knee chondrocytes are acceptable as a positive control in terms				
327	of chondrogenesis. It is important to mention that the pellet assay cannot fully predict the				
328	behavior of cells that migrated in the fibrin gel, given the 2-D in vitro expansion step required				
329	to obtain enough cells. Flap chondrocytes were shown to dedifferentiate during in vitro				
330	culture but to re-differentiate upon a switch to 3-D culture [11]. The pellet assay, although not				
331	using a hydrogel as a matrix, still provides a 3-D culture environment, which may				
332	compensate for the need of expansion. The observations in pellets can, however, be				
333	strengthened by the finding that the gels used for the migration assay were very soft and				
334	difficult to handle after the 4 weeks of culture, which strongly suggests the absence of matrix				
335	produced by the migrating cells. It is evident that the free-floating condition during the				
336	culture of the fibrin gels drastically differs from a hip environment [20, 26]. The free-floating				
337	condition can be a reason for absence of deposited ECM, but it is necessary to investigate				
338	specifically the property of the fibrin sealant to support migration of the cells growing out of				
339	the flaps.				

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

340	It was not possible to exactly define the duration of the mechanical degradation, but all				
341	patients were symptomatic for at least 1 year (Table 2). Considering the fact that the onset of				
342	FAI symptoms appear once advanced structural changes have happened, we assume that				
343	patients had lived at least 1 year with damaged articular cartilage. All patients had cam FAI				
344	(Fig. 2), which minimized population heterogeneity.				
345	Chondrocyte Viability in Cartilage Flaps and Quality of ECM				
346	We found lower chondrocyte viability in delaminated flaps compared with control tissues				
347	(50% $\pm$ 19% versus 76 $\pm$ 6%), confirming other reports of low viability in flaps (39% $\pm$ 14%				
348	[18] and 54.6% $\pm$ 25.6% [35]), although these data contrast with the high viability (87% $\pm$				
349	10%) found by Wright et al. [43]. We found there was higher viability in non-delaminated				
350	controls, which was not shown in an earlier report (38.8%) [35]. The presence of viable cells				
351	is important to consider when deciding whether to re-fixate chondral flaps since residing				
352	chondrocytes are key players in the regeneration process, through migration and proliferation				
353	[28, 34]. The variable results found in previous studies may be attributed to the quantification				
354	technique, because we observed higher viability with the trypan blue exclusion assay on				
355	isolated cells (Supplemental Fig. 1; supplemental materials are available with the online				
356	version of CORR) than with the live-dead assay on whole cartilage samples. Finally, if the				
357	reported viability in allografts after the maximum allowed storage time of 45 days (~ $66\%$				
358	[39]) was considered as a threshold for tissue implantation or refixation, the viability in flaps				
359	would be too low to be considered for refixation.				
360	Our study showed that chondrocytes in chondral flaps exist in a degraded ECM environment.				
361	A novel aspect of this work is the observation of the flaps in their entirety, which highlighted				
362	a loss of GAG and the presence of fibrillation either only at the delaminated edge of the flap				

363 or in the whole sample in the worst cases. The structural organization of the ECM, crucial for AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

364	the function of the cartilage as a load absorber [16], is lost throughout the flap, not only at the				
365	surface or delaminated edge. These features likely revealed a mechanically damaged tissue,				
366	as described by Kurz et al. [24]. The presence of various amounts of degenerated, fibrillated				
367	tissue in chondral flaps has already been reported, although without consideration of the				
368	whole flap structure [23, 35]. Our study confirmed the GAG depletion [18, 35], loss of				
369	columnar distribution of the cells, and the disorganized structure of the ECM [20, 35] that has				
370	been described. The drastic loss of GAG provides an environment in which cells are more				
371	prone to die in case of mechanical injury [32], which could explain the low viability				
372	observed. Our findings regarding the increased area stained with Type II collagen in all tested				
373	samples are in line with a published biochemical quantification of collagens in flaps, which				
374	showed increased amounts of collagen in flaps that have a degenerated gross appearance [18].				
375	However, previously published histology data in flaps showed a loss of Type II collagen in				
376	flap samples with high Osteoarthritis Research Society International grades [35]. Increased				
377	Type II collagen production seems more consistent with the known correlation between				
378	mechanical stimulation and anabolic metabolism of chondrocytes [8] on the one hand, and				
379	between FAI and development of osteoarthritis on the other. During the early stage of				
380	osteoarthritis, the collagen synthesis rate increases [41] and a transition from Type II to Type				
381	I collagen occurs [25], which motivated our assessment of the presence of Type I collagen, as				
382	it had not been done before. We observed the presence of Type I collagen in two of seven				
383	samples in the form of islands; one of these two samples originated from a patient with a				
384	Tönnis Grade of 1.				

**385** Regeneration Potential of Cells in Cartilage Flaps, in Terms of Migration and Matrix

386 Deposition

387 We observed chondrocyte migration out of the flap ECM into fibrin in all five biological

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

388	samples, notwithstanding the relatively high fibrin concentration in gels, which is used to				
389	increase the gel's stability [15]. Many in vitro studies have investigated the capacity of				
390	chondrocytes to migrate from healthy tissue into fibrin glue as a repair strategy for cartilage				
391	lesions [3, 9, 42, 44]. Absence of outgrowth has been reported [3, 44], unless additional				
392	mechanical [42] or chemical [9] cues were provided. The reported need for an ECM digestion				
393	step to promote cellular outgrowth from a healthy ECM in cartilage [29, 33, 36] strengthens				
394	our theory that the loss of GAG and degenerative features in flap samples promote				
395	outgrowth. This could be an argument in favor of refixation; however, the fibrin gels after				
396	culture were very soft, which is most probably because of the absence of ECM production by				
397	the outgrown cells. We thus investigated the capacity of residing chondrocytes to produce				
398	cartilaginous matrix with a pellet assay.				
399	When testing the ECM deposition by cells isolated from flaps, we observed not only limited				
400	deposition of cartilaginous matrix but also Type I collagen. Pellets of flap chondrocytes				
401	displayed smaller sizes as well as less-homogeneously distributed cells and ECM compared				
402	with positive control pellets. The limited capacity to produce cartilaginous ECM, even in the				
403	presence of supraphysiologic amounts of chondrogenic transforming growth factor $\beta$ 3, is a				
404	drawback to the refixation of flaps as they may not be able to deposit ECM within the fibrin				
405	sealant to strengthen the fixation to the subchondral bone. These findings are in contrast to				
406	those of Giannini et al. [17] who demonstrated the promising use of cells harvested from				
407	detached osteochondral fragments for autologous chondrocyte implantation procedures in the				
408	ankle. However, in their study, the histologic appearance of the detached tissue appeared to				
409	be hyaline-like, in contrast to the disrupted ECM of the flaps. Biant and Bentley [7]				
410	investigated the chondrogenic potential of cells isolated from cartilage debrided during				
411	autologous chondrocyte implantation and cultured until Passage 2. They showed that DNA,				

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

412	GAG, and Type II collagen amounts in pellets of chondrocytes from debrided samples were			
413	similar to the amounts obtained in pellets of chondrocytes from healthy, non-load-bearing			
414	cartilage samples. The results of Bretschneider et al. [11] also indicate that flap chondrocytes			
415	de-differentiate but can recover their phenotype upon encapsulation in a chondro-inductive			
416	biomaterial. Nevertheless, fibrin was shown to fail at promoting chondrocyte redifferentiation			
417	[6], which challenges the use of this biomaterial. Of note, the study of ECM deposition by			
418	flaps chondrocytes cannot predict whether the flap will heal upon refixation, due to the			
419	technical limitation of chondrocyte expansion in vitro. However, it can be expected that the			
420	flaps would not heal, considering the well-known limited capacity of cartilage tissue to			
421	regenerate upon damage [12] and the very slow collagen turnover in healthy and pathologic			
422	joints [19]. Studies reporting results of flap refixation do not focus on the quality of the			
423	refixated tissue or the newly generated tissue at the interface with the subchondral bone, but			
424	rather on clinical scores [37, 40].			
425	Conclusions			

We found that chondrocytes isolated from hip flaps can outgrow into fibrin glue, however, their cell viability is low (< 70%) and they have limited capacity to deposit cartilage matrix. These data suggest that the fibrin refixation approach should be questioned, as it may not lead to long-term attachment and repair; however, the results should be considered cautiously due to the small sample size. A study in a large animal model would give more realistic insights into the long-term efficacy of refixation with fibrin to ultimately provide a more definitive surgical recommendation.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

#### Acknowledgments

We thank Prof. Dr. med Matthias Steinwachs for providing the knee cartilage samples and Dr. Emma Cavalli for isolating the chondrocytes from these knee cartilage samples. The authors acknowledge the use of the Scientific Center for Optical and Electron Microscopy for histology imaging. Finally, we thank Angela Bonato, <u>MSc</u> and Benjamin Kessel, <u>MSc</u> for their blinded observations of the histological sections.

**Kommentiert [CO11]:** AU: If possible, please provide the final academic degree for each of these individuals.

Author response: The two missing titles were added.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

#### References

- 433 [1] Agricola R, Waarsing JH, Arden NK, Carr AJ, Bierma-Zeinstra SMA, Thomas GE,
- 434 Weinans H, Glyn-Jones S. Cam impingement of the hip a risk factor for hip
- 435 osteoarthritis. *Nat Rev Rheumatol.* 2013;9:630–634.
- 436 [2] Anderson LA, Peters CL, Park BB, Stoddard GJ, Erickson JA, Crim JR. Acetabular
- 437 cartilage delamination in femoroacetabular impingement. Risk factors and magnetic
- 438 resonance imaging diagnosis. *J Bone Joint Surg Am.* 2009;91:305–313.
- 439 [3] Andjelkov N, Hamberg H, Bjellerup P. No outgrowth of chondrocytes from non-digested
- 440 particulated articular cartilage embedded in commercially available fibrin matrix: An in
- 441 vitro study. J Orthop Surg Res. 2016;11:1–5.
- 442 [4] Beaulé PE, Le Duff MJ, Zaragoza E. Quality of life following femoral head-neck
- 443 osteochondroplasty for femoroacetabular impingement. *J Bone Joint Surg Am.*444 2007;89:773–779.
- 445 [5] Beck M, Leunig M, Parvizi J, Boutier V, Wyss D, Ganz R. Anterior Femoroacetabular
- 446 Impingement: Part II. Midterm Results of Surgical Treatment. *Clin Orthop Relat Res.*447 2004;418:67–73.
- 448 [6] Bianchi VJ, Lee A, Anderson J, Parreno J, Theodoropoulos J, Backstein D, Kandel R.
- 449 Redifferentiated Chondrocytes in Fibrin Gel for the Repair of Articular Cartilage
  450 Lesions. *Am J Sports Med.* 2019;47:2348–59.
- 451 [7] Biant LC, Bentley G. Stem cells and debrided waste: Two alternative sources of cells for
- 452 transplantation of cartilage. *J Bone Joint Surg Br.* 2007;89:1110–1114.
- 453 [8] Bleuel J, Zaucke F, Brüggemann GP, Niehoff A. Effects of cyclic tensile strain on

454 chondrocyte metabolism: a systematic review. *PLoS One.* 2015;10:e0119816.
 AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

- [9] Bonasia DE, Marmotti A, Rosso F, Collo G, Rossi R. Use of chondral fragments for one
  stage cartilage repair: A systematic review. *World J Orthop. 2015;6*:1006–1011.
- 457 [10] Bredella MA, Stoller DW. MR imaging of femoroacetabular impingement. *Magn Reson*
- 458 Imaging Clin N Am. 2005;13:653–664.
- 459 [11] Bretschneider H, Stiehler M, Hartmann A, Boger E, Osswald C, Mollenhauer J,
- 460 Gaissmaier C, Günther KP. Characterization of primary chondrocytes harvested from
- 461 hips with femoroacetabular impingement. *Osteoarthritis Cartilage*. 2016;24:1622–1628.
- 462 [12] Buckwalter JA, Mankin HJ. Articular Cartilage. Part II: Degeneration and
- 463 Osteoarthrosis, Repair, Regeneration, and Transplantation. *J Bone Joint Surg Am*.
  464 1997;79:612-632.
- 465 [13] Cassar-Gheiti AJ, Byrne DP, Kavanagh E, Mulhall KJ. Comparison of four chondral
- repair techniques in the hip joint: A biomechanical study using a physiological human
  cadaveric model. *Osteoarthritis Cartilage*. 2015;23:1018–1025.
- 468 [14] Cavalli E, Levinson C, Hertl M, Broguier N, Brück O, Mustjoki S, Gerstenberg A,
- 469 Weber D, Salzmann G, Steinwachs M, Barreto G, Zenobi-Wong M. Characterization of
- 470 polydactyly chondrocytes and their use in cartilage engineering. *Sci Rep.* 2019;9:1–15.
- 471 [15] Douglas AM, Fragkopoulos AA, Gaines MK, Lyon LA, Fernandez-Nieves A, Barker
- 472 TH. Dynamic assembly of ultrasoft colloidal networks enables cell invasion within
- 473 restrictive fibrillar polymers. *Proc Natl Acad Sci USA*. 2017;114:885–890.
- 474 [16] Fox JS, Bedi A, Scott AR. The Basic Science of Articular Cartilage: structure,
- 475 composition and function. *Sports Health*. 2009;1:461–468.
- 476 [17] Giannini S, Buda R, Grigolo B, Vannini F, De Franceschi L, Facchini A. The detached
- 477 osteochondral fragment as a source of cells for autologous chondrocyte implantation
   AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

478 (ACI) in the ankle joint. Osteoarthritis Cartilage. 2005;13:601–607.

- 479 [18] Hariri S, Truntzer J, Smith RL, Safran MR. Biochemical and cellular assessment of
- 480 acetabular chondral flaps identified during hip arthroscopy. *Arthroscopy*. 2015;31:1077–
  481 1083.
- 482 [19] Heinemeier KM, Schjerling P, Heinemeier J, Moller MB, Krogsgaard MR, Grum-
- 483 Schwensen T, Petersen MM, Kjaer M. Radiocarbon dating reveals minimal collagen
- 484 turnover in both healthy and osteoarthritic human cartilage." *Sci Transl Med.* 2016;
  485 8:346ra90.
- 486 [20] Jannelli E, Parafioriti A, Acerbi A, Ivone A, Fioruzzi A, Fontana A. Acetabular
  487 Delamination: Epidemiology, Histological Features, and Treatment. *Cartilage*.
- 488 2018;10:314-320.

493

489 [21] Johnston TL, Schenker ML, Briggs KK, Philippon MJ. Relationship Between Offset

Angle Alpha and Hip Chondral Injury in Femoroacetabular Impingement. *Arthroscopy*.
2008;24:669–675.

492 [22] Johnstone B, Hering TM, Caplan AI, Goldberg VM, Yoo JU. In Vitro Chondrogenesis of

Bone Marrow-Derived Mesenchymal Progenitor Cells. Exp Cell Res. 1998;238:265-72.

- 494 [23] Kohl S, Hosalkar HS, Mainil-Varlet P, Krueger A, Buechler L, Siebenrock K. Histology
- 495 of damaged acetabular cartilage in symptomatic femoroacetabular impingement: An
  496 observational analysis. *Hip Int.* 2011;21:154–162.
- 497 [24] Kurz B, Lemke AK, Fay J, Pufe T, Grodzinsky AJ, Schünke M. Pathomechanisms of
- 498 Cartilage Destruction by Mechanical Injury. *Ann Anat.* 2005;187:473–85.
- 499 [25] Lahm A, Mrosek E, Spank H, Erggelet C, Kasch R, Esser J, Merk H. Changes in content

500 and synthesis of collagen types and proteoglycans in osteoarthritis of the knee joint and AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

- 501 comparison of quantitative analysis with Photoshop-based image analysis. *Arch Orthop*
- 502 *Trauma Surg.* 2010;130:557–564.
- 503 [26] Lavigne M, Parvizi J, Beck M, Siebenrock KA, Ganz R, Leunig M. Anterior
- Femoroacetabular Impingement: part I. Techniques of joint preserving surgery. *Clin Orthop Relat Res.* 2004;418:61-66.
- 506 [27] Levinson C, Cavalli E, Sindin DM, Kessel B, Zenobi-Wong M, Preiss S, Salzmann G,
- 507 Neidenbach P. Chondrocytes From Device-Minced Articular Cartilage Show Potent
- 508 Outgrowth Into Fibrin and Collagen Hydrogels." *Orthop J Sports Med.* 2019;7:1–12.
- 509 [28] Lu Y, Dhanaraj S, Wang Z, Bradley DM, Bowman SM, Cole BJ, Binette F. Minced
- 510 Cartilage without Cell Culture Serves as an Effective Intraoperative Cell Source for
- 511 Cartilage Repair. J Orthop Res. 2006;24:1261–1270.
- 512 [29] McGregor AJ, Amsden BG, Waldman SD. Chondrocyte repopulation of the zone of
- 513 death induced by osteochondral harvest. *Osteoarthritis Cartilage*. 2011;19:242–248.
- 514 [30] Morales TI. Chondrocyte moves: clever strategies? Osteoarthritis Cartilage.
- 515 2007;15:861-871.
- 516 [31] Nepple JJ, Carlisle JC, Nunley RM, Clohisy JC. Clinical and radiographic predictors of
- 517 intra-articular hip disease in arthroscopy. *Am J Sports Med.* 2011;39:296–303.
- 518 [32] Otsuki S, Brinson DC, Creighton L, Kinoshita M, Sah RL, D'Lima D, Lotz M. The
- effect of glycosaminoglycan loss on chondrocyte viability: A study on porcine cartilage
  explants. *Arthritis Rheum*. 2008;58:1076–1085.
- 521 [33] Qiu W, Murray MM, Shortkroff S, Lee CR, Martin SD, Spector M. Outgrowth of
- 522 chondrocytes from human articular cartilage explants and expression of alpha-smooth
- 523 muscle actin. Wound Repair Regen. 2000;8:383–391.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

524	[34] Ribitsch I, Mayer RL	, Egerbacher M,	Gabner S, Kańduła N	IM, Rosser J. Fetal articular
-----	---------------------------	-----------------	---------------------	-------------------------------

- 525 cartilage regeneration versus adult fibrocartilaginous repair: secretome proteomics
- 526 unravels molecular mechanisms in an ovine model. *Dis Model Mech.* 2018;11:1–11.
- 527 [35] Rodriguez-Fontan F, Payne KA, Chahla J, Mei-Dan O, Richards A, Uchida S, Pascual-
- 528 Garrido C. Viability and tissue quality of cartilage flaps from patients with

529 femoroacetabular hip impingement. Orthop J Sports Med. 2017;5:1–10.

- 530 [36] Sage A, Chang AA, Schumacher BL, Sah RL, Watson D. Cartilage outgrowth in fibrin
- 531 scaffolds. *Am J Rhinol Allergy*. 2009;23:486–491.
- 532 [37] Stafford GH, Bunn JR, Villar RN. Arthroscopic repair of delaminated acetabular
- 533articular cartilage using fibrin adhesive. Results at one to three years. *Hip Int.*
- **534** 2011;21:744–750.
- [38] Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res.* 1985;198:43-49.
- 537 [39] Torrie AM, Kesler WW, Elkin J, Gallo RA. Osteochondral allograft. Curr Rev
- 538 *Musculoskelet Med.* 2015;8:413–422.
- 539 [40] Tzaveas AP, Villar RN. Arthroscopic repair of acetabular chondral delamination with
- 540 fibrin adhesive. *Hip Int.* 2010;20:115–119.
- 541 [41] Videman T, Eronen I, Candolin T. [3 H]proline incorporation and hydroxyproline
- 542 concentration in articular cartilage during the development of osteoarthritis caused by
- 543 immobilization. A study in vivo with rabbits. *Biochem J.* 2015;200:435–440.
- 544 [42] Wang N, Grad S, Stoddart MJ, Niemeyer P, Reising K, Schmal H, Südkamp NP, Alini
- 545 M, Salzma, GM. Particulate cartilage under bioreactor-induced compression and shear.
- 546 *Int Orthop.* 2014;38:1105–1111.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

- 547 [43] Wright VJ, McCrum CL, Li H, Tranovich MJ, Huard J. Significant Chondrocyte
- 548 Viability Is Present in Acetabular Chondral Flaps Associated With Femoroacetabular
- 549 Impingement. Am J Sports Med. 2018;46:149–152.
- 550 [44] Zingler C, Carl HD, Swoboda B, Krinner S, Hennig F, Gelse K. Limited evidence of
- chondrocyte outgrowth from adult human articular cartilage. *Osteoarthritis Cartilage*.
  2015;24:124–128.

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

#### Legends

**Fig. 1** This figure shows an overview of the study's procedures. (A) Chondral flaps were harvested from the hips of adult patients with FAI and sutured on the delaminated edge. (B) We used oriented hip flaps to investigate the properties of (C) whole tissue pieces (viability and ECM distribution), (D) particulated pieces embedded in fibrin glue, casted in a PDMS (polydimethylsiloxane) for the migration assay, and (E) isolated chondrocytes cultured in vitro (chondrogenesis in pellets).

**Fig. 2** Radiographs of hips from patients included in the study. The sex (M/F) and age are indicated at the top of the radiograph. The sample number is placed at the bottom right of the images.

**Fig. 3** Chondrocyte viability was decreased in flaps samples compared with controls. (A) These representative images show the (I) highest and (II) lowest microscopic images of livedead staining of non-delaminated hip samples (top) and flaps (bottom). Live cells are depicted in green and dead cells in red. The sex, age, and viability result are given at the bottom left. Scale bar: 100  $\mu$ m. (B) This summary table shows the viability results for each sample (six biologic replicates for flaps and three for controls, with three technical replicates), obtained from quantification of live-dead images. (C) This plot shows the viability results (dots represent the average of the three technical replicates of each donor analyzed), comparing viability between non-delaminated controls and flaps (unpaired t-test with Welch's correction, \* p < 0.05).

**Fig. 4** The ECM structure was fibrillated and depleted of glycosaminoglycan (GAG) in chondral flaps. Histologic staining of 5-μm-thick sections of oriented flap samples are shown. From left to right: these images show hematoxylin and eosin, Safranin O (GAG stains), and

immunohistochemistry results for Type II and Type I collagen (hematoxylin counterstain, AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

**Kommentiert [CO12]:** AU: In the figure, please use sentence case for all text, meaning that only the first letter of the first word is capitalized, unless there is a proper now.

In Fig. 1C, please delete the ampersand symbol and use the word "and" instead.

Author response: We implemented the changes, notably by capitalizing the following:

- Fig. 1A: "Marks"
- Fig. 1C: "Superficial"; "Attached"
  Fig. 1D: "Minced" and "Casting"
- Fig. 1E: "Isolation", "Expansion" and "Centrifugation"
- Fig. IE. Isolation, Expansion and Centifugation

**Kommentiert [CO13]:** AU: In the figure, please capitalize the first letter of "Female" and "Male" at each instance.

Please put a space before and after the equal sign in the p value.

Please include a note about the asterisk in the legend.

Author response: we capitalized the letters as requested and added the spaces around the "=" sign. The note about the asterisk was added in the legend in replacement of the exact p value, which is already mentioned in the figure.

hat gelöscht: p = 0.02,

positive staining in brown). The \* is positioned at the side of delamination, on the side of the acetabular cartilage surface. The first row shows the staining of a non-delaminated hip cartilage sample and the three following rows show three representative images of flap samples (sex and age indicated on the left side), from the best- (M21) to the worst- (M32) looking conditions, from top to bottom. On the hematoxylin and eosin staining pictures, the dotted line separates the fibrillated and smooth areas. n = 7 oriented samples and n = 3 control samples. Scale bars: 500 µm; H&E = hematoxylin and eosin.

**Fig. 5** This figure shows the outgrowth potential of chondrocytes in flap samples. (A) A scoring system was used to investigate the outgrowth potential of the cells. Gels were imaged with 10 x magnification, and outgrowth values were calculated as the ratio of the area covered with cells (Ac) to the total area of the gel (Ag). (B) This plot shows the outgrowth score (five biologic replicates with two or three technical replicates for each biologic replicate). (C) These image shows an example of the best score (left), an intermediate score (middle), and a low score (right). Scale bars: 500 µm.

**Fig. 6** Chondrocytes from chondral flaps display low chondrogenic potential. These images show histologic staining of 5- $\mu$ m-thick cryosections of pellets after a 4-week culture period in vitro for glycosaminoglycan (GAG) (Alcian blue) and Types I and II collagen. (A) This image shows pellets made of flap chondrocytes at Passage 2, n = 5. (B) This shows pellets made of chondrocytes from non-delaminated knee articular cartilage samples—Knee AC at Passage 2, n = 3. Two negative controls (no primary antibody) are shown for immunostaining (bottom right). Sex and age (years) are indicated at the bottom left of all images. Arrows indicate areas where fainter or absent staining was observed, and dotted lines circle cell clusters. Scale bar: 100  $\mu$ m (close up) and 500  $\mu$ m (inset). (C) This image shows the immunohistochemistry of the negative control sample (no primary antibody). Scale bar: 100

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.

**Kommentiert [CO14]:** AU: Please put a space before and after the equal sign at each instance.

Author response: we added the spaces

**Kommentiert [CO15]:** AU: Please put a space before and after the equal sign in the p value.

Author response: this was done

 $\mu$ m (close up) and 500  $\mu$ m (inset). (D) This plot shows the measured area of the pellets, determined with an unpaired t-test with Welch's correction.

Supplemental Fig. 1 The chondrocytes' viability depends on the quantification technique. (A) This plot shows the viability results (dots are the average of the three technical replicates of each donor analyzed), comparing viability of the flaps using either live-dead or trypan blue exclusion assays (paired t test, \* p < 0.05). (B) This is a summary table of viability results for each sample (five biological replicates, three technical replicates for each).

**Kommentiert [CO16]:** AU: In the actual image, please use sentence case for capitalizing words (that is, only capitalize the first letter of each word.

Please put a space before and after the equal sign in the p value.

Please explain the asterisk in the A image in the legend.

For the column labeled "Age", please spell out "years" and put it in parentheses.

Please change "Gender" to "Sex", and include (M/F) in the column heading.

Please provide measurement labels for the columns "Viability live-dead" (note the addition of the hyphen; that is how you have written it in all other instances) and "Viability trypan blue"; are the measurements percentages? If so, include this in the column head: (%). It helps the reader to immediately understand the data you are presenting.

Author response: thank you for all the suggestions, all modifications were made

hat gelöscht: p = 0.03,

AU: Please do not delete query boxes or remove line numbers; ensure you address each query in the query box. You may modify text within selected text or outside the selected text (as appropriate) without deleting the query.