

Climate change impacts across a large forest enterprise in the Northern Pre-Alps: dynamic forest modelling as a tool for decision support

Journal Article**Author(s):**

Thrippleton, Timothy; Lüscher, Felix; Bugmann, Harald

Publication date:

2020-06

Permanent link:

<https://doi.org/10.3929/ethz-b-000416238>

Rights / license:

In Copyright - Non-Commercial Use Permitted

Originally published in:

European Journal of Forest Research 139(3), <https://doi.org/10.1007/s10342-020-01263-x>

1 **TITLE:**

2 Climate change impacts across a large forest enterprise in the Northern Pre-Alps - dynamic
3 forest modelling as a tool for decision support

4

5 Timothy Thrippleton^{1*} (timothy.thrippleton@usys.ethz.ch, +41 44 632 57 59)

6 ORCID IDs: 0000-0002-1017-7083

7 Felix Lüscher² (felix.luescher@oak-schwyz.ch)

8 Harald Bugmann¹ (harald.bugmann@env.ethz.ch)

9

10 **AFFILIATIONS AND POSTAL ADDRESSES:**

11 1: Swiss Federal Institute of Technology (ETH Zurich), Department of Environmental

12 Systems Science, Forest Ecology, Universitätstrasse 16, CH-8092 Zürich, Switzerland.

13 2: Oberallmeindkorporation Schwyz (OAK), Postfach 449, CH-6431 Schwyz, Switzerland.

14 *** Corresponding author**

15

16 **KEYWORDS:** Mountain forest, climate change impacts, Switzerland, dynamic vegetation
17 model, ungulate browsing.

18

19 **ACKNOWLEDGEMENTS:**

20 We thank Dominic Michel for support in all IT-related questions and the employees of the OAK

21 Schwyz for conducting the measurements of forest structure in the field. David I. Forrester and

22 the Experimental Forest Management (EFM) Project of the Swiss Federal Research Institute

23 WSL are gratefully acknowledged for providing data on the current stand structure of the site

24 FW (Riemenstalden). Two anonymous reviewers are gratefully acknowledged for their

25 thoughtful comments and helpful suggestions.

26

27 **ABSTRACT:**

28 Mountain forest managers face the challenge to anticipate climate change (CC) impacts across
29 large elevational ranges. For management planning, information on site-specific long-term
30 responses to CC as well as the consequences for protection functions is particularly crucial. We
31 used the process-based model ForClim to provide projections of forest development and their
32 protective function as decision support for a large forest enterprise in the Northern Pre-Alps.
33 Specifically, we investigated the impact of three climate scenarios (present climate, low- and
34 high-impact CC) at five representative sites along an elevational gradient (700 to 1450 m a.s.l.).
35 Relatively small changes to current forest structure and composition were evident under present
36 climate, but divergent trajectories occurred under CC: while the low-elevation sites (≤ 1000 m)
37 were affected by drought-related mortality, high-elevation sites benefited from the warming.
38 Changes at low-elevation sites were accompanied by shifts in species composition, favouring
39 in particular *Tilia* ('low impact' CC) and *Pinus sylvestris* ('high impact' CC). Forest
40 management accelerated the shift towards climate-adapted tree species, thereby reducing
41 detrimental effects of the 'low-impact' CC scenario. Under the 'high-impact' scenario,
42 however, drastic decreases in protective function occurred for the late 21st century at low
43 elevations. A set of exemplary disturbance scenarios (windthrow and bark beetle) demonstrated
44 the importance of forest management and browsing for post-disturbance resilience in mountain
45 forests.

46 Overall, our results underline the potential of process-based forest models as decision support
47 tools for forest enterprises, providing local projections of CC impacts across large elevational
48 ranges at the site-specific resolution required by forest managers.

49

50 **INTRODUCTION:**

51 The increasing impacts of climate change on forests worldwide (e.g., Allen et al. 2010) have
52 brought forest managers in the difficult position to balance multiple demands for forest
53 ecosystem services (ES, e.g., timber production, biodiversity, recreation) in the face of
54 uncertainties about the degree of future climate change (e.g., Yousefpour et al. 2017).
55 Maintaining the multitude of ES under climate change is particularly important for mountain
56 forests, which provide important ES for the mountain regions themselves (e.g., protection
57 functions against gravitational hazards, Elkin et al. 2013, Irauschek et al. 2017a) as well as
58 downstream (EEA 2010, Langner et al. 2017).

59 Climate change impacts on mountain forests can be highly heterogeneous in space, i.e. varying
60 in particular with elevation and local topography (Lindner et al. 2010). While low-elevation
61 valley bottoms are likely to become increasingly prone to drought-induced forest die-off (e.g.,
62 Bigler et al. 2006, Jump et al. 2006), higher elevations on the contrary tend to benefit from
63 longer vegetation periods and thus better growing conditions (Lindner et al. 2010, Bugmann et
64 al. 2014). However, high spatial heterogeneity (e.g., in terms of topography) and local site and
65 stand conditions (e.g., soil conditions, species composition) can superimpose these larger-scale
66 trends (Lindner et al. 2010) and potentially lead to complex, site-specific forest responses to
67 climate change (e.g., Etzold et al. 2019). Previous studies providing climate impact assessments
68 for mountain forest managers therefore emphasized the importance of local site and stand
69 conditions at a relatively fine scale and high resolution (Irauschek et al. 2017a, Klopčič et al.
70 2017, Mina et al. 2017b).

71 The rising awareness of the importance of mountain forests and their multiple ES has induced
72 an increasing research interest in the past years (Lexer and Bugmann 2017). Due the expected
73 extent of environmental changes (e.g., SCNAT 2016) and the long planning horizons in
74 mountain forests, dynamic vegetation models (DVMs) have become a central tool for

75 assessments of climate change impacts (e.g., Seidl et al. 2011a). Besides their suitability to
76 climate change applications, a further advantage of most DVMs is their ability to consider a
77 wide range of species (Pretzsch et al. 2008) and explore the benefits of species mixtures
78 (Forrester et al. 2017). Furthermore, various DVMs have been expanded over the last years to
79 represent a diverse array of management techniques (e.g., Rasche et al. 2011, Lafond et al.
80 2014, Irauschek et al. 2017b). Altogether, these developments have substantially improved the
81 suitability of DVMs to assess the effect of management and climate change on future forest
82 development and ecosystem service provisioning in a mountain context (e.g., Maroschek et al.
83 2015, Mina et al. 2017a).

84 In spite of the availability of assessments at larger (e.g., national) scales (e.g., Lexer et al. 2002,
85 Bircher et al. 2015), still only few studies are available for mountain forests at the local scale,
86 providing decision support at the level required by an individual forest enterprise (see e.g.,
87 Maroschek et al. 2015). Furthermore, findings from one case study area may apply only to a
88 limited degree to another area, particularly if environmental conditions are markedly different
89 (see e.g., Elkin et al. 2013). Mountain forest managers thus face the challenge to estimate how
90 studies from other regions or assessments at larger scales apply to their specific enterprise, and
91 they may therefore benefit significantly from DVM assessments that account for the specific
92 local conditions within their forest enterprise.

93 Initiated and co-developed by the local forest manager, we provide and evaluate the utility of a
94 DVM application for decision support in a large forest enterprise in the Northern Pre-Alps (the
95 Oberallmeindkorporation Schwyz, OAK-SZ) in Switzerland. The study area covers a large
96 environmental gradient from drought-affected sub-montane to currently temperature-limited
97 subalpine stands. As in many other mountain regions of Europe, a primary goal of forest
98 management in the OAK-SZ is to maintain their protection function against gravitational
99 hazards, in particular against rockfall and avalanches (see also Bebi et al. 2016). While some

100 stands may benefit from a warming under a moderate climate change scenario, a higher
101 frequency and intensity of extreme drought events may reverse this trend and cause abrupt
102 changes up to the point of complete forest dieback (Allen et al. 2015), with potentially fatal
103 consequences for the forests' protection function (Elkin et al. 2013, Bebi et al. 2016). An aspect
104 of particular importance for forest management is therefore whether stands are more likely to
105 change in a gradual or abrupt way, and at which time horizons these changes are to be expected
106 (Temperli et al. 2012). Besides the direct effects of altered temperature and precipitation, further
107 climate-related processes are likely to impact future forest development, most notably
108 overstorey disturbances (e.g., bark beetle outbreaks, Temperli et al. 2013, Seidl et al. 2017) as
109 well as intensified browsing pressure due to high ungulate populations (Côté et al. 2004,
110 Schulze et al. 2014).

111 The purpose of this study is therefore to address a set of stakeholder-defined management
112 questions at a high local resolution within the forest enterprise, which thus complements larger-
113 scale assessments (Bircher et al. 2015) and results from local-scale studies from other mountain
114 areas (Irauschek et al. 2017a, Mina et al. 2017a). Due to its particular location in an area of
115 steep environmental gradients from water-limited to temperature-limited forests, the study
116 furthermore provides an informative case for other managed mountain forests in the Northern
117 Alps, where contrasting climate change impacts can be expected to occur along elevational
118 gradients (cf. Lexer and Bugmann 2017).

119 Using the DVM ForClim (Bugmann 1996, Huber 2019), which has been developed for climate
120 change applications in mountain forests of the Central European Alps, we addressed the
121 following research questions, which were defined as the basis for long-term planning and future
122 decision making by the local forest manager:

123 (1) *Where* are the largest changes to be expected in terms of basal area and species
124 composition for two contrasting climate change scenarios ('low' and 'high impact')?

125 (2) *When* are changes in structure and composition to be expected? Are these changes
126 gradual or abrupt?

127 (3) *Which* effect will large-scale disturbances (windthrow, bark beetles) have at spruce-
128 dominated sites under a ‘high impact’ climate scenario? How is post-disturbance
129 recovery affected by previous forest management and the level of ungulate browsing?

130

131 **METHODS:**

132 *Forest model ForClim*

133 ForClim is a climate-sensitive dynamic vegetation model developed for short- and long-term
134 simulation of forest dynamics (Bugmann 1996). It belongs to the group of ‘forest gap models’,
135 which simulate forest properties emerging from individual-level interactions under the
136 influence of site-specific environmental conditions (e.g., temperature, precipitation,
137 topography, soil conditions, etc.). A forest is represented by multiple small patches, with the
138 patch size equivalent to the area dominated by a single large tree individual (Botkin et al. 1972)
139 based on the concept of patch dynamics (Watt 1947). Within each patch, tree demography is
140 simulated explicitly at annual time steps in the form of establishment, growth and mortality of
141 tree cohorts (i.e., groups of trees of the same species and age, Bugmann, 1996). Environmental
142 effects on tree growth are represented via growth-reduction factors, i.e. environmental
143 conditions that deviate from the optimum reduce species-specific growth (see also Bugmann,
144 2001).

145 ForClim has been applied across various temperate forests in Europe and other parts of the
146 world (e.g., Gutierrez et al., 2016; Mina et al., 2017; Huber 2019) and undergone thorough
147 evaluation under a wide range of species and site conditions (Rasche et al. 2012, Huber 2019).
148 Over the past decade, the capacities of ForClim to represent forest management have been
149 continuously expanded and evaluated in several mountain regions across Europe (Rasche et al.

150 2011, Mina et al. 2017a). All simulations of this study were conducted with ForClim Version
151 4.0.1 (Huber 2019); see Online Resource 1 for further details.

152

153 *Study areas*

154 Within the planning unit of the ‘Oberallmeindkorporation Schwyz’ (subsequently abbreviated
155 as OAK-SZ), five sites were selected to represent (1) the most frequent forest communities
156 (after Ellenberg and Klötzli 1972) per elevation zone (see Table 1) which were characteristic
157 for the forest enterprise, and (2) stands in the timber stage of development (i.e. dominant DBH
158 > 30 cm), which had highest priority from a forest management perspective. The study region
159 is located in the Northern Alps and features a pronounced precipitation gradient over a relatively
160 short distance, ranging from moderate annual precipitation amounts of ca. 1100 mm at the
161 southeastern part of Lake Lucerne to >2000 mm in the valleys in its east and northeast (HADES
162 2015). Furthermore, soil conditions differ considerably among the sites due to distinct
163 differences in climate, geology, vegetation and topography (particularly slope angle). While the
164 study sites Brünischart (BS) and Fronwald (FW) are characterized by relatively low soil water
165 holding capacity (SWHC) due to steep slopes and shallow soil depth on calcareous bedrock
166 (Hantke and Kuriger 2003), deeper soils with a higher SWHC occur at higher elevations, i.e.
167 the study site of Herrenwald (HW), Tröliger Wald (TW) and Schwarz Stock (SSt, see Table 1).
168 In terms of species composition, the lowest elevation site BS is characterized by a broadleaf-
169 dominated forest, while the higher elevation sites are dominated by coniferous species
170 (particularly *Abies alba* and *Picea abies*, see Table 1 for details). The stand structure of each
171 site was measured within 1 ha plots during July to September 2018, with the exception of FW
172 (Riemenstalden site 01-053.001, Schwitter 2006), where data were provided by the
173 Experimental Forest Management project under the lead of David I. Forrester (Forrester et al.
174 2019). Details about the stand initialisation are provided in Online Resource 1.




Figure 1 Location of study sites within the management area of the forest enterprise ‘Oberallmeindkorporation Schwyz’ (OAK-SZ), Switzerland. © Swisstopo.

Table 1 Elevation zones (following the definition of Frehner et al. 2005), environmental conditions (MAT: mean annual temperature; AP: annual precipitation sum), dominant tree species and respective forest community (after Ellenberg and Klötzli 1972) for the study sites (as described in ATRAGENE 1997, 1999, 2004). Species abbreviations: QuPetr: *Quercus petraea*, PiSylv: *Pinus sylvestris*, AcPseu: *Acer pseudoplatanus*, AbAlba: *Abies alba*, PiAbie: *Picea abies*.

Site	Elevation (m. a.s.l.)	Elev. Zone	MAT (°C)	AP (mm)	Slope (°)	Aspect	Soil type	Soil depth (cm)	Dom. tree species	Forest community (Ellenberg & Klötzli, 1972)
Brünischart (BS)	700	sub-montane	8.63	1178	>30	SE	Rendzina	40-60	<i>QuPetr.</i> , <i>PiSylv.</i> , <i>AcPseu</i>	40C
Fronwald (FW)	1000	lower- montane	6.11	1360	>30	SW	Rendzina	50-60	<i>AbAlba</i> , <i>PiAbie</i>	12
Herrenwald (HW)	1200	upper- montane	5.02	2000	10-30	S	Brown soil	100	<i>PiAbie</i> , <i>AbAlba</i>	19
Tröliger Wald (TW)	1350	high- montane	4.67	2010	10-30	N	Brown soil	80	<i>PiAbie</i>	46D
Schwarz Stock (SSt)	1450	sub-alpine	4.42	2033	>30	N	Podzol	70	<i>PiAbie</i>	57S

180 *Small-scale 'mountain forest plentering' silviculture*

181 Forest management in mountain regions is particularly difficult, since management options
182 become increasingly constrained towards higher elevations and promoting sufficient
183 regeneration is a considerable challenge (Schwitter, 2013). The management technique of
184 'mountain forest plentering' (MFP) silviculture has been developed to cope with the specific
185 situation in mountain forests, in particular to induce regeneration (Schwitter 2013, Leuch et al.
186 2017). MFP represents a small-scale removal of collectives of trees (rather than individual trees
187 as done in lowland plenter forests, Schwitter 2013) with the objective to foster regeneration via
188 improved light and temperature conditions at the forest floor (Leuch et al. 2017).

189 A new MFP management module was therefore designed for ForClim to harvest tree collectives
190 in small patches of 400 m² (i.e. the ForClim patch size in this study) within the forest. Although
191 not being spatially explicit, this routine mimics the approach underlying cable yarding, where
192 the goal is to remove a pre-defined fraction of the standing volume per stand. The harvest
193 intensity is therefore defined as the target timber volume (Vol_{Target}), representing the fraction of
194 volume within the entire stand (i.e., all patches) to be harvested per intervention. Besides, a
195 species-specific target diameter (DBH_{Target}) has to be defined by the user, representing the
196 diameter threshold above which trees can be harvested. Furthermore, the time interval between
197 the management interventions has to be defined by the user. The MFP module assures that
198 scheduled harvest interventions are only carried out if sufficient harvestable volume (i.e. the
199 prescribed Vol_{Target}) is available in the entire stand (i.e. all patches) to mimic the situation that
200 expensive mountain forest harvest interventions are only conducted if the intervention is cost-
201 effective, i.e. sufficient timber can be harvested. If sufficient harvestable timber is available for
202 the intervention, the module progressively harvests all trees within randomly chosen patches
203 above the user-defined DBH_{Target} , thus creating gaps with favourable light conditions for
204 regeneration while not removing the regeneration that is present already (if any; so-called

205 “advance regeneration”). Notably, the module assures that the same patch is only harvested
206 once per rotation cycle, i.e. a patch is not harvested in two consecutive interventions as long as
207 other unharvested patches are available. Further details about the MFP module can be found in
208 Online Resource 1.

209 All MFP-management interventions were scheduled at 20-year intervals for all sites, starting
210 from 2019. The management prescriptions differed between the lower-elevation sites (BS and
211 FW) and the higher elevation sites (HW, TW and SSt) accounting for the different conditions
212 along the elevation gradient. For the lower-elevation sites, all interventions were carried out
213 with the same management intensity (Vol_{Target} of 15%) for all harvest interventions, applying a
214 DBH_{Target} of 40 cm for *Picea abies* and 60 cm for all other species. For the higher-elevation
215 sites, a higher intensity intervention was carried out in the first harvest year 2019 (Vol_{Target} of
216 25%), followed by lower intensity harvests (Vol_{Target} of 15%) in the subsequent interventions. At
217 the high elevations, a DBH_{Target} of 12 cm was defined for all species.

218

219 *Calculation of ecosystem services*

220 Changes in aboveground biomass, species composition and forest structure were measured in
221 terms of basal area ($m^2 ha^{-1}$), protection against gravitational hazards was calculated as a
222 dimensionless protection indices, i.e. avalanche protection index (API) and rockfall protection
223 index (RFPI), developed by Elkin et al. (2013) and Schmid (2014) based on Frehner et al.
224 (2005). The API is derived from an interception component (calculated from stand leaf area
225 index, LAI, and the relative share of coniferous trees in the stand) and a stability component
226 associated to stem density (based on the number of trees with a $DBH > 8$ cm). The resulting
227 index varies between 0 and 100, with an API of 100 representing maximum protection. The
228 RFPI is based on stem density and diameter distribution, which determines the capacity of the
229 stand to protect against rocks of certain sizes. For the present study, a collective index was

230 calculated for all rock size classes, as described in Schmid (2014). As for the avalanche
231 protection index, the RFPI index varies between 0 (no protection) and 100 (max. protection
232 function). Further details and formulae are given in Online Resource 1.

233

234 *Simulation scenarios*

235 Simulations of future forest dynamics were conducted for present climate and two contrasting
236 climate change scenarios that represented a ‘low impact’ scenario (RCP3PD, compliant to the
237 targets of the Paris Agreement) and a ‘high impact’ scenario (A2, i.e. unabated emissions),
238 based on the CH2011 report for Switzerland (CH2011). For present climate conditions, the time
239 series from 1931 – 2017 from the WSL database of spatially interpolated climate data derived
240 using DAYMET (Thornton et al. 1997) was applied (cf. Online Resource 1). The ‘low impact’
241 scenario features a moderate increase in annual mean temperature by +1.5 °C and decrease of
242 annual precipitation sum by -10% until the end of the 21st century (relative to the baseline period
243 of 1980 to 2011), while the ‘high impact’ scenario represents a temperature increase by +4 °C
244 and precipitation decrease of -25% (CH2011). Further details about the climate change
245 scenarios are provided in Online Resource 1.

246 Furthermore, two contrasting scenarios for ungulate browsing pressure (‘low browsing’ and
247 ‘high browsing’) were considered in the simulation setup, since browsing damage to tree
248 regeneration is a considerable problem in many mountain forests (Kupferschmid et al. 2015).
249 Browsing in ForClim affects the density and species composition of tree regeneration,
250 depending on browsing pressure and a species-specific browsing tolerance (Didion et al. 2011,
251 see also Online Resource 1, section 'Browsing scenarios' for details).

252 To assess the importance of management on forest dynamics under climate change, two
253 scenarios were compared: one without any management interventions and one including the

254 MFP management (cf. above and Online Resource 1 for details). All simulations were run for
255 150 years, as in Mina et al. (2017b). Since close-to-nature forestry is an important management
256 guideline in many parts of Europe (Bauhus et al. 2013), simulations determining the potential
257 natural vegetation (PNV) at each site were conducted to provide additional information on this
258 ‘natural reference state’ for forest management. These simulations were run for 1000 years
259 under all climate scenarios, in the absence of management, and under a low browsing pressure.
260 In addition to the direct effects of temperature and precipitation changes, climate change is also
261 likely to lead to intensified disturbance regimes (Seidl et al. 2017). Since predicting the exact
262 timing of disturbance occurrence is practically impossible (Mina et al. 2017b), we investigated
263 forest dynamics following a set of exemplary large-scale disturbances by windthrow and bark-
264 beetle outbreaks, two key disturbance types in Central European forests (Seidl et al. 2011b).
265 These simulations were carried out for the site TW (1350 m), a typical mature, spruce-
266 dominated stand that is structurally prone to both windthrow and bark-beetle disturbances. To
267 evaluate the effects of the timing of disturbance occurrence under a strong warming scenario,
268 exemplary disturbance scenarios were simulated under the ‘high impact’ climate change
269 scenario for the years 2030 and 2070 with and without pre-disturbance management (cf. Online
270 Resource 1 for detailed information).

271 All results refer to forest structure and composition at the end of the simulation period (year
272 2150), unless stated otherwise. The simulations were carried out with ForClim version 4.0.1
273 (Huber 2019), all analyses and visualisations were carried out with R Version 3.4.3 (R
274 Development Core Team 2017).


275

276 **RESULTS:**

277 *Elevation-specific magnitude of changes in structure and composition*

278 For the ‘low impact’ climate scenario, only small changes in basal area, but noticeable shifts in
279 species composition occurred across the five sites until 2150 (Fig. 2). Species shifts were most
280 pronounced at the sites ≤ 1000 m a.s.l. (i.e., BS and FW), where warmth-adapted broadleaved
281 species (particularly *Fagus sylvatica* and *Tilia cordata*) increased in abundance. Higher
282 elevation sites experienced an increase in *Fagus sylvatica*, partly at the expense of *Picea abies*
283 (e.g., site SSt). The PNV simulations for BS and FW (Fig. 2, top row) showed that these trends
284 indicate a long-term shift in species dominance.

285 Simulations including forest management (Fig. 2, bottom row) showed a higher share of early
286 successional species (e.g., *Betula pendula*) as well as a shift towards more thermophilic
287 broadleaved species in the regeneration layer (e.g., *Fagus* and *Acer*, Online Resource 3, Fig.
288 A3.14ff.), representing an earlier transition towards a species composition in equilibrium with
289 climate (see PNV results). However, this transition in the regeneration layer was inhibited by
290 high browsing pressure (Fig. 2, ‘high browsing pressure’; cf. Online Resource 3), which
291 reduced basal area and substantially reduced regeneration density (Online Resource A3, e.g.,
292 Fig. A3.15).




293

Figure 2 Stand basal area under the ‘Low impact’ climate scenario (RCP3PD) at the five study sites for unmanaged (top row) and managed (bottom row) conditions. ‘Present state’ refers to measured initial state at year 2018, ‘Future state’ (with ‘low’ and ‘high’ browsing pressure) refers to stand state at simulation year 2150, and ‘PNV’ refers to potential natural vegetation establishing after a simulation time of 1000 years under novel climatic conditions (only for unmanaged conditions).

294

295 For the ‘high impact’ climate scenario, drastic reductions in basal area occurred for the lower
 296 elevation sites BS and FW, while the other sites located at higher elevations increased in basal
 297 area until the year 2150 (Fig. 3, top row). For BS, the previously broadleaf-dominated stand
 298 experienced a complete transition to a *Pinus sylvestris*-dominated woodland, which featured
 299 low basal area ($< 15 \text{ m}^2 \text{ ha}^{-1}$) and was characterized by low densities of large trees ($>30 \text{ cm}$
 300 DBH), particularly under ‘high browsing’ conditions (Online Resource 3, Fig. A3.3 and Fig.
 301 A3.5). In terms of species composition, all other sites showed increasing shares of broadleaved
 302 trees (particularly *Tilia* for the site FW and *Fagus* at the other sites). Simulations including
 303 forest management showed higher shares of these species in the regeneration layer (Fig. 3 and
 304 Online Resource 3, Fig. A3.9ff.) and, by the year 2150, were closer to the PNV composition
 305 (Fig. 3).




306

Figure 3 Stand basal area under the ‘High impact’ climate scenario (A2) at the five study sites for unmanaged (top row) and managed (bottom row) conditions. ‘Present state’ refers to measured initial state at year 2018, ‘Future state’ (with ‘low’ and ‘high’ browsing pressure) refers to stand state at simulation year 2150, and ‘PNV’ refers to potential natural vegetation establishing after a simulation time of 1000 years under novel climatic conditions (only for unmanaged conditions).

307

308 These climate-induced changes affected the basal area and protection function of all sites,
 309 although the magnitudes of change differed between the two climate scenarios (Fig. 4, Online
 310 Resource 2, Fig. A2.4-2.6). For the ‘low impact’ climate scenario, only slight decreases in basal
 311 area resulted for the low-elevation sites (BS and FW), and higher basal area as well as rockfall
 312 and avalanche protection for the high-elevation sites (Fig.4). These patterns were much more
 313 pronounced for the ‘high impact’ scenario, where basal area and protection functions decreased
 314 dramatically for the low-elevation sites. At the high-elevation sites, basal area as well as rockfall
 315 and avalanche protection increased. In general, similar patterns emerged under either browsing
 316 scenarios, although the magnitude of changes was higher in the ‘high browsing pressure’
 317 scenario, particularly for avalanche protection (Fig. 4 and Online Resource 2, Fig. A2.4-2.6).



318


Figure 4 Changes in basal area, rockfall protection (RFP) and avalanche protection (AP) index for all sites at year 2150 under the ‘low impact’ (RCP3PD) and ‘high impact’ (A2) scenario relative to present climate. Colours indicate the direction of change (red: decrease, blue: increase). Note that results shown here refer to the ‘with management’ and ‘high browsing’ simulations. Results for the ‘no management’ and ‘low browsing’ simulations feature similar patterns and are provided in Online Resource 2.

319

320 *Temporal patterns of change*

321 Temporal patterns of change in basal area differed substantially between the five sites and
 322 among the two climate change scenarios and were consistent for unmanaged (Fig.5) and
 323 managed stands (Online Resource 3, Fig. A3.4ff.). While the ‘low impact’ climate scenario
 324 showed only gradual changes until the year 2150, abrupt changes occurred under the ‘high
 325 impact’ climate scenario for the low-elevation sites BS and FW. Furthermore, the onset of a
 326 decreasing trend for BS and FW differed between the climate scenarios: for the ‘low impact’
 327 scenario, basal area remained nearly constant until 2080, whereas it started to decrease already

328 around 2050 for the ‘high impact’ scenario (Fig. 5). Moreover, abrupt changes in basal area
329 under the ‘high impact’ scenario occurred at different time points depending on the site. While
330 BS experienced an abrupt dieback already by 2060-2080, the site FW experienced a strong
331 drought-related decrease only towards the end of the 21st century, and to a smaller extent (Fig.5
332 and Online Resource 2, Table A2.1, Fig. A2.9).




333

Figure 5 Development of basal area over time under the ‘low impact’ (RCP3PD) and ‘high impact’ (A2) scenario, shown as the difference to the respective trajectory under present climatic conditions for the five study sites under unmanaged conditions.

334

335 *Disturbance effects*

336 Both exemplary disturbance simulations for windthrow and bark beetle at the spruce-dominated
337 site TW (1350 m) under a ‘high impact’ climate scenario had a similarly drastic effect, reducing
338 stand basal area to <10 m² per ha (Fig. 6 and Online Resource 3, Fig. A3.26ff.). The reduction
339 of stand basal area was higher when the disturbances occurred earlier (i.e. in 2030 compared to
340 2080), which was due to a lower abundance of advance regeneration prior to year 2030 (Online
341 Resource 3, Fig. A3.26 and Fig. A3.30). Furthermore, tree regeneration and growth benefited
342 more from the warmer conditions prevailing in 2080 compared to 2030, leading to a faster
343 recovery of basal area (Fig. 6). The recovery was however substantially impeded by high
344 browsing pressure (Fig. 6).




345

Figure 6 Post-disturbance recovery of basal area for the exemplary barkbeetle and windthrow disturbance scenarios at the site TW (1350 m). Different colors indicate the different time point of disturbance occurrence (2030 or 2080). Different line type indicates the browsing intensity. Simulations assumed a ‘high impact’ (A2) climate change scenario without pre-disturbance management.

346

347 Regarding the effect of pre-disturbance forest management, basal area was up to 30% higher at
348 the end of the first decade after disturbance if the stand had previously been managed by
349 mountain forest plentering than if no management had been conducted (Fig. 7 and Online
350 Resource 2, Fig. A2.7). This was due to a higher abundance of advance regeneration in the

351 managed stands, which contributed substantially to post-disturbance recovery (Online Resource
352 A3, cf. Fig. A3.26 and Fig. A3.28). The legacy effect of previous management decreased with
353 time since disturbance, but high browsing pressure continued to shape the trajectory of long-
354 term recovery (Fig. 6 and Online Resource 2, Fig. A2.8).



355

Figure 7 Basal area at the end of the first decade after disturbance in 2080 for the two exemplary disturbance scenarios (windthrow and bark beetle) under low and high browsing, as well as managed (“WithMng”) and unmanaged (“NoMng”) conditions at the site TW (1350 m). Results for the disturbance in year 2030 are shown in Fig. A2.5 (Online Resource A2).

356

357 **DISCUSSION**

358 Our results showed contrasting responses to climate change along the elevational gradient
359 within the forest enterprise, with negative impacts likely to occur at the lower elevation sites
360 (≤ 1000 m a.s.l.), whereas higher elevation sites are more likely to benefit from warmer
361 conditions in the absence of disturbances. In contrast to the ‘low impact’ climate scenario,
362 abrupt negative impacts occurred for low-elevation sites under the ‘high impact’ scenario from
363 the mid-21st century onwards, leading to substantial shifts in species composition and stand
364 structure as well as a severe loss of protection function. Following our research questions, we
365 first discuss (1) the elevation-specific magnitude of change and (2) their temporal dynamics,
366 including the response to the exemplary disturbance scenarios. Ultimately, we discuss (3) the
367 limitations of this study and further capabilities of DVMs to provide decision support for forest
368 managers.

369

370 *Elevation-specific magnitude of change*

371 Elevation and topography are key factors altering the local climate in mountain landscapes due
372 to, e.g., the temperature gradient with elevation, orographic rainfall and varying incident solar
373 irradiation with aspect and slope (Whiteman 2000, Zou et al. 2007). The five stands of our study
374 were located across a gradient from drought-prone, steep south-facing sites at lower elevation
375 to moist, north-facing sites at high elevation. These contrasting conditions were reflected in the
376 differential climate change impacts, with detrimental effects predominating at the two low-
377 elevation sites with southerly aspect (BS and FW). Empirical studies from dry inner-alpine
378 valleys in Central Europe show similar patterns under contemporary climate change, i.e.
379 significantly higher drought-induced mortality at elevations < 1000 m (Minerbi et al. 2006,
380 Rigling et al. 2013) and decreased regeneration densities on low-elevation south-facing slopes
381 (e.g., Wohlgemuth and Moser 2018). However, the comparison of the simulated climate change

382 impacts for the site BS with the results from dry inner alpine valleys is restricted by the
383 distinctly different climatic conditions between the respective regions (e.g., Elkin et al. 2013).
384 At the site BS, high precipitation amounts during summer prevent severe droughts under present
385 climate conditions (see Online Resource 2, Fig.A2.9) and permit the growth of a mixed
386 deciduous forest with relatively high basal area in spite of the low soil water holding capacity
387 (ATRAGENE 1997). Nevertheless, our simulation results indicate that a number of severe
388 droughts under future climate change may change the situation rapidly, and emphasize the
389 particular vulnerability of the low-elevation, south-facing site BS among the gradient of
390 simulated sites within the forest enterprise.

391 This site-specific response exemplifies the challenge that mountain forest managers are facing
392 and shows how DVMs can provide decision support for forest planning with a local resolution.
393 On the one hand, large-scale assessments provide important information on climate change
394 impacts (e.g., Bugmann et al. 2014, Bircher et al. 2015, SCNAT 2016), which may however
395 not be at a sufficiently detailed level for mountain forest enterprises located in topographically
396 complex and environmentally contrasting settings. As shown by the recent empirical study by
397 Etzold et al. (2019), patterns of climate-induced forest mortality in Switzerland are highly
398 complex and depend on the combination of species effects and small-scale site conditions. An
399 evaluation based on species composition alone may for instance conclude that the stand BS
400 (dominated by drought-tolerant oak and pine) would be more resistant to drought impacts than
401 FW (dominated by the less drought-tolerant *Abies alba* and *Picea abies*, e.g., Leuschner and
402 Meier 2018). While the ‘low impact’ climate scenario indeed caused little change in forest
403 structure and composition at BS, the ‘high impact’ climate scenario induced drastic diebacks at
404 this site, despite the predominance of drought-tolerant species. In contrast, the higher elevation
405 site FW was less impacted despite the predominance of more drought-sensitive conifers. Our
406 study thus supports the findings of context-specific climate change impacts, which depend

407 strongly on the specific abiotic and biotic conditions (Condes and del Rio 2015, Sanchez-
408 Salguero et al. 2015, Etzold et al. 2019).

409 The simulated ‘high impact’ climate change effects at low elevations have pronounced
410 consequences for the services provided by the respective stands, since the decrease of basal area
411 implies a loss of harvestable timber as well as a severe loss of protection function against
412 gravitational hazards. In the absence of a continuous forest cover, avalanche release risk is
413 substantially higher since the forest cover decreases snowpack depth (due to higher snow
414 interception), alters microclimatic conditions and increases surface roughness (Frehner et al.
415 2005). Similarly, rockfall risk increases drastically as stands formerly characterized by a large
416 range of size and densities lose their well-structured characteristics (Dorren et al. 2005).
417 According to Bebi et al. (2016), the importance of rockfall (as well as landslides and erosion)
418 is expected to increase more than the importance of avalanches under climate change. Although
419 the seasonal time window with critical snowcover for avalanches is likely to decrease with
420 climate change, avalanche risks at lower elevations should not be underestimated, particularly
421 regarding snow-gliding on steep south-facing slopes (Bebi et al. 2016), e.g. at the site BS. In
422 the case of rockfall, an adaption towards a coppice management could however diminish this
423 increasing risk and improve the protective function of the lower elevation, broadleaf-dominated
424 forests (e.g., Radtke et al. 2014).

425 An additional level of decision support that can be provided by a DVM-based approach is the
426 local-scale composition of potential natural vegetation (PNV), which represents the emerging
427 species compositions under novel climate conditions. Although the drastic dieback at the site
428 BS was projected to lead to very low basal areas for the end of the 21st century, the PNV
429 simulations imply that a *Pinus sylvestris* forest would re-establish in the long-term even under
430 a ‘high impact’ climate scenario. It is however possible that besides *Pinus sylvestris*, more
431 drought tolerant sub-Mediterranean tree species could immigrate at the low-elevation BS site

432 in the future, such as pubescent oak (*Quercus pubescens*), which is becoming an increasingly
433 important tree species in dry central Alpine valleys (Rigling et al. 2013). As recently
434 demonstrated by Huber (2019), the immigration of more climate-adapted species could play an
435 important role in buffering negative impacts of climate change. Altogether, the combination of
436 insights from empirical studies (e.g., Frank et al. 2017, Wohlgemuth et al. 2018), as well as
437 from DVM-based assessments can thus provide important decision support regarding which
438 tree species to favour at a specific location in the long-term.

439 In contrast to the detrimental impacts at low elevations, the ‘high impact’ climate scenario led
440 to increases in basal area and a shift in species composition towards more thermophilic
441 broadleaved species at the high-elevation sites (> 1000 m a.s.l.). Although the increase in basal
442 area suggests a positive development for forest management (i.e., provisioning of more
443 harvestable timber), simulated species composition showed a strong discrepancy compared to
444 the PNV simulations by the year 2150, indicating that the high fraction of Norway spruce (*Picea*
445 *abies*) remaining in the simulated stands is unlikely to represent a climatically suitable forest
446 composition (cf. Bugmann et al. 2014). Particularly under unmanaged conditions, stand
447 structure in the year 2150 was characterized by a high number of tall, old spruce trees, indicating
448 a high susceptibility to windthrow (Seidl et al. 2014, Schuler et al. 2019). Furthermore, climate
449 change likely leads to an increase in the risk of bark beetle outbreaks at higher elevations due
450 to an increase in the number of generations per year and prolonged annual flight periods
451 (Temperli et al. 2013, Bugmann et al. 2014). The advantages of increased growth may thus be
452 counteracted by negative impacts of enhanced biotic or abiotic disturbances, as shown in our
453 exemplary disturbance simulations. Including forest management in our simulations however
454 induced the regeneration of more climate-adapted species (particularly *Fagus* and *Acer*) and led
455 to a faster transition towards a less vulnerable mixed species forest at higher elevations.

456 With respect to species responses to climate change along the elevation gradient, our simulation
457 results showed an upslope spread of thermophilic, broadleaved species, particularly under the
458 ‘high impact’ climate scenario. This is in line with an increasing number of observations under
459 contemporary climate change that are reporting an upward expansion of the distribution ranges
460 of trees (e.g., Penuelas and Boada 2003, Vitasse et al. 2012), although species-specific
461 responses can be complex and site-specific (e.g., Gazda et al. 2019). Furthermore,
462 palaeoecological studies show similar upward shifts of broadleaved species during periods with
463 higher temperatures, e.g. for *Tilia* during the Holocene temperature maximum (Thöle et al.
464 2016), and suggest an increase in the abundance of *Fagus sylvatica* at high elevations under
465 future climate change (Schwörer et al. 2014). The projected shifts of dominant species are thus
466 in general agreement with patterns found in empirical studies of past and contemporary climate
467 change, although the responses of some subdominant species (e.g., increase of *Betula pendula*
468 at high-elevation managed sites) may be overestimated (cf., e.g., Wohlgemuth and Moser
469 2018).

470

471 *Temporal patterns of change*

472 Estimating the most likely pathway of future climate change is difficult due to uncertainties in
473 the future of socio-economic and technological developments (IPCC 2013). While trajectories
474 of temperature changes show relatively small differences between the emission scenarios in the
475 first half of the 21st century, they differ substantially until the end of the century for Switzerland
476 (CH2011). A similar trend was evident from the simulated forest properties, with little change
477 until around 2040, but substantial differences between the climate change scenarios thereafter.
478 The rates of change are of particular importance in this context (cf., Reyer et al. 2015), tending
479 to be gradual for the ‘low impact’ scenario, but more abrupt for the ‘high impact’ scenario at
480 sites ≤ 1000 m a.s.l. While the specific time points of these drastic changes are subject to

481 considerable uncertainty, several studies suggest a much higher probability and frequency of
482 drought events in the twenty-first century (e.g., Dai 2013). The combination of higher
483 temperatures and lower precipitation is likely to lead to tipping point dynamics, resulting in
484 sudden drought-related tree mortality (Allen et al. 2015). The risk of this sudden dieback is
485 particularly high for tree species currently growing under unsuitable climatic conditions, as
486 shown e.g. for *Picea abies* at low elevations (e.g., Levesque et al. 2013). In accordance with
487 these findings, our results suggest that a ‘high impact’ climate change scenario leads to a much
488 higher risk of rapid drought-related forest dieback for the late 21st century.

489 If the current trend of failing to meet the targets of the Paris agreement continues (UNEP 2019),
490 low- to mid-elevation mountain forests may be at particular risk of drought-related tree
491 mortality and substantial losses in their protective function against gravitational hazards. To
492 avoid or at least mitigate this situation, forest management should particularly promote the
493 regeneration with climate-adapted species (SCNAT 2016). Our results demonstrate how a
494 DVM approach can provide decision support in the selection of potentially suitable species and
495 help to estimate the timespan remaining for forest management to address these issues before
496 negative changes are becoming evident in the forest. Both aspects, i.e. identifying suitable
497 species and estimating available time windows for taking action, are key for enterprise-level
498 forest planning (Streit et al. 2017). Based on the results from BS, considerable impacts of
499 climate change are likely to be expected in our case-study enterprise from around the mid-21st
500 century under a ‘high impact’ climate scenario, suggesting that forest management has less than
501 three decades to establish sufficient advance regeneration, which is a critically short time
502 window in mountain forests (Ott et al. 1997).

503 Besides the direct impacts of climate change on low elevation sites, abiotic and biotic
504 disturbances by windthrow and bark-beetle attacks pose a threat to the spruce forests at higher
505 elevations (Seidl et al. 2011b, Temperli et al. 2013). As demonstrated by our exemplary

506 disturbance simulations, the benefit of timely forest management and reduced browsing
507 pressure also plays a key role for post-disturbance recovery in high elevation forests. Notably,
508 a higher post-disturbance basal area and a faster recovery was simulated when the disturbance
509 (windthrow or bark beetle) occurred in the late 21st century (year 2080), which was due to a
510 higher abundance of advance regeneration induced by management and the additional effect of
511 the warmer climate. These findings are supported by empirical studies, e.g. by Schwitter et al.
512 (2015), who evaluated 24 years of forest dynamics at windthrow sites in Switzerland and found
513 that post-disturbance recovery was much faster when advance regeneration was present in the
514 pre-disturbance stands. This initial advantage in the first years after disturbance is of particular
515 importance for the protection function of a forest, since trees of a certain minimum height are
516 required to provide protection against avalanches and rockfall (Noack et al. 2004). Furthermore,
517 browsing pressure is a key factor that can reduce establishment and slow down post-disturbance
518 forest succession substantially, as shown by various empirical (e.g., Kupferschmid and
519 Bugmann 2005) and modelling studies (e.g., Kupferschmid et al. 2006, Thrippleton et al. 2018).
520 It has to be noted, however, that rather than providing a realistic projection of future forest
521 dynamics under changing disturbance regimes, our exemplary disturbance simulations only
522 aimed to test the broad effects of timing of disturbance, previous management and browsing
523 pressure on recovery. Despite these limitations, our results demonstrate at a general level that
524 reduced browsing pressure and adequate forest management are key to improve forest resilience
525 to disturbances by windthrow and bark beetle attacks in susceptible spruce-dominated mountain
526 forests.

527

528 *Using DVMs as decision support tools for mountain forest managers*

529 Our study exemplifies how a DVM can provide key insights into climate change impacts at the
530 scale of interest of a forest enterprise. For the present study, a relatively simple approach was

531 applied, which requires only moderate efforts in terms of data acquisition from the forest
532 enterprise (i.e. full calipering of a handful of representative stands). However, DVMs can
533 provide a wealth of further decision support under changing environmental conditions (see also
534 review by Fontes et al. 2010) from lowland to mountain forests (e.g., Irauschek et al. 2017a,
535 Lexer and Bugmann 2017, Gutsch et al. 2018).

536 With respect to spatial scale, measurements of current stand structure can be obtained at high
537 resolution and large spatial extents by combining inventory data and airborne laser scanning
538 (LiDAR), as shown by Maroschek et al. (2015). Rather than using the spatially limited approach
539 of representative stands, as in our study, some DVMs are capable of representing the spatial
540 arrangement of forest patches at much larger spatial scales, e.g. the models LANDIS
541 (Mladenoff 2004), iLand (Seidl et al. 2012) or LandClim (Schumacher et al. 2004). This type
542 of landscape models furthermore allows to simulate exogenous disturbances as self-emergent
543 properties of the system (e.g., Seidl et al. 2014, Temperli et al. 2015), which was not possible
544 in our stand-scale DVM. However, the increasing degree of spatial detail comes with its own
545 challenges, e.g. a strongly increasing data demand (see also review by Keane et al. 2015), which
546 reduces the likelihood of such models being used in the absence of a dedicated and specifically
547 funded research project.

548 Another aspect that was not investigated in detail in this study is the use of DVMs to explore
549 alternative forest management approaches to optimize the provisioning of multiple goods and
550 services. The capacity of DVMs to assess trade-offs and synergies between multiple ecosystem
551 services has been addressed more recently, e.g. by studies in different mountain areas within
552 Europe by Mina et al. (2017a) and Langner et al. (2017). An overview of the management
553 capacities of different DVMs is given in Fontes et al. (2010), although the capabilities of some
554 DVMs have increased considerably in the last years (e.g., Irauschek et al. 2017b, Mina et al.
555 2017b). Ultimately, multi-criteria decision analyses can be used in combination with DVM

556 simulations to facilitate the identification of appropriate management alternatives for a wide
557 range of forest goods and services (e.g., Wolfslehner and Seidl 2010).

558 Overall, DVMs are increasingly suitable tools for providing decision support in mountain
559 forests. A broader coverage of studies from different mountain regions would thus be highly
560 valuable from a forest management perspective as well as from a scientific perspective, e.g. for
561 synthesizing overarching conclusions on the impact of climate change in mountain regions
562 (e.g., Price et al. 2011).

563

564 **CONCLUSION**

565 Although climate change impact assessments are increasingly becoming available at larger
566 (e.g., national) scales, only a few studies have focused on the climate change impacts on
567 mountain forests at the level of a forest enterprise, taking into account the combined effect of
568 climate, specific site conditions, initial forest structure, and management. Dynamic forest
569 models are capable of accounting for these aspects and thus are promising tools to provide
570 assessments at a relatively high local resolution, which is required by mountain forest managers.

571 The present study demonstrates how a DVM approach can provide information on site-specific
572 forest vulnerability, assessments of the timing and magnitude of change, quantification of
573 changes in protective function as well as information regarding the choice of climate-adapted
574 tree species; all these aspects are highly relevant for long-term planning within the forest
575 enterprise. For our case study forest enterprise, key conclusions were: (1) low elevation sites
576 (≤ 1000 m a.s.l.) were most vulnerable to adverse climate change impacts and should thus
577 receive particular attention. (2) Forest management was of high importance to induce sufficient
578 advance regeneration, which is key to ensure forest resilience and specifically the protective
579 function in the long term. (3) Management measures must be taken in the near future (i.e., the

580 coming 20 to 30 years) to avoid a severe loss of the protective function particularly under a
581 ‘high impact’ climate scenario. Although higher elevation sites benefited from the warming
582 climate, simulations for potential natural vegetation indicate that the current vegetation
583 composition becomes climatically unsuitable, thus suggesting that risks of catastrophic
584 disturbances (e.g., windthrow and barkbeetle) may rise accordingly.

585 Ultimately, the results from this study emphasize that international efforts to reach the goals of
586 the Paris Agreement are of crucial importance for mountain forests, as unabated climate change
587 is likely to severely deteriorate forest health as well as the services that are provided by these
588 forests.

589

590 **ELECTRONIC SUPPLEMENTARY MATERIALS:**

591 ESM_1: Extended material and methods

592 ESM_2: Additional simulation results

593 ESM_3: Detailed site-specific projections

594

595 **REFERENCES:**

- 596 Allen, C. D., D. D. Breshears, and N. G. McDowell. 2015. On underestimation of global vulnerability to
 597 tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* **6**:1-55.
- 598 Allen, C. D., A. K. Macalady, H. Chenchouni, D. Bachelet, N. McDowell, M. Vennetier, T. Kitzberger, A.
 599 Rigling, D. D. Breshears, E. H. Hogg, P. Gonzalez, R. Fensham, Z. Zhang, J. Castro, N.
 600 Demidova, J. H. Lim, G. Allard, S. W. Running, A. Semerci, and N. Cobb. 2010. A global
 601 overview of drought and heat-induced tree mortality reveals emerging climate change risks
 602 for forests. *Forest Ecology and Management* **259**:660-684.
- 603 ATRAGENE. 1997. Standortkundliche Erhebungen, Waldbau C-Projekte Gersau, Fallenbach und
 604 Sitiwald, Forstkreis 2, Rigi-Rossberg, Kanton Schwyz.
- 605 ATRAGENE. 1999. Standortkundliche Erhebungen, Forstkreis 1, Muotathal-Fronalpstock, Kanton
 606 Schwyz.
- 607 ATRAGENE. 2004. Standortkundliche Erhebungen, Rothenthurm-Rossberg, Kanton Schwyz.
- 608 Bauhus, J., K. J. Puettmann, and C. Kühne. 2013. Close-to-nature forest management in Europe.
 609 Pages 187-213 *in* C. Messier, K. J. Puettmann, and K. D. Coates, editors. *Managing Forests as*
 610 *Complex Adaptive Systems: Building to the Challenge of Global Change*. The Earthscan Forest
 611 Library, New York.
- 612 Bebi, P., H. Bugmann, P. Lüscher, B. Lange, and P. Brang. 2016. Auswirkungen des Klimawandels auf
 613 Schutzwald und Naturgefahren. Pages 269-285 *in* A. R. Pluess, S. Augustin, and P. Brang,
 614 editors. *Wald im Klimawandel. Grundlagen für Adaptationsstrategien*. Bundesamt für
 615 Umwelt BAFU, Bern, Eidgenössische Forschungsanstalt für Wald, Schnee und Landschaft
 616 WSL, Birmensdorf, Bern, Stuttgart, Wien.
- 617 Bigler, C., O. U. Braker, H. Bugmann, M. Dobbertin, and A. Rigling. 2006. Drought as an inciting
 618 mortality factor in Scots pine stands of the Valais, Switzerland. *Ecosystems* **9**:330-343.
- 619 Bircher, N., M. Cailleret, M. Huber, and H. Bugmann. 2015. Empfindlichkeit typischer Schweizer
 620 Waldbestände auf den Klimawandel. *Schweizerische Zeitschrift für Forstwesen* **166**:408-419.
- 621 Botkin, D. B., J. F. Janak, and J. R. Wallis. 1972. Some ecological consequences of a computer model
 622 of forest growth. *Journal of Ecology* **60**:849-872.
- 623 Bugmann, H. 1996. A simplified forest model to study species composition along climate gradients.
 624 *Ecology* **77**:2055-2074.
- 625 Bugmann, H., P. Brang, C. Elkin, P. Henne, O. Jakoby, M. Levesque, H. Lischke, A. Psomas, A. Rigling,
 626 B. Wermelinger, and N. E. Zimmermann. 2014. Climate change impacts on tree species,
 627 forest properties and ecosystem services. *in* CH2014-Impacts, editor. *Towards Quantitative*
 628 *Scenarios of Climate Change Impacts in Switzerland*. OCCR, FOEN, MeteoSwiss, C2SM,
 629 Agroscope, and ProClim, Bern, Switzerland.
- 630 CH2011. 2011. *Swiss Climate Change Scenarios CH2011*. C2SM, MeteoSwiss, ETH, NCCR Climate, and
 631 OcCC, Zurich, Switzerland.
- 632 Condes, S., and M. del Rio. 2015. Climate modifies tree interactions in terms of basal area growth and
 633 mortality in monospecific and mixed *Fagus sylvatica* and *Pinus sylvestris* forests. *European*
 634 *Journal of Forest Research* **134**:1095-1108.
- 635 Côté, S. D., T. P. Rooney, J. P. Tremblay, C. Dussault, and D. M. Waller. 2004. Ecological impacts of
 636 deer overabundance. *Annual Review of Ecology Evolution and Systematics* **35**:113-147.
- 637 Dai, A. G. 2013. Increasing drought under global warming in observations and models. *Nature*
 638 *Climate Change* **3**:52-58.
- 639 Didion, M., A. D. Kupferschmid, A. Wolf, and H. Bugmann. 2011. Ungulate herbivory modifies the
 640 effects of climate change on mountain forests. *Climatic Change* **109**:647-669.
- 641 Dorren, L., F. Berger, and B. Maier. 2005. Der Schutzwald als Steinschlagnetz. *LWF aktuell* **50**:25-27.
- 642 EEA. 2010. *Europe's Ecological Backbone: Recognising the True Value of Our Mountains*. European
 643 Environment Agency, Copenhagen.

644 Elkin, C., A. G. Gutierrez, S. Leuzinger, C. Manusch, C. Temperli, L. Rasche, and H. Bugmann. 2013. A 2
645 degrees C warmer world is not safe for ecosystem services in the European Alps. *Global*
646 *Change Biology* **19**:1827-1840.

647 Ellenberg, H., and F. Klötzli. 1972. *Waldgesellschaften und Waldstandorte der Schweiz*.
648 Schweizerische Anstalt für das Forstliche Versuchswesen, Birmensdorf, ZH.

649 Etzold, S., K. Zieminska, B. Rohner, A. Bottero, A. K. Bose, N. K. Ruehr, A. Zingg, and A. Rigling. 2019.
650 One Century of Forest Monitoring Data in Switzerland Reveals Species- and Site-Specific
651 Trends of Climate-Induced Tree Mortality. *Frontiers in Plant Science* **10**.

652 Fontes, L., J. D. Bontemps, H. Bugmann, M. Van Oijen, C. Gracia, K. Kramer, M. Lindner, T. Rotzer, and
653 J. P. Skovsgaard. 2010. Models for supporting forest management in a changing
654 environment. *Forest Systems* **19**:8-29.

655 Forrester, D. I., C. Ammer, P. J. Annighofer, A. Avdagic, I. Barbeito, K. Bielak, G. Brazaitis, L. Coll, M.
656 del Rio, L. Drossler, M. Heym, V. Hurt, M. Lof, B. Matovic, F. Meloni, J. den Ouden, M. Pach,
657 M. G. Pereira, Q. Ponette, H. Pretzsch, J. Skrzyszewski, D. Stojanovic, M. Svoboda, R. Ruiz-
658 Peinado, G. Vacchiano, K. Verheyen, T. Zlatanov, and A. Bravo-Oviedo. 2017. Predicting the
659 spatial and temporal dynamics of species interactions in *Fagus sylvatica* and *Pinus sylvestris*
660 forests across Europe. *Forest Ecology and Management* **405**:112-133.

661 Forrester, D. I., J. Nitzsche, and H. Schmid. 2019. The Experimental Forest Management project: An
662 overview and methodology of the long-term growth and yield plot network. Swiss Federal
663 Institute of Forest, Snow and Landscape Research WSL, Birmensdorf.

664 Frank, A., G. T. Howe, C. Sperisen, P. Brang, J. B. St Clair, D. R. Schmatz, and C. Heiri. 2017. Risk of
665 genetic maladaptation due to climate change in three major European tree species. *Global*
666 *Change Biology* **23**:5358-5371.

667 Frehner, M., B. Wasser, and R. Schwitter. 2005. *Nachhaltigkeit im Schutzwald (Projekt NaiS) -*
668 *Bundesamt für Umwelt BAFU. Bern.*

669 Gazda, A., P. Koscielniak, M. Hardy, E. Muter, K. Kedra, J. Bodziarczyk, M. Fraczek, K. Chwistek, W.
670 Rozanski, and J. Szwagrzyk. 2019. Upward expansion of distribution ranges of tree species:
671 Contrasting results from two national parks in Western Carpathians. *Science of the Total*
672 *Environment* **653**:920-929.

673 Gutsch, M., P. Lasch-Born, C. Kollas, F. Suckow, and C. P. O. Reyer. 2018. Balancing trade-offs
674 between ecosystem services in Germany's forests under climate change. *Environmental*
675 *Research Letters* **13**:1-12.

676 Gutierrez, A., R.S. Snell, H. Bugmann (2016) Using a dynamic forest model to predict tree species
677 distributions. *Global Ecol Biogeogr* **25**:347–358

678 HADES. 2015. *Hydrologischer Atlas der Schweiz*. University of Bern, Bern, Switzerland.

679 Hantke, R., and E. Kuriger. 2003. Überblick über die Geologie des Kantons Schwyz und seiner
680 Nachbargebiete. Pages 9-17 *in* SCNAT, editor. *Geologie und Geotope im Kanton Schwyz*.
681 Schweizerische Naturforschende Gesellschaft, Einsiedeln.

682 Huber, N. 2019. Towards robust projections of future forest dynamics: why there is no silver bullet to
683 cope with complexity. PhD Thesis, ETH Zurich.

684 IPCC. 2013. *Climate Change 2013: The Physical Science Basis*. Contribution of Working Group I to the
685 Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
686 University Press, Cambridge, United Kingdom and New York, NY, USA.

687 Irauschek, F., W. Rammer, and M. J. Lexer. 2017a. Can current management maintain forest
688 landscape multifunctionality in the Eastern Alps in Austria under climate change? *Regional*
689 *Environmental Change* **17**:33-48.

690 Irauschek, F., W. Rammer, and M. J. Lexer. 2017b. Evaluating multifunctionality and adaptive capacity
691 of mountain forest management alternatives under climate change in the Eastern Alps.
692 *European Journal of Forest Research* **136**:1051-1069.

693 Jump, A. S., J. M. Hunt, and J. Penuelas. 2006. Rapid climate change-related growth decline at the
694 southern range edge of *Fagus sylvatica*. *Global Change Biology* **12**:2163-2174.

695 Keane, R. E., D. McKenzie, D. A. Falk, E. A. H. Smithwick, C. Miller, and L. K. B. Kellogg. 2015.
696 Representing climate, disturbance, and vegetation interactions in landscape models.
697 *Ecological Modelling* **309**:33-47.

698 Klopčič, M., M. Mina, H. Bugmann, and A. Boncina. 2017. The prospects of silver fir (*Abies alba* Mill.)
699 and Norway spruce (*Picea abies* (L.) Karst) in mixed mountain forests under various
700 management strategies, climate change and high browsing pressure. *European Journal of*
701 *Forest Research* **136**:1071-1090.

702 Kupferschmid, A. D., P. Brang, W. Schönenberger, and H. Bugmann. 2006. Predicting tree
703 regeneration in *Picea abies* snag stands. *European Journal of Forest Research* **125**:163-179.

704 Kupferschmid, A. D., and H. Bugmann. 2005. Effect of microsites, logs and ungulate browsing on
705 *Picea abies* regeneration in a mountain forest. *Forest Ecology and Management* **205**:251-265.

706 Kupferschmid, A. D., C. Heiri, M. Huber, M. Fehr, M. Frei, P. Gmür, N. Imesch, J. Zinggeler, P. Brang, J.-
707 C. Clivaz, and O. Odermatt. 2015. Einfluss wildlebender Huftiere auf die Waldverjüngung: ein
708 Überblick für die Schweiz. *Schweizerische Zeitschrift für Forstwesen* **166**:420-431.

709 Lafond, V., G. Lagarrigues, T. Cordonnier, and B. Courbaud. 2014. Uneven-aged management options
710 to promote forest resilience for climate change adaptation: effects of group selection and
711 harvesting intensity. *Annals of Forest Science* **71**:937-937.

712 Langner, A., F. Irauschek, S. Perez, M. Pardos, T. Zlatanov, K. Ohman, E. M. Nordstrom, and M. J.
713 Lexer. 2017. Value-based ecosystem service trade-offs in multi-objective management in
714 European mountain forests. *Ecosystem Services* **26**:245-257.

715 Leuch, B. A., K. Streit, and P. Brang. 2017. Naturnaher Waldbau im Klimawandel. *Merkblatt für die*
716 *Praxis* **59**:1-8.

717 Leuschner, C., and I. C. Meier. 2018. The ecology of Central European tree species: Trait spectra,
718 functional trade-offs, and ecological classification of adult trees. *Perspectives in Plant Ecology*
719 *Evolution and Systematics* **33**:89-103.

720 Levesque, M., M. Saurer, R. Siegwolf, B. Eilmann, P. Brang, H. Bugmann, and A. Rigling. 2013. Drought
721 response of five conifer species under contrasting water availability suggests high
722 vulnerability of Norway spruce and European larch. *Global Change Biology* **19**:3184-3199.

723 Lexer, M. J., and H. Bugmann. 2017. Mountain forest management in a changing world. *European*
724 *Journal of Forest Research* **136**:981-982.

725 Lexer, M. J., K. Honninger, H. Scheifinger, C. Matulla, N. Groll, H. Kromp-Kolb, K. Schadauer, F.
726 Starlinger, and M. Englisch. 2002. The sensitivity of Austrian forests to scenarios of climatic
727 change: a large-scale risk assessment based on a modified gap model and forest inventory
728 data. *Forest Ecology and Management* **162**:53-72.

729 Lindner, M., M. Maroschek, S. Netherer, A. Kremer, A. Barbati, J. Garcia-Gonzalo, R. Seidl, S. Delzon,
730 P. Corona, M. Kolstrom, M. J. Lexer, and M. Marchetti. 2010. Climate change impacts,
731 adaptive capacity, and vulnerability of European forest ecosystems. *Forest Ecology and*
732 *Management* **259**:698-709.

733 Maroschek, M., W. Rammer, and M. J. Lexer. 2015. Using a novel assessment framework to evaluate
734 protective functions and timber production in Austrian mountain forests under climate
735 change. *Regional Environmental Change* **15**:1543-1555.

736 Mina, M., H. Bugmann, T. Cordonnier, F. Irauschek, M. Klopčič, M. Pardos, and M. Cailleret. 2017a.
737 Future ecosystem services from European mountain forests under climate change. *Journal of*
738 *Applied Ecology* **54**:389-401.

739 Mina, M., H. Bugmann, M. Klopčič, and M. Cailleret. 2017b. Accurate modeling of harvesting is key
740 for projecting future forest dynamics: a case study in the Slovenian mountains. *Regional*
741 *Environmental Change* **17**:49-64.

742 Minerbi, S., A. Cescatti, P. Cherubini, K. Hellrigl, G. Markart, M. Saurer, and C. Mutinelli. 2006. Scots
743 Pine dieback in the Isarco Valley due to severe drought in the summer of 2003. *Forest*
744 *Observer* **2**:89-144.

745 Mladenoff, D. J. 2004. LANDIS and forest landscape models. *Ecological Modelling* **180**:7-19.

746 Noack, A., W. Schönenberger, and P. Thee. 2004. Schützen Windwurfflächen vor Lawinen und
747 Steinschlag? *Wald und Holz* **10**:43-46.

748 Ott, E., M. Frehner, H. U. Frey, and P. Lüscher. 1997. Gebirgsnadelwälder: ein praxisorientierter
749 Leitfaden für eine standortgerechte Waldbehandlung. P. Haupt Verlag, Bern, Switzerland.

750 Penuelas, J., and M. Boada. 2003. A global change-induced biome shift in the Montseny mountains
751 (NE Spain). *Global Change Biology* **9**:131-140.

752 Pretzsch, H., R. Grote, B. Reineking, T. Roetzer, and S. Seifert. 2008. Models for forest ecosystem
753 management: A European perspective. *Annals of Botany* **101**:1065-1087.

754 Price, M. F., G. Gratzer, L. A. Duguma, T. Kohler, D. Maselli, and R. Romeo. 2011. Mountain Forests in
755 a Changing World - Realizing Values, addressing challenges., Food and Agriculture
756 Organization of the United Nations, FAO, Rome.

757 R Development Core Team. 2017. R: A language and environment for statistical computing. R
758 Foundation for Statistical Computing, Vienna, Austria.

759 Radtke, A., D. Toe, F. Berger, S. Zerbe, and F. Bourrier. 2014. Managing coppice forests for rockfall
760 protection: lessons from modeling. *Annals of Forest Science* **71**:485-494.

761 Rasche, L., L. Fahse, A. Zingg, and H. Bugmann. 2011. Getting a virtual forester fit for the challenge of
762 climatic change. *Journal of Applied Ecology* **48**:1174-1186.

763 Rasche, L., L. Fahse, A. Zingg, and H. Bugmann. 2012. Enhancing gap model accuracy by modeling
764 dynamic height growth and dynamic maximum tree height. *Ecological Modelling* **232**:133-
765 143.

766 Reyer, C. P. O., N. Brouwers, A. Rammig, B. W. Brook, J. Epila, R. F. Grant, M. Holmgren, F.
767 Langerwisch, S. Leuzinger, W. Lucht, B. Medlyn, M. Pfeifer, J. Steinkamp, M. C. Vanderwel, H.
768 Verbeeck, and D. M. Vilella. 2015. Forest resilience and tipping points at different spatio-
769 temporal scales: approaches and challenges. *Journal of Ecology* **103**:5-15.

770 Rigling, A., C. Bigler, B. Eilmann, E. Feldmeyer-Christe, U. Gimmi, C. Ginzler, U. Graf, P. Mayer, G.
771 Vacchiano, P. Weber, T. Wohlgemuth, R. Zweifel, and M. Dobbertin. 2013. Driving factors of a
772 vegetation shift from Scots pine to pubescent oak in dry Alpine forests. *Global Change*
773 *Biology* **19**:229-240.

774 Sanchez-Salguero, R., J. C. Linares, J. J. Camarero, J. Madrigal-Gonzalez, A. Hevia, A. Sanchez-Miranda,
775 J. A. Ballesteros-Canovas, R. Alfaro-Sanchez, A. I. Garcia-Cervigon, C. Bigler, and A. Rigling.
776 2015. Disentangling the effects of competition and climate on individual tree growth: A
777 retrospective and dynamic approach in Scots pine. *Forest Ecology and Management* **358**:12-
778 25.

779 Schuler, L. J., H. Bugmann, G. Petter, and R. S. Snell. 2019. How multiple and interacting disturbances
780 shape tree diversity in European mountain landscapes. *Landscape Ecology* **34**:1279-1294.

781 Schulze, E. D., O. Bouriaud, J. Waldchen, N. Eisenhauer, H. Walentowski, C. Seele, E. Heinze, U.
782 Pruschitzki, G. Danila, G. Marin, D. Hessenmoller, L. Bouriaud, and M. Teodosiu. 2014.
783 Ungulate browsing causes species loss in deciduous forests independent of community
784 dynamics and silvicultural management in Central and Southeastern Europe. *Annals of Forest*
785 *Research* **57**:267-288.

786 Schmid, U. 2014. Bewirtschaftung ausgewählter Wälder der Schweiz unter Klimawandel. Master-
787 Thesis. ETH Zurich, Zurich

788 Schumacher, S., H. Bugmann, and D. Mladenoff. 2004. Improving the formulation of tree growth and
789 succession in a spatially explicit landscape model. *Ecological Modelling* **180**:175-194.

790 Schwitter, R. 2006. Behandlung von Jungwaldbeständen auf Sturmflächen im Schutzwald unter
791 Berücksichtigung der langfristigen Entwicklung. Schweizerische Gebirgswaldpflegegruppe
792 (GWG), Schwyz and Maienfeld.

793 Schwitter, R. 2013. Gebirgswald- und Schutzwaldpflege: Eine Orientierungshilfe für die Praxis., ibW
794 Bildungszentrum Wald, Maienfeld, Fachstelle für Gebirgswald (GWP).

795 Schwitter, R., A. Sandri, P. Bebi, T. Wohlgemuth, and P. Brang. 2015. Lehren aus Vivian für den
796 Gebirgswald - im Hinblick auf den nächsten Sturm. *Schweizerische Zeitschrift für Forstwesen*
797 **166**:159-167.

798 Schwörer, C., P. D. Henne, and W. Tinner. 2014. A model-data comparison of Holocene timberline
799 changes in the Swiss Alps reveals past and future drivers of mountain forest dynamics. *Global*
800 *Change Biology* **20**:1512-1526.

801 SCNAT. 2016. Brennpunkt Klima Schweiz. Grundlagen, Folgen und Perspektiven. Swiss Academy of
802 Sciences, Bern, Switzerland.

803 Seidl, R., W. Rammer, and K. Blennow. 2014. Simulating wind disturbance impacts on forest
804 landscapes: Tree-level heterogeneity matters. *Environmental Modelling & Software* **51**:1-11.

805 Seidl, R., W. Rammer, and M. J. Lexer. 2011a. Climate change vulnerability of sustainable forest
806 management in the Eastern Alps. *Climatic Change* **106**:225-254.

807 Seidl, R., W. Rammer, R. M. Scheller, and T. A. Spies. 2012. An individual-based process model to
808 simulate landscape-scale forest ecosystem dynamics. *Ecological Modelling* **231**:87-100.

809 Seidl, R., M. J. Schelhaas, and M. J. Lexer. 2011b. Unraveling the drivers of intensifying forest
810 disturbance regimes in Europe. *Global Change Biology* **17**:2842-2852.

811 Seidl, R., D. Thom, M. Kautz, D. Martin-Benito, M. Peltoniemi, G. Vacchiano, J. Wild, D. Ascoli, M.
812 Petr, J. Honkaniemi, M. J. Lexer, V. Trotsiuk, P. Mairota, M. Svoboda, M. Fabrika, T. A. Nagel,
813 and C. P. O. Reyer. 2017. Forest disturbances under climate change. *Nature Climate Change*
814 **7**:395-402.

815 Streit, K., B. A. Leuch, and P. Brang. 2017. Der richtige Eingriff zur richtigen Zeit. *Wald und Holz* **98**:32-
816 35.

817 Temperli, C., H. Bugmann, and C. Elkin. 2012. Adaptive management for competing forest goods and
818 services under climate change. *Ecological Applications* **22**:2065-2077.

819 Temperli, C., H. Bugmann, and C. Elkin. 2013. Cross-scale interactions among bark beetles, climate
820 change, and wind disturbances: a landscape modeling approach. *Ecological Monographs*
821 **83**:383-402.

822 Temperli, C., T. T. Veblen, S. J. Hart, D. Kulakowski, and A. J. Tepley. 2015. Interactions among spruce
823 beetle disturbance, climate change and forest dynamics captured by a forest landscape
824 model. *Ecosphere* **6**:1-20.

825 Thöle, L., C. Schwörer, D. Colombaroli, E. Gobet, P. Kaltenrieder, J. van Leeuwen, and W. Tinner.
826 2016. Reconstruction of Holocene vegetation dynamics at Lac de Bretaye, a high-mountain
827 lake in the Swiss Alps. *Holocene* **26**:380-396.

828 Thornton, P. E., S. W. Running, and M. A. White. 1997. Generating surfaces of daily meteorological
829 variables over large regions of complex terrain. *Journal of Hydrology* **190**:214-251.

830 Thrippleton, T., H. Bugmann, and R. S. Snell. 2018. Herbaceous competition and browsing may induce
831 arrested succession in central European forests. *Journal of Ecology* **106**:1120-1132.

832 UNEP. 2019. Emissions Gap Report 2019. Executive summary., United Nations Environment
833 Programme, Nairobi.

834 Vitasse, Y., G. Hoch, C. F. Randin, A. Lenz, C. Kollas, and C. Korner. 2012. Tree recruitment of
835 European tree species at their current upper elevational limits in the Swiss Alps. *Journal of*
836 *Biogeography* **39**:1439-1449.

837 Watt, A. S. 1947. Pattern and Process in the Plant Community. *Journal of Ecology* **35**:1-22.

838 Whiteman, D. C. 2000. Mountain meteorology : fundamentals and applications. Oxford University
839 Press, New York.

840 Wohlgenuth, T., V. Doublet, C. Nussbaumer, L. Feichtinger, and A. Rigling. 2018. Baumartenwechsel
841 in den Walliser Waldföhrenwäldern verstärkt nach grossen Störungen. *Schweizerische*
842 *Zeitschrift für Forstwesen* **169**:260-268.

843 Wohlgenuth, T., and B. Moser. 2018. Ten years of vegetation dynamics in a forest fire patch in Leuk
844 (Valais). *Schweizerische Zeitschrift für Forstwesen* **169**:279 - 289.

845 Wolfslehner, B., and R. Seidl. 2010. Harnessing Ecosystem Models and Multi-Criteria Decision
846 Analysis for the Support of Forest Management. *Environmental Management* **46**:850-861.

847 Yousefpour, R., C. Temperli, J. B. Jacobsen, B. J. Thorsen, H. Meilby, M. J. Lexer, M. Lindner, H.
848 Bugmann, J. G. Borges, J. H. N. Palma, D. Ray, N. E. Zimmermann, S. Delzon, A. Kremer, K.
849 Kramer, C. P. O. Reyer, P. Lasch-Born, J. Garcia-Gonzalo, and M. Hanewinkel. 2017. A
850 framework for modeling adaptive forest management and decision making under climate
851 change. *Ecology and Society* **22**:1-14.

852 Zou, C. B., G. A. Barron-Gafford, and D. D. Breshears. 2007. Effects of topography and woody plant
853 canopy cover on near-ground solar radiation: Relevant energy inputs for ecohydrology and
854 hydrology. *Geophysical Research Letters* **34**:1-6.
855