


Synchronized broadcast spawning by six invertebrates (Echinodermata and Mollusca) in the north-western Red Sea

Journal Article

Author(s):

Webb, Alice E.; Engelen, Aschwin H.; Bouwmeester, Jessica; van Dijk, Inge; Geerken, Esmee; [Lattaud, Julie](#) ; Engelen, Dario; de Bakker, Bernadette S.; de Bakker, Didier M.

Publication date:

2021

Permanent link:

<https://doi.org/10.3929/ethz-b-000479154>

Rights / license:

[Creative Commons Attribution 4.0 International](#)

Originally published in:

Marine Biology 168(5), <https://doi.org/10.1007/s00227-021-03871-6>



Synchronized broadcast spawning by six invertebrates (Echinodermata and Mollusca) in the north-western Red Sea

Alice E. Webb¹ · Aschwin H. Engelen² · Jessica Bouwmeester^{3,4} · Inge van Dijk⁵ · Esmee Geerken¹ · Julie Lattaud⁶ · Dario Engelen^{1,2,3,4,5,6,7,8} · Bernadette S. de Bakker⁷ · Didier M. de Bakker⁸ 

Received: 26 November 2020 / Accepted: 21 March 2021
© The Author(s) 2021

Abstract

On the evenings of June 11 and 12, 2019, 5 and 6 days before full moon, broadcast spawning by four echinoderm species and two mollusc species was observed on the Marsa Shagra reef, Egypt (25° 14' 44.2" N, 34° 47' 49.0" E). Water temperature was 28 °C and the invertebrates were observed at 2–8 m depth. The sightings included a single basket star *Astroboa nuda* (Lyman 1874), 2 large *Tectus dentatus* (Forsk. 1775) sea snails, 14 individuals of the *Leiaster* cf. *leachi* (Gray 1840) sea star and 1 *Mithrodia clavigera* (Lamarck 1816) sea star, 3 *Pearsonothuria graeffei* (Semper 1868) sea cucumbers, and 2 giant clams, *Tridacna maxima* (Röding 1798). The observations presented here provide relevant information on broadcast spawning of non-coral invertebrate taxa in the Red Sea, where spawning is considerably less well documented than in other tropical geographical regions such as the Indo-Pacific and Caribbean.

Introduction

The sedentary nature of many invertebrates on coral reefs renders them heavily reliant on dispersal during the pelagic larval period of their life cycle to ensure connectivity

amongst reef populations (Atchison et al. 2008; Nunes et al. 2011). Broadcast spawning occurs across most marine phyla (e.g. Hagman and Vize 2003; Ritson-Williams et al. 2005; Hoffman et al. 2011; Bouwmeester 2013) and is a mode of reproduction in which large quantities of male and female gametes are released into the water column within a short period of time. Success of the strategy, however, relies on reproductive synchrony among intraspecific individuals within a few hours of specific nights each year. Such annual spawning events have been shown to be closely tied to factors such as annual temperature variation, lunar cycles and photoperiod length (Boch et al. 2011; Sweeney et al. 2011; Keith et al. 2016). To date, invertebrate species other than scleractinian corals have received noticeably less attention with regard to reproduction and this is particularly relevant for the Red Sea, where only very few studies have been conducted on multi-species broadcast spawning (Bouwmeester et al. 2016). Here we describe spawning observations of 23 non-coral invertebrates belonging to six species and two phyla.

Responsible Editor: J. Grassle Reviewers: undisclosed experts.

✉ Didier M. de Bakker
didierdebakker@gmail.com

- ¹ Department of Ocean Systems, NIOZ-Royal Netherlands Institute for Sea Research, and Utrecht University, 1790 AB Den Burg, The Netherlands
- ² Centro de Ciências Do Mar Do Algarve (CCMAR), University of Algarve, 8005 139 Faro, Portugal
- ³ Smithsonian Conservation Biology Institute, National Zoological Park Department, Front Royal, VA 22630, USA
- ⁴ Hawaii Institute of Marine Biology, Kaneohe, HI 96744, USA
- ⁵ Alfred-Wegener-Institut, Helmholtz-Zentrum Für Polar-Und Meeresforschung, 27570 Bremerhaven, Germany
- ⁶ Biogeoscience Group, ETH Zurich, 8092 Zurich, Switzerland
- ⁷ Department of (Neuro)Pathology, Amsterdam UMC, University of Amsterdam, 1105AZ Amsterdam, The Netherlands
- ⁸ Wageningen Marine Research, Wageningen University and Research, 1780 AB Den Helder, The Netherlands

Materials and methods

On the evenings of June 11 and 12, 2019 at 19:00–20:30 h, broadcast spawning by six invertebrate species (Mollusca and Echinodermata) was observed on the Marsa Shagra reef,

Egypt (~25 km north of Marsa Alam, 25° 14' 44.2" N 34° 47' 49.0" E). This fringing reef is characterized by a wide (~1000 m) and shallow (< 1 m at high tide) reef flat underlain by a large carbonate framework. The reef flat is interrupted by a sand channel (bay) shouldered by gradually downward sloping reefs on both sides, and which transitions into a reef shelf at ~20 m depth. All spawning was observed on the sloping edges of the channel at 2–8 m depth (Table 1). The reef slope is characterized by a complex structural morphology and a 60–70% cover of scleractinian coral (mainly *Acropora* spp., *Porites* spp. and *Pocillopora* spp., pers. obs. DdB). Water temperature at 5 m depth was 28 °C, measured with a Suunto D4i dive computer. All observations were made while SCUBA diving and an underwater photographic record was taken for each observation. Species identification, and the identification of sperm or egg spawn, was performed visually in situ. Sperm release was characterized by a “smoky” cloud emitted by the individuals, and egg release was characterized by a “non-smoky” cloud of tiny conspicuous eggs. Taxonomic identifications were later confirmed by Gustav Paulay (FLMNH, USA) based on the extensive photographic and video material. Due to permit limitations and the limited number of individuals that were observed spawning, no voucher specimens were collected for posteriori identification.

Results

Spawning observations

A single basket star, *Astroboa nuda* (Lyman 1874), was observed on June 11 perched on top of a fire coral releasing female gametes (Fig. 1a, Table 1). When encountered, the individual was in a typical feeding position with arm branches extended and was releasing a vast number of eggs (~1 mm diameter) that were dispersed by the prevailing current. When illuminated, it continued spawning but drew in its arms towards the central disk, which is typical for this extremely light-sensitive species (Tsurumal and Marder 1966; Hendler et al. 1995). A few minutes later, two large top-shell snails (Trochidae), *Tectus dentatus* (Forsk. 1775)

were encountered releasing a jet of male gametes into the water column (Fig. 1b). They were found 10 m apart, both perched on top of a coral mound (Table 1).

On June 12, 14 starfish *Leiaster* cf. *leachi* (Gray 1840) were seen broadcast spawning. Several individuals released sperm, while other individuals were observed releasing a cloud of eggs (Table 1). Individuals were several meters apart from each other (i.e. no clustering) and generally on top of coral mounds. Most of them exhibited the same spawning behaviours with raised discs and perched on the tips of their arms (Fig. 1c). Another sea star species, *Mithrodia clavigera* (Lamarck 1816), was encountered releasing male gametes in a flattened position on top of a coral mound (Table 1).

During the same evening, spawn was observed emanating from three *Pearsonothuria graeffei* (Semper 1868) sea cucumbers (Table 1). One was observed on a sandy bottom with half of its body lifted off the seafloor in a vertical position. It swayed with the waves' surging motion while releasing a cloud of male gametes. The two other individuals were encountered briefly emerging from the cryptic reef environment to release a single cloud of male gametes before rapidly retreating into the reef (ESM_1). Lastly, two giant clams (*Tridacna maxima*, Röding 1798) were encountered, releasing a cloud of male gametes. Discharge lasted for only a few seconds and occurred through the excurrent siphons. The two individuals were in close proximity (1 m apart), embedded in the same coral mound. They spawned within seconds of each other (ESM_1).

Discussion

Spawning times of four out of the six species encountered here were not yet described for the Red Sea and to our knowledge, this is the first field observation of a basket star releasing female gametes. Relatively little is known about the life history of *A. nuda*, but our observations suggest that this species—like many Ophiuroids—relies on broadcast spawning. Apart from our description of mass egg release in the north-western Red Sea, the only other in situ description of spawning appears to be a single *A. nuda* individual

Table 1 Timetable and specifications of the recorded invertebrate spawning

Days before full moon	Species	Number of individuals	Time of spawning	Type of gametes	Depth (m)
6 (June 11)	<i>Astroboa nuda</i>	1 ♀	19:45	Eggs	8
	<i>Tectus dentatus</i>	2 ♂	20:00–20:10	Sperm	3–5
5 (June 12)	<i>Leiaster</i> cf. <i>leachi</i>	5 ♂ & 9 ♀	19:30–20:15	Sperm & eggs	3–8
	<i>Mithrodia clavigera</i>	1 ♂	20:15	Sperm	2
	<i>Tridacna maxima</i>	2 ♀	19:50	Sperm	3
	<i>Pearsonothuria graeffei</i>	3 ♂	19:50–20:10	Sperm	2–5




Fig. 1 Nocturnal spawning of non-coral invertebrates on the shallow reef of the Egyptian Red Sea. **a** *Astroboa nuda* releasing eggs. **b** Continuous jet-like release of male gametes from the siphon of a *Tectus*

dentatus sea snail. **c** *Leiaster* cf. *leachi* discharging female gametes in typical upright position. Video footage of all observed species is provided as Supplementary Resource 1 (ESM_1)

releasing sperm two nights before the full moon (July 2012) on the Red Sea reef Al Fahal (Saudi Arabia) (Bouwmeester 2013). *Astroboa nuda* are known to return to the same location on the reef at night to feed (Tsurumal and Marder

1966), and this behaviour did not appear to be affected by their spawning activity.

Observations of natural spawning of the two sea star species have previously been described in the Red Sea on Al

Fahal Reef (Bouwmeester et al. 2016). They found a single individual of *M. clavigera* and four individuals of *L. leachi* that released sperm in April, 4–5 days after the full moon, and a single *L. leachi* that spawned two days after the May full moon in 2014 (Bouwmeester et al. 2016). Together with our observations in June, this might indicate that both species have a relatively extended period of potential spawning, yet a more thorough assessment is needed to support this hypothesis. Other sea star species from the Red Sea have been observed to spawn in winter, spring and summer (Pearse 1968). Although both species have a relatively broad distribution range, observations of spawning are scarce. Komatsu (1973) describes a single *L. leachi* sea star in an aquarium, which released eggs for ~ 1 h from the dorsal–lateral parts of the arms (August 11, 1972, at 10:00 h). Very little is known about the reproduction and development of these sea stars—and tropical sea stars in general (Bos et al. 2008)—except that both species appear to produce relatively small eggs (~ 140 µm diameter) and have a planktonic-feeding larval stage (Yamaguchi 1975; Chia et al. 1993).

The holothurian *P. graeffei* is found throughout the wider tropical Indo-Pacific and has been recorded spawning in various locations, including Thailand (Scott et al. 2015), the South China Sea (Purwati 2003), Great Barrier Reef (Purcell et al. 2012), the Maldives, the Seychelles and Mozambique (Friedman 2008). Described individuals are generally found exhibiting the typical characteristics of sea cucumber spawning, with the anterior part of the body stretched and in a vertical position, while the posterior part of the body remained on the substrate. Similar to our observation, this species has been described spawning simultaneously with other echinoderms (Scott et al. 2015) and in most recordings two or more individuals were encountered a few meters apart. The observed *P. graeffei* on the Marsa Shagra reef are the first to our knowledge to be found exhibiting spawning behaviour in the Red Sea. Previously described encounters in other geographical regions occurred between 15:00 h and sunset, and the lunar phase was usually between waning crescent and new moon. The individuals encountered in the Red Sea differ from these previous sightings at they were found releasing gametes in full darkness around 20:00 h and under a waxing gibbous moon.

The only description of reproduction in *T. dentatus* is a case of stimulated egg release by exposing several individuals to variable temperatures in a controlled aquarium environment (Eisawy 1970). Each female released approximately 1 million eggs. The same article provides some details regarding observations of *T. dentatus* spawning in its natural environment, but extensive descriptions are essentially absent. Eisawy (1970) described a spawning period ranging from April until July with an approximate lunar periodicity (and at an optimum water temperature of 27–28 °C), which complies with our observations in

June. The reproduction in *T. dentatus* appears to be comparable to other Trochidae species. *Trochus niloticus* males have also been observed to release sperm in the evening (18:00–22:00 h), and for an extended period of time (10 min to an hour) on the Great Barrier Reef, while females spawned for up to 15 min (Heslinga and Hillmann 1981). Although the sexes of *T. dentatus* are separate (Eisawy 1970), we did not observe any females releasing eggs. Spawning in Trochidae species is generally initiated by males, although females do not always spawn in response to sperm release (Heslinga and Hillmann 1981). It is possible that *T. dentatus* females released eggs later in the night or even several days later. Until our spawning observation, *T. dentatus* snails had rarely been observed on top of the reef of Marsa Shagra. Like many of the closely related Trochidae species, *T. dentatus* are known to spend most of their time hidden. Their increased activity and displacement to the reef surface at nights around the full moon, however, has also been described for *T. niloticus* (based on anecdotal evidence from fishermen).

The spawning behaviour of *T. maxima* appears to correspond to previous descriptions of natural gamete release in the Red Sea. The reproductive strategy of *T. maxima* is relatively well described (See Roa-Quiaoit (2005) and references therein). This is partly because these clams are fished and cultured for commercial purposes (food, shell craft industry and the aquarium trade) (Lucas 1994). Throughout the Red Sea, *T. maxima* spawn June–December (Kilada et al. 1998), although the length of the spawning season can differ due to latitudinal variations in water temperature or the availability of food (Roa-Quiaoit 2005). Female gametes reach their largest diameter in June (~ 95 µm) shortly before spawning—which coincides with the timing of our observations—and are generally released within hours after the male gametes. Like many hermaphroditic broadcast spawners, this strategy reduces the chance of self-fertilization (Lucas 1994).

Spawning observations reported here contribute to the still poorly documented spawning of non-coral invertebrates in the Red Sea and shed light on their spawning behaviour. Previous reports combined with our findings provide evidence for synchronized inter-taxa mass spawning on the coral reefs of the Red Sea. Only a few dedicated surveys to document spawning in invertebrates have been conducted in the Red Sea, and it is to be expected that many more species spawn around the same period.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00227-021-03871-6>.

Acknowledgements We thank the team of Red Sea Diving Safaris at Marsa Shagra Village and in particular Mohamed Ezz for sharing his valuable local insights. We would also like to extend our gratitude to Professor Gustav Paulay, Florida Museum of Natural History, for his

help in identifying the encountered species and members of the Marsa Shagra diving group. We thank the Editor and reviewer for their careful reading of the manuscript and their constructive remarks.

Author contribution Conceptualisation: AEW and DMdB. Methodology: AEW, AHE, and DMdB. Data collection: AEW, AHE, IvD, EG, JL, DE, BSdB, and DMdB. Writing—original draft preparation: AEW and DMdB. Writing—review and editing: AEW, AHE, JB, IvD, EG, JL, DE, BSdB, and DMdB. Funding acquisition: AHE. Supervision: AEW, JB, and DMdB.

Funding This study received Portuguese national funds from FCT—Foundation for Science and Technology through project UIDB/04326/2020 and contract CEECINST/00114/2018.

Availability of data and materials All data and relevant footage collected during this study are presented in this article or the accompanying electronic supplementary materials. Additional photographs are available on request.

Code availability Not applicable.

Declarations

Conflict of interest The authors have no conflict of interest to declare.

Ethical approval Access to the study site was provided under the authorisation of the Ecological Team of the Red Sea Diving Safari dive centre. All precautions were taken to minimise the impact on the marine habitat at the study site. As this study was purely observational, no additional permits or ethical approval were required.

Consent to participate Not applicable.

Consent for publication Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Atchison AD, Sammarco PW, Brazeau DA (2008) Genetic connectivity in corals on the Flower Garden Banks and surrounding oil/gas platforms, Gulf of Mexico. *J Exp Mar Biol Ecol* 365:1–12. <https://doi.org/10.1016/j.jembe.2008.07.002>
- Boch CA, Ananthasubramaniam B, Sweeney AM, Doyle Iii FJ, Morse DE (2011) Effects of light dynamics on coral spawning synchrony. *Biol Bull* 220:161–173. <https://doi.org/10.1086/BBLv220n3p161>
- Bos AR, Gumanao GS, Alipoyo JC, Cardona LT (2008) Population dynamics, reproduction and growth of the Indo-Pacific horned sea star, *Protoreaster nodosus* (Echinodermata; Asteroidea). *Mar Biol* 156:55–63. <https://doi.org/10.1007/s00227-008-1064-2>
- Bouwmeester J (2013) Reproduction patterns of scleractinian corals in the central Red Sea. Dissertation, King Abdullah University of Science and Technology. <https://doi.org/10.25781/KAUST-73163>
- Bouwmeester J, Gatins R, Giles EC, Sinclair-Taylor TH, Berumen ML (2016) Spawning of coral reef invertebrates and a second spawning season for scleractinian corals in the central Red Sea. *Invertebr Biol* 135:273–284. <https://doi.org/10.1111/ivb.12129>
- Chia FS, Oguro C, Komatsu M (1993) Sea-star (asteroid) development. *Oceanogr Mar Biol Annu Rev* 31:223–257
- de Lamarck JBPA (1816) *Asterie*. Histoire naturelle des animaux sans vertèbres 2. Verdrière, Paris, pp 547–568
- Eisawy A (1970) The spawning and development of *Trochus* (*Tectus*) *dentatus* Forskal. *Bull Nat Inst Oceanogr Fish* 1:379–394
- Forsk P (1775) *Descriptiones animalium avium, amphibiorum, piscium, insectorum, vermium quae in itinere orientali observavit*. Hauniae, Copenhagen
- Friedman K (2008) Pacific sea cucumber dataset from the Secretariat of the Pacific Community, Reef Fisheries Observatory. *S Pac Comm Beche-de-Mer Inf Bull* 27:38–39
- Gray JE (1840) A synopsis of the genera and species of the class Hypostoma (*Asterias*, Linnaeus). *Ann Mag Nat Hist* 6:275–290
- Hagman DK, Vize PD (2003) Mass spawning by two brittle star species, *Ophioderma rubicundum* and *O. squamosissimum* (Echinodermata: Ophiuroidea), at the Flower Garden Banks. *Gulf of Mexico Bull Mar Sci* 72:871–876
- Hendler G, Miller JE, Pawson DL, Kier PM (1995) *Echinoderms of Florida and the Caribbean: Sea stars, sea urchins, and allies*. Smithsonian Institution Press, Washington. <https://doi.org/10.1086/419711>
- Heslinga G, Hillmann A (1981) Hatchery culture of the commercial top snail *Trochus niloticus* in Palau, Caroline Islands. *Aquac* 22:35–43. [https://doi.org/10.1016/0044-8486\(81\)90131-9](https://doi.org/10.1016/0044-8486(81)90131-9)
- Hoffman JI, Clarke A, Linse K, Peck LS (2011) Effects of brooding and broadcasting reproductive modes on the population genetic structure of two Antarctic gastropod molluscs. *Mar Biol* 158:287–296. <https://doi.org/10.1007/s00227-010-1558-6>
- Keith SA, Maynard JA, Edwards AJ, Guest JR, Bauman AG, Van Hooi-donk R, Heron SF, Berumen ML, Bouwmeester J, Piromvaragorn S et al (2016) Coral mass spawning predicted by rapid seasonal rise in ocean temperature. *Proc R Soc B* 283:20160011. <https://doi.org/10.1098/rspb.2016.0011>
- Kilada R, Zakaria S, Farghalli ME (1998) Distribution and abundance of the giant clam *Tridacna maxima* (Bivalvia: Tridacnidae) in the northern Red Sea. *Bull Natl Inst Oceanogr Fish* 24:221–240
- Komatsu MA (1973) A preliminary report on the development of the sea-star, *Leiaster leachii*. *Proc Jap Soc Systemat Zool* 9:55–58. https://doi.org/10.19004/pjssz.9.0_55
- Lucas JS (1994) The biology, exploitation, and mariculture of giant clams (Tridacnidae). *Rev Fish Sci Aquac* 2:181–223. <https://doi.org/10.1080/10641269409388557>
- Lyman T (1874) Ophiuridae and Astrophytidae, new and old. *Bull Mus Comp Zool* 3:250
- Nunes FL, Norris RD, Knowlton N (2011) Long distance dispersal and connectivity in amphi-Atlantic corals at regional and basin scales. *PLoS ONE* 6:e22298. <https://doi.org/10.1371/journal.pone.0022298>
- Pearse J (1968) Patterns of reproductive periodicities in four species of Indo-Pacific echinoderms. *Proc Natl Acad Sci India Sect B Biol Sci* 67:247–279. <https://doi.org/10.1007/BF03052195>
- Purcell SW, Samyn Y CC (2012) Commercially important sea cucumbers of the world. *FAO Species Catalogue for Fishery Purposes*, FAO, Rome, p 39
- Purwati P (2003) Natural spawning observations of *Pearsonothuria graeffei*. *S Pac Comm Beche-de-Mer Inf Bull* 18:38

- Ritson-Williams R, Becerro MA, Paul VJ (2005) Spawning of the giant barrel sponge *Xestospongia muta* in Belize. *Coral Reefs* 24:160. <https://doi.org/10.1007/s00338-004-0460-4>
- Roa-Quiaoit HAF (2005) Ecology and culture of giant clams (Tridacnidae) in the Jordanian sector of the Gulf of Aqaba, Red Sea. Dissertation, Universitat Bremen
- Röding PF (1798) *Museum Boltenianum sive Catalogus cimeliorum e tribus regnis naturæ quæ olim collegerat Joa. Fried Bolten, M. D. p. d. per XL. annos proto physicus Hamburgensis. Pars secunda continens Conchylia sive Testacea univalvia, bivalvia & multivalvia. Trappii, Hamburg, pp 199*
- Scott CM, Mehrotra R, Urgell P (2015) Spawning observation of *Acanthaster planci* in the Gulf of Thailand. *Mar Biodivers* 45:621–622. <https://doi.org/10.1007/s12526-014-0300-x>
- Semper C (1868) Holothurien. In: Semper C (ed) *Reisen im Archipel der Philippinen Zweiter Theil Wissenschaftliche Resultate*. W. Engelmann, Leipzig, p 288. <https://doi.org/10.5962/bhl.title.11687>
- Sweeney AM, Boch CA, Johnsen S, Morse DE (2011) Twilight spectral dynamics and the coral reef invertebrate spawning response. *J Exp Biol* 214:770–777. <https://doi.org/10.1242/jeb.043406>
- Tsurnamal M, Marder J (1966) Observations on the basket star *Astroboa nuda* (Lyman) on coral reefs at Elat (Gulf of Aqaba). *Isr J Ecol Evol* 15:9–17. <https://doi.org/10.1080/00212210.1966.10688225>
- Yamaguchi M (1975) Coral-reef asteroids of Guam. *Biotropica* 7:12–23. <https://doi.org/10.2307/2989795>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.