ETH zürich

The Solar System calcium isotopic composition inferred from Ryugu samples

Journal Article

Author(s):

Moynier, Frédéric; Dai, W.; Yokoyama, T.; Hu, Y.; Paquet, Matthieu; Abe, Y.; Aléon, Jérôme; Alexander, Conel M.O'D.; Amari, Sachiko; Amelin, Yuri; Bajo, Ken-ichi; Bizzarro, Martin; Bouvier, Aurelie; Carlson, Robert W.; Chaussidon, M.; Choi, Byeon-Gak; Dauphas, Nicolas; Davis, Andrew M.; Di Rocco, Tommaso; Fujiya, Wataru; Schönbächler, Maria; et al.

Publication date: 2022

Permanent link: https://doi.org/10.3929/ethz-b-000581252

Rights / license: Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International

Originally published in: Geochemical Perspectives Letters 24, <u>https://doi.org/10.7185/geochemlet.2238</u>

This page was generated automatically upon download from the <u>ETH Zurich Research Collection</u>. For more information, please consult the <u>Terms of use</u>.



 $$\ensuremath{\mathbb{C}}\xspace$ 2022 The Authors Published by the European Association of Geochemistry

The Solar System calcium isotopic composition inferred from Ryugu samples

F. Moynier^{1*}, W. Dai¹, T. Yokoyama², Y. Hu¹, M. Paquet¹, Y. Abe³, J. Aléon⁴, C.M.O'D. Alexander⁵, S. Amari⁶, Y. Amelin⁷, K.-I. Bajo⁸, M. Bizzarro^{9,1}, A. Bouvier¹⁰, R.W. Carlson⁵, M. Chaussidon¹, B.-G. Choi¹¹, N. Dauphas¹², A.M. Davis¹², T. Di Rocco¹³, W. Fujiya¹⁴, R. Fukai¹⁵, I. Gautam², M.K. Haba², Y. Hibiya¹⁶, H. Hidaka¹⁷, H. Homma¹⁸, P. Hoppe¹⁹, G.R. Huss²⁰, K. Ichida²¹, T. Iizuka²², T.R. Ireland²³, A. Ishikawa², M. Ito²⁴, S. Itoh²⁵, N. Kawasaki⁸, N.T. Kita²⁶, K. Kitajima²⁶, T. Kleine²⁷, S. Komatani²¹, A.N. Krot²⁰, M.-C. Liu²⁸, Y. Masuda², K.D. McKeegan²⁸, M. Morita²¹, K. Motomura²⁹, I. Nakai²⁹, K. Nagashima²⁰, D. Nesvorný³⁰, A. Nguyen³¹, L. Nittler⁵, M. Onose²¹, A. Pack¹³, C. Park³², L. Piani³³, L. Qin³⁴, S.S. Russell³⁵, N. Sakamoto³⁶, M. Schönbächler³⁷, L. Tafla²⁸, H. Tang²⁸, K. Terada³⁸, Y. Terada³⁹, T. Usui¹⁵, S. Wada⁸, M. Wadhwa⁴⁰, R.J. Walker⁴¹, K. Yamashita⁴², Q.-Z. Yin⁴³, S. Yoneda⁴⁴, E.D. Young²⁸, H. Yui⁴⁵, A.-C. Zhang⁴⁶, T. Nakamura⁴⁷, H. Naraoka⁴⁸, T. Noguchi²⁴, R. Okazaki⁴⁸, K. Sakamoto¹⁵, H. Yabuta⁴⁹, M. Abe¹⁵, A. Miyazaki¹⁵, A. Nakato¹⁵, M. Nishimura¹⁵, T. Okada¹⁵, T. Yada¹⁵, S.-I. Watanabe¹⁷, M. Yoshikawa¹⁵, S. Tachibana⁵¹, H. Yurimoto⁸

OPEN OACCESS



The Hayabusa2 spacecraft has returned samples from the Cb-type asteroid (162173) Ryugu to Earth. Previous petrological and chemical analyses support a close link between Ryugu and CI chondrites that are presumed to be chemically the most primitive meteorites with a solar-like composition. However, Ryugu samples are highly enriched in Ca compared to typical CI chondrites. To identify the cause of this discrepancy, here we report stable Ca isotopic data (expressed as $\delta^{44/40}Ca_{SRM915a}$) for returned Ryugu samples collected from two sites. We found that samples from both sites have similar $\delta^{44/40}Ca_{SRM915a}$ (0.58 ± 0.03 ‰ and 0.55 ± 0.08 ‰, 2 s.d.) that fall within the range defined by CIs. This isotopic similarity suggests that the Ca budget of CIs and Ryugu samples is dominated by carbonates, and the variably higher Ca

contents in Ryugu samples are due to the abundant carbonates. Precipitation of carbonates on Ryugu likely coincided with a major episode of aqueous activity dated to have occurred ~5 Myr after Solar System formation. Based on the pristine Ryugu samples, the average $\delta^{44/40}Ca_{SRM915a}$ of the Solar System is defined to be 0.57 ± 0.04 ‰ (2 s.d.).

Received 28 June 2022 | Accepted 19 September 2022 | Published 19 October 2022

Introduction

Abstract

In December 2020, the JAXA Hayabusa2 spacecraft returned to Earth with the first samples collected from a Cb-type asteroid, (162173) Ryugu (Tachibana *et al.*, 2022; Yada *et al.*, 2022). Chemical, mineralogical, petrological, and isotopic analyses of these samples suggest that they are closely related to CI

chondrites. In particular, their bulk Cr and Ti isotopic signatures and the chemical abundances of most elements are within the range of CI chondrites (Nakamura E. *et al.*, 2022; Yokoyama *et al.*, 2022). Among meteorites, CI chondrites have chemical compositions that most closely resemble the Sun; therefore, they are the most representative samples of the solar nebula composition with the exception of volatile elements (Palme *et al.*, 2014).

* Corresponding author (email: moynier@ipgp.fr)



Т

^{1.} Université Paris Cité, Institut de Physique du Globe de Paris, CNRS, 75005 Paris, France

^{2.} Department of Earth and Planetary Sciences, Tokyo Institute of Technology, Tokyo 152-8551, Japan

^{3.} Graduate School of Engineering Materials Science and Engineering, Tokyo Denki University, Tokyo 120-8551, Japan

Given that the Ryugu samples did not experience any terrestrial alteration, they are likely the chemically most pristine Solar System samples available estimating the original isotopic compositions of most elements in the Solar System (Yokoyama *et al.*, 2022).

Despite many aspects of similarities with the CIs, the Ryugu samples display an apparent excess of Ca by over 50 %, which may be related to a heterogeneous distribution of carbonates (dolomite and calcite) (Nakamura T. et al., 2022; Yokoyama et al., 2022) between Ryugu, Orgueil and other CIs. Calcium is a major constituent of carbonates and can be isotopically fractionated during aqueous alteration and carbonate precipitation, leading to more than 1 ‰ variations in the ⁴⁴Ca/⁴⁰Ca ratio in terrestrial carbonates (e.g., Fantle and Tipper, 2014; Blättler and Higgins, 2017). In addition, Ca exhibits large isotopic variations among bulk carbonaceous chondrites (CC), with the 44Ca/40Ca ratio spanning a range of 1 ‰. This range is likely related to a combination of the variable modal abundances of refractory inclusions among CC (Hezel et al., 2008) that can be enriched in the lighter Ca isotopes by several per mille (Niederer and Papanastassiou, 1984; Huang et al., 2012) and the heterogeneous distribution of carbonates (Simon and DePaolo, 2010; Valdes et al., 2014; Dauphas and Pourmand, 2015). Therefore, stable Ca isotopes could be useful for investigating the origin of Ca excess in Ryugu samples compared to CIs.

Here we have analysed the stable Ca isotopic compositions of Ryugu samples collected from the first and second touchdown sites, using the collision cell equipped multicollection inductively-coupled plasma mass spectrometer (CC-MC-ICP-MS), Nu Sapphire.

- Earth and Planets Laboratory, Carnegie Institution for Science, Washington, DC, 20015, USA
- McDonnell Center for the Space Sciences and Physics Department, Washington University, St. Louis, MO 63130, USA
- Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, GD 510640, China
- 8. Natural History Sciences, IIL, Hokkaido University, Sapporo 001-0021, Japan
- 9. Centre for Star and Planet Formation, GLOBE Institute, University of Copenhagen, Copenhagen, K 1350, Denmark
- 10. Bayerisches Geoinstitut, Universität Bayreuth, Bayreuth 95447, Germany
- Department of Earth Science Education, Seoul National University, Seoul 08826, Republic of Korea
- Department of the Geophysical Sciences and Enrico Fermi Institute, The University of Chicago, 5734 South Ellis Avenue, Chicago 60637, USA
- Faculty of Geosciences and Geography, University of Göttingen, Göttingen, D-37077, Germany
- 14. Faculty of Science, Ibaraki University, Mito 310-8512, Japan
- 15. ISAS/JSEC, JAXA, Sagamihara 252-5210, Japan
- 16. General Systems Studies, The University of Tokyo, Tokyo 153-0041, Japan
- 17. Earth and Planetary Sciences, Nagoya University, Nagoya 464-8601, Japan
- Osaka Application Laboratory, SBUWDX, Rigaku Corporation, Osaka 569-1146, Japan
- 19. Max Planck Institute for Chemistry, Mainz 55128, Germany.
- Hawai'i Institute of Geophysics and Planetology, University of Hawai'i at Mānoa, Honolulu, HI 96822, USA
- 21. Analytical Technology, Horiba Techno Service Co., Ltd., Kyoto 601-8125, Japan
- 22. Earth and Planetary Science, The University of Tokyo, Tokyo 113-0033, Japan
- School of Earth and Environmental Sciences, The University of Queensland, St Lucia QLD 4072, Australia
- 24. Kochi Institute for Core Sample Research, JAMSTEC, Kochi 783-8502, Japan
- 25. Earth and Planetary Sciences, Kyoto University, Kyoto 606-8502, Japan
- 26. Geoscience, University of Wisconsin-Madison, Madison, WI 53706, USA
- 27. Max Planck Institute for Solar System Research; 37077 Göttingen, Germany
- 28. Earth, Planetary, and Space Sciences, UCLA, Los Angeles, CA 90095, USA

Samples and Methods

The samples returned by the Hayabusa2 spacecraft consist of ~5 g of materials from the Ryugu asteroid recovered during two touchdowns (Tachibana *et al.*, 2022; Yada *et al.*, 2022). Approximately 3 g of samples representing the surface materials of Ryugu were collected during the first touchdown and stored in Chamber A. Approximately 2 g of samples likely representing a mixture of materials from the surface and subsurface were collected into Chamber C at a site that was close to the crater formed by the Small Carry-on Impactor, a kinetic impact experiment of the Hayabusa2 mission (Saiki *et al.*, 2017; Arakawa *et al.*, 2020). Two Ryugu samples, A0106-A0107 (Chamber A) and C0108 (Chamber C), were analysed in this study (for information on the mineralogy see https://jaxa.repo.nii.ac.jp/?action=repository_uri&item_id=48255&file_id=31&file_no=1).

Sample A0106-A0107 was prepared from a mixed aggregate of A0106 (1.6 mg) and A0107 (27.3 mg). In addition to the Ryugu samples, fusion-crust free bulk samples of six CC, Orgueil (CI1), Alais (CI1), Tarda (C2-ungrouped), Tagish Lake (C2ungrouped), Murchison (CM2), and Allende (CV3), were analysed in the same way for comparison. See Table S-1 for the weights and providers of the meteorite samples. All samples were dissolved in PFA vials with a mixture of concentrated HF and HNO₃ at the Tokyo Institute of Technology (Yokoyama *et al.*, 2022).

After dissolution, aliquots of ~0.15 % of the solutions containing ~5 μ g of Ca were transferred and dedicated for our study. All the sample aliquots were dried and redissolved in 0.4 ml of 4 mol/L HNO₃ in preparation for Ca chemical purification and isotopic measurements at the Institut de Physique du Globe, following Dai *et al.* (2022) (see Supplementary Information).

- Therman Analysis, Rigadu Corporation, Tokyo 190-0000, Japan
 Department of Space Studies, Southwest Research Institute, Boulder, CO 80302,
- USA 31. Astromaterials Research and Exploration Science, NASA Johnson Space Center, Houston, TX 77058, USA
- Earth-System Sciences, Korea Polar Research Institute, Incheon 21990, Korea
- Centre de Recherches Pétrographiques et Géochimiques, CNRS Université de Lorraine, 54500 Nancy, France
- University of Science and Technology of China, School of Earth and Space Sciences, Anhui 230026, China
- 35. Department of Earth Sciences, Natural History Museum, London, SW7 5BD, UK
- 36. IIL, Hokkaido University, Sapporo 001-0021, Japan
- 37. Institute for Geochemistry and Petrology, Department of Earth Sciences, ETH Zurich, Zurich, Switzerland
- 38. Earth and Space Science, Osaka University, Osaka 560-0043, Japan
- Spectroscopy and Imaging, Japan Synchrotron Radiation Research Institute, Hyogo 679-5198 Japan
- 40. School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85281, USA
- 41. Geology, University of Maryland, College Park, MD 20742, USA
- 42. Graduate School of Natural Science and Technology, Okayama University, Okayama 700-8530, Japan
- 43. Earth and Planetary Sciences, University of California, Davis, CA 95616, USA
- Science and Engineering, National Museum of Nature and Science, Tsukuba 305-0005, Japan
- 45. Chemistry, Tokyo University of Science, Tokyo 162-8601, Japan
- 46. School of Earth Sciences and Engineering, Nanjing University, Nanjing 210023,
- China
- 47. Department of Earth Science, Tohoku University, Sendai, 980-8578, Japan
- Department of Earth and Planetary Sciences, Kyushu University, Fukuoka 819-0395, Japan
- 49. Earth and Planetary Systems Science Program, Hiroshima University, Higashi-Hiroshima, 739-8526, Japan
- 50. Kanagawa Institute of Technology, Atsugi 243-0292, Japan
- 51. UTokyo Organization for Planetary and Space Science, University of Tokyo, Tokyo 113-0033, Japan

Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Sorbonne Université, Museum National d'Histoire Naturelle, CNRS UMR 7590, IRD, 75005 Paris, France

^{29.} Thermal Analysis, Rigaku Corporation, Tokyo 196-8666, Japan

We report both the mass dependent deviation and the radiogenic ingrowth on ^{40}Ca from the decay of ^{40}K . For mass dependent deviation, the data are reported as $\delta^{x/y}Ca$:

$$\delta^{x/y} Ca = \left(\frac{({}^{x}Ca/{}^{y}Ca)_{sample}}{({}^{x}Ca/{}^{y}Ca)_{sRM915b}} - 1\right) \times 10^{3},$$

with x and y = 40, 42, 43 or 44. Since most of the published Ca isotope data are measured against the SRM 915a standard, the $\delta^{44/40}$ Ca values reported here are re-normalised to SRM915a to facilitate comparison.

The radiogenic ingrowth on ${}^{40}Ca$ is reported using the epsilon notation,

$$\epsilon^{40}Ca = \left(\frac{({}^{40}Ca/{}^{44}Ca)_{n \text{ sample}}}{({}^{40}Ca/{}^{44}Ca)_{n \text{ SRM915b}}} - 1\right) \times 10^4,$$

with $({}^{40}Ca/{}^{44}Ca)_n$ representing the ${}^{40}Ca/{}^{44}Ca$ ratio corrected from the mass dependent isotopic fractionation after being normalised to the ${}^{42}Ca/{}^{44}Ca$ ratio using the exponential law and ${}^{42}Ca/{}^{44}Ca = 0.31221$ (Russell *et al.*, 1978).

The effect of concentration mismatch on the Sapphire is more significant than on traditional MC-ICP-MS (Moynier *et al.*, 2021), and all the samples were analysed with Ca concentrations within 1 % of the standard.

Results and Discussion

The Ca isotopic compositions of the two Ryugu samples and the six CC are reported in Table 1, along with literature values for the chondrites where available. The radiogenic ingrowth on ⁴⁰Ca from ⁴⁰K decay was corrected using the K and Ca abundances of the samples (from Yokoyama *et al.*, 2022) and $\delta^{44/40}$ Ca (age corrected) ratios are also presented in Table 1. The following discussion focuses on these corrected mass dependent isotopic variations. The $\delta^{44/40}$ Ca difference between SRM915b and SRM915a is 0.72 ‰ (Heuser and Eisenhauer, 2008). Neither the Ryugu samples nor the meteorites analysed here show any ⁴⁰Ca anomalies (after age corrections), which is consistent with the literature (*e.g.*, Simon and DePaolo, 2010; Huang and Jacobsen, 2017).

In a plot of $\delta^{44/42}$ Ca *vs*. $\delta^{44/40}$ Ca (age corrected), the data fall along a mass dependent line, regardless of whether the slope for equilibrium fractionation (1/2.1 as shown in Fig. 1) or kinetic fractionation is used. Likewise, variations between $\delta^{44/42}$ Ca and $\delta^{44/43}$ Ca are mass dependent within error (Fig. S-2). Therefore, the Ca isotopic variations observed among the samples analysed primarily reflect mass dependent isotopic fractionation.

The meteorite data reported here are consistent with literature values (Fig. 2 and Table 1), but it should be noted that literature Ca isotopic values are variable, especially for Orgueil and Allende. The variability may reflect interlaboratory bias, but more likely it reflects isotopic heterogeneity at the sample scale analysed. This is particularly the case for Allende, which contains abundant calcium-aluminum-rich inclusions (CAIs). Since our Allende sample was obtained from the Smithsonian Museum's large batch of homogenised powder, and our measured Ca isotopic composition falls in the middle of the range previously reported, it is likely representative of the bulk (Fig. 2). For Orgueil, part of the interlaboratory variability may be controlled by the variable distribution of secondary phases produced by aqueous alteration since Ca may be isotopically fractionated during alteration and carbonate precipitation (Blättler and Higgins, 2017). However, none of the studies that report Ca isotopic data include the Ca contents of their Orgueil analyses. We report here the first $\delta^{44/40}$ Ca values for Tarda and



Figure 1 The $\delta^{44/42}$ Ca values plotted against $\delta^{44/40}$ Ca values for the various samples analysed in this study, including the Ryugu samples. All the samples fall on a mass dependent line within error.

Error bars represent 2 sigma standard deviation.



Figure 2 A comparison of age corrected $\delta^{44/40}$ Ca values for the samples analysed here (in colour, see Fig. 1) and from the literature (grey). Ryugu samples from Chambers A and C are similar within error and fall within the range defined by the CI chondrites. The literature data are from Table 1, BSE estimate from Kang *et al.* (2017) and chondrules data from Amsellem *et al.* (2017). Error bars are 2 x standard deviation.

Tagish Lake, which are within error of one another and overlap with Orgueil.

The two Ryugu samples have Ca isotopic compositions within error of one another ($\delta^{44/40}Ca = 0.58 \pm 0.03$ ‰ for C0108 and 0.55 ± 0.08 ‰ for A0106-A0107; uncertainties represent 2 s.e. for n = 5 and 6, respectively). They are also within the range of published $\delta^{44/40}Ca$ values for CIs (Fig. 2). Notably, Ryugu sample A0106-A0107 (Ca/Al ~ 1.9) has almost twice the amount of Ca compared to the average CI (Ca/Al ~ 1.06) (Fig. 3), and ~20 % more than Ryugu sample C0108 (Ca/Al ~ 1.55). If the excess Ca in A0106-A0107 is mainly stored in secondary carbonates (Yokoyama *et al.*, 2022 and our discussion below), then these carbonates must contribute significantly to the bulk Ca



Table 1 Calcium isotopic data from this study and literature (Simon and DePaolo, 2010; Valdes *et al.*, 2014; Amsellem *et al.*, 2017; Huang and Jacobsen, 2017). 2 s.d. = 2 x standard deviation and 2 s.e. = 2 x standard error ($2sd/\sqrt{n}$). n is number of measurements.

Sample names	$\delta^{40/44} Ca_{SRM915b}$	2 s.d.	$\delta^{42/44}$ Ca _{SRM915b}	2 s.d.	$\delta^{43/44}Ca_{SRM915b}$	2 s.d.	ε ⁴⁰ Ca	2 s.e	n	$\delta^{44/40}Ca_{SRM915a}$	$\delta^{44/40}$ Ca _{SRM915a} (age corrected)
Ryugu C0108	0.22	0.03	0.12	0.03	0.04	0.07	-0.17	0.25	5	0.50	0.58
Ryugu A0106-A0107	0.23	0.08	0.11	0.05	0.06	0.07	0.09	0.22	6	0.49	0.55
Murchison	0.12	0.07	0.02	0.07	0.01	0.10	0.85	0.46	5	0.60	0.67
Murchison (Valdes+)											0.84
Murchison (Huang+.)										0.72	
Allende	0.36	0.05	0.10	0.08	0.06	0.13	1.47	0.79	4	0.36	0.39
Allende (Simon+)							0.52	0.58	2	0.49	0.54
Allende (Valdes+)											0.55
Allende (Amsellem+)											0.26
Allende (Amsellem+)											0.10
Allende (Amsellem+)											0.44
Allende (Huang+)										0.28	
Alais	0.41	0.06	0.17	0.06	0.10	0.08	0.56	0.34	6	0.31	0.42
Tarda	0.34	0.06	0.15	0.07	0.05	0.07	0.25	0.37	6	0.38	0.45
Tagish Lake	0.30	0.03	0.13	0.03	0.06	0.07	0.33	0.26	6	0.42	0.48
Orgueil	0.30	0.04	0.13	0.02	0.06	0.08	0.44	0.40	4	0.42	0.48
Orgueil (Amsellem+)											0.45
Orgueil (Valdes+)											0.65
Orgueil (Huang+)										0.75	



Figure 3 $\delta^{44/40}$ Ca plotted against the Ca/Al ratio of the samples. The Ca/Al ratios of bulk Chamber A and Chamber C samples from Yokoyama *et al.* (2022) are used for Ryugu samples in this work and are taken from the average value in Barrat *et al.* (2012) for Orgueil, as it was not available for specific samples used here.

isotopic composition. Therefore, our results imply that the Ca isotopic composition of the Ryugu samples and CIs are not significantly modified by the dissolution of primary Ca-bearing phases and precipitation as carbonates during aqueous alteration. One reason for the similar Ca isotopic compositions in Ryugu samples and CIs is that their Ca isotopic compositions are mostly dominated by the high abundance and the composition of the carbonates.

Several studies have quantified the modal abundances of minerals in CIs, but there is no systematic study and consensus

on the Ca carriers. Calcium sulfates are usually not detected in CIs but can be present at up to ~1 vol. % (e.g., Endress and Bischoff, 1996; Morlok, et al., 2006). Even when present, they were suggested to have formed during terrestrial alteration (Gounelle and Zolensky, 2001). Ca-rich phosphates exist in CI chondrites (Morlok et al., 2006), but they appear to be quite rare (0–0.05 vol. %; King et al., 2015; Alfing et al., 2019) and therefore are unlikely to have a strong control on the total Ca budget. Thus, carbonates are the most likely major carriers of Ca in CIs (Endress and Bischoff, 1996; Morlok et al., 2006, Alfing et al., 2019). Scanning electron microscopic (SEM) analyses of 18 sections of CIs (including Orgueil) point to an average carbonate abundance of ~5 vol. % and the carbonates are dominated by dolomite (Endress and Bischoff, 1996). However, analyses of CO2 released by phosphoric acid dissolution of ~100 mg of Orgueil only returned ~0.1 wt. % of carbonate C (Alexander et al., 2015), which is equivalent to ~0.8 wt. % carbonate (although carbonate abundance in Ivuna estimated by a similar method is three times higher). X-ray diffraction (detection limit ~1 vol. %) did not reveal carbonates in three Orgueil samples (from 50 to 200 mg), but 2 vol. % in Alais (200 mg) and 3 vol. % in Ivuna (50 mg) (King et al., 2015), while Bland et al. (2004) detected no carbonates in Orgueil (200-300 mg samples). Given the variability in the modal mineralogy in the literature, Alfing et al. (2009) focused on phases >5 µm (which only represent ~6 vol. % of CIs) and found ~0.5 wt. % of carbonates in CIs. A variability in the abundance of carbonates in CIs is consistent with variable Ca concentrations (from 0.77 to 0.96 wt. %) measured even in large (0.5-1 g) Orgueil bulk samples (Barrat et al., 2012). Considering that the most abundant carbonates in Orgueil are dolomites with ~20 wt. % Ca (Endress and Bishoff, 1996), the presence of ~4 wt. % carbonates in CIs would be sufficient to dominate their Ca budget, less if calcites or aragonites are involved. Despite the variability in carbonate abundances of CIs reported in the literature, we suggest that the major fraction of Ca in CIs is stored in carbonates.

Nakamura, T. et al. (2022) estimated the mineral abundances and compositions of the main phases of the Ryugu samples by SEM observations of two ~10 mm² sections from a sample from the second touchdown site (sample C0002) (see their Tables S6 and S7). No Ca sulfates were found in these sections, and a simple mass balance using their data shows that carbonates account for 75-80 % of the Ca budget, with apatite and phyllosilicates accommodating the remaining Ca more or less equally. This calculation may underestimate the Ca fraction in carbonates because the samples also contain small Ca carbonate grains (e.g., Table S7 of Nakamura T. et al., 2022), the abundances of which could not be quantified here. C0002 is the third largest sample among all returned grains containing the major lithology (Nakamura T. et al., 2022), suggesting that the Ca budget in the Ryugu samples is dominated by carbonates. It should be noted that another study also found several vol. % of carbonate minerals in samples from both touchdown sites, with a large variability in Ca content between ~1 mg grains (Nakamura E. et al., 2022) and a correlation between Ca content and the dolomite abundances of the grains (Fig. S-1). The two heaviest Ca isotopic compositions for Orgueil samples from Valdes et al. (2014) (0.65 \pm 0.17 ‰, 2 s.d.) and Huang and Jacobsen (2017) (0.75 ± 0.11 ‰, 2 s.d.) may reflect different proportions of carbonates. Unfortunately, these two studies did not report the Ca contents of their Orgueil fractions, so it is not possible to test this hypothesis.

The similar Ca isotopic compositions between the two Ryugu samples and CIs are most simply explained if the Ca excesses observed in the bulk Ryugu samples are due to the heterogeneous distribution of carbonates, and if these carbonates have similar Ca isotopic compositions to the bulk samples. This explanation is consistent with an episode of fluid circulation and carbonate precipitation in the Ryugu samples that occurred 2.5 to 5 Myr after CAIs formation, as dated using ⁵³Mn-⁵³Cr chronometry in carbonate phases (Nakamura E. et al., 2022; Yokoyama et al., 2022). Hence, at present the average of the two Ryugu samples $(0.57 \pm 0.04 \%, 2 \text{ s.d.})$ represents the best estimate of Ryugu's and Solar System Ca isotopic composition. Future work should test whether this value is representative of the whole body by analysing other Ryugu fragments containing fewer carbonate phases and less total Ca, such as the Ryugu material in section 5 from C0002 (Nakamura T. et al., 2022), which only contains ~75 % of its Ca in carbonates.

Acknowledgements

This work was partly supported by the IPGP analytical platform PARI, Ile-de-France SESAME Grants 12015908, DIM ACAV+, the ERC grant 101001282 (METAL), Labex UnivEarth (FM), JSPS Kaken-hi grants (ST, HY, TY) and the CNES. We thank Dominik Hezel and one anonymous reviewer for constructive comments that helped improve the manuscript.

Editor: Anat Shahar

Additional Information

Supplementary Information accompanies this letter at https://www.geochemicalperspectivesletters.org/article2238.



© 2022 The Authors. This work is distributed under the Creative Commons Attribution Non-Commercial No-Derivatives 4.0

License, which permits unrestricted distribution provided the original author and source are credited. The material may not be adapted (remixed, transformed or built upon) or used for commercial purposes without written permission from the author. Additional information is available at https://www. geochemicalperspectivesletters.org/copyright-and-permissions.

Cite this letter as: Moynier, F., Dai, W., Yokoyama, T., Hu, Y., Paquet, M., Abe, Y., Aléon, J., Alexander, C.M.O'D., Amari, S., Amelin, Y., Bajo, K.-I., Bizzarro, M., Bouvier, A., Carlson, R.W., Chaussidon, M., Choi, B.-G., Dauphas, N., Davis, A.M., Di Rocco, T., Fujiya, W., Fukai, R., Gautam, I., Haba, M.K., Hibiya, Y., Hidaka, H., Homma, H., Hoppe, P., Huss, G.R., Ichida, K., Iizuka, T., Ireland, T.R., Ishikawa, A., Ito, M., Itoh, S., Kawasaki, N., Kita, N.T., Kitajima, K., Kleine, T., Komatani, S., Krot, A.N., Liu, M.-C., Masuda, Y., McKeegan, K.D., Morita, M., Motomura, K., Nakai, I., Nagashima, K., Nesvorný, D., Nguyen, A., Nittler, L., Onose, M., Pack, A., Park, C., Piani, L., Qin, L., Russell, S.S., Sakamoto, N., Schönbächler, M., Tafla, L., Tang, H., Terada, K., Terada, Y., Usui, T., Wada, S., Wadhwa, M., Walker, R.J., Yamashita, K., Yin, Q.-Z., Yoneda, S., Young, E.D., Yui, H., Zhang, A.-C., Nakamura, T., Naraoka, H., Noguchi, T., Okazaki, R., Sakamoto, K., Yabuta, H., Abe, M., Miyazaki, A., Nakato, A., Nishimura, M., Okada, T., Yada, T., Yogata, K., Nakazawa, S., Saiki, T., Tanaka, S., Terui, F., Tsuda, Y., Watanabe, S.-I., Yoshikawa, M., Tachibana, S., Yurimoto, H. (2022) The Solar System calcium isotopic composition inferred from Ryugu samples. Geochem. Persp. Let. 24, 1-6. https://doi.org/10.7185/ geochemlet.2238

References

- ALFING, J., PATSEK, M., BISCHOFF, A. (2019) Geochemistry Modal abundances of coarse-grained (>5µm) components within CI-chondrites and their individual clasts - Mixing of various lithologies on the CI parent body(ies). *Geochemistry* 39, 3–16. https://doi.org/10.1016/j.chemer.2019.08.004
- ALEXANDER, C.M.O'D., BOWDEN, R., FOGEL, M.L., HOWARD, K. (2015) Carbonate abundances and isotopic compositions in chondrites. *Meteoritics & Planetary Science* 50, 810–833. https://doi.org/10.1111/maps.12410
- AMSELLEM, E., MOYNIER, F., PRINGLE, E., BOUVIER, A., CHEN, H., DAY, J.M.D. (2017) Testing the chondrule-rich accretion model for planetary embryos using calcium isotopes. *Earth and Planetary Science Letters* 469, 75–83. https:// doi.org/10.1016/j.epsl.2017.04.022
- ARAKAWA, M., SAIKI, T., WADA, K., OGAWA, K., KADONO, T. et al. (2020) An artificial impact on the asteroid (162173) Ryugu formed a crater in the gravitydominated regime. Science 368, 67–71. https://doi.org/10.1126/science. aaz1701
- BARRAT, J.A., ZANDA, B., MOYNIER, F., BOLLINGER, C., LIORZOU, C. et al. (2012) Geochemistry of CI chondrites: Major and trace elements, and Cu and Zn Isotopes. Geochimica et Cosmochimica Acta 83, 79–92. https://doi.org/ 10.1016/j.gca.2011.12.011
- BLAND, P., CRESSEY, G., MENZIES, O. (2010) Modal mineralogy of carbonaceous chondrites by X-ray diffraction and Mossbauer spectroscopy. *Meteoritics and Planetary Science* 39, 3–16. https://doi.org/10.1111/j.1945-5100.2004. tb00046.x
- BLÄTTLER, C., HIGGINS, J. (2017) Testing Urey's carbonate–silicate cycle using the calcium isotopic composition of sedimentary carbonates. *Earth and Planetary Science Letters* 479, 241–251. https://doi.org/10.1016/j.epsl.2017.09.033
- DAI, W., MOYNIER, F., PAQUET, M., MOUREAU, J., DEBRET, B., SIEBERT, J., GERARD, Y., ZHAO, Y. (2021) Calcium isotope measurements using a collision cell (CC)-MC-ICP-MS. *Chemical Geology* 590, 120688. https://doi.org/10. 1016/j.chemgeo.2021.120688
- DAUPHAS, N., POURMAND, A. (2015) Thulium anomalies and rare earth element patterns in meteorites and Earth: Nebular fractionation and the nugget effect. *Geochimica et Cosmochimica Acta* 163, 234–261. https://doi.org/10.1016/j. gca.2015.03.037
- ENDRESS, M., BISHOFF, A. (1996) Carbonates in CI chondrites: Clues to parent body evolution. *Geochimica et Cosmochimica Acta* 60, 489–507. https://doi.org/10. 1016/0016-7037(95)00399-1
- FANTLE, M.S., TIPPER, E.T. (2014) Calcium isotopes in the global biogeochemical Ca cycle: Implications for development of a Ca isotope proxy. *Earth-Science Reviews* 129, 148–177. https://doi.org/10.1016/j.earscirev.2013.10.004

- GOUNELLE, M., ZOLENSKY, M. (2001) A terrestrial origin for sulfate veins in CI1 chondrites. *Meteoritics & Planetary Science*. 36, 1321–1329. https://doi.org/10. 1111/j.1945-5100.2001.tb01827.x
- HEUSER, A., EISENHAUER, A. (2008) The calcium isotope composition (delta Ca-44/40) of NIST SRM 915b and NIST SRM 1486. Geostandards and Geoanalytical Research 32, 311–315. https://doi.org/10.1111/j.1751-908X.2008.00877.x
- HEZEL, DC., RUSSELL, S.S., ROSS, A.J., KEARSLEY, A.T. (2008) Modal Abundances of CAIs: Implications for bulk chondrite element abundances and fractionations. *Meteoritics & Planetary Science* 43, 1879–1894. https://doi.org/10. 1111/j.1945-5100.2008.tb00649.x
- HUANG, S., JACOBSEN, S.B. (2017) Calcium isotopic compositions of chondrites. Geochimica et Cosmochimica Acta 201, 364–376. https://doi.org/10.1016/j. gca.2016.09.039
- HUANG, S., FARKAS, J., YU, G., PETAEV, M.I., JACOBSEN, S.B. (2012) Calcium isotopic ratios and rare earth elements abundances from refractory inclusions from the Allende CV3 chondrite. *Geochimica et Cosmochimica Acta* 77, 252–265. https://doi.org/10.1016/j.gca.2011.11.002
- KANG, J.T., IONOV, D.A., LIU, F., ZHANG, C.L., GOLOVIN, A.V., QIN, L.-P., ZHANG, Z.-F., HUANG, F. (2017) Calcium isotopic fractionation in mantle peridotites by melting and metasomatism and Ca isotope composition of the Bulk Silicate Earth. Earth and Planetary Science Letters 474, 128–137. https:// doi.org/10.1016/j.epsl.2017.05.035
- KING, A., SCHOFIELD, P.F., HOWARD, K., RUSSELL, S.S. (2015) Modal mineralogy of CI and CI-like chondrites by X-ray diffraction. *Geochimica et Cosmochimica Acta* 165, 148–160. https://doi.org/10.1016/j.gca.2015.05.038
- MORLOK, A., BISCHOFF, A., STEPHAN, T., FLOSS, C., ZINNER, E., JESSBERGER, E.K. (2006) Brecciation and chemical heterogeneities of CI chondrites. *Geochimica* et Cosmochimica Acta 70, 5371–5394. https://doi.org/10.1016/j.gca.2006.08. 007
- MOYNIER, F., HU, Y., WANG, K., ZHAO, Y., GÉRARD, Y., DENG, Z., MOUREAU, J., LI, W., SIMON, J.I., TENG, F.-Z. (2021) Potassium isotopic composition of various samples using a dual-path collision-cell-capable multiple-collector inductively coupled plasma mass spectrometer. *Chemical Geology* 571, 120144. https://doi.org/10.1016/j.chemgeo.2021.120144
- NAKAMURA, E., KOBAYASHI, K., TANAKA, R., KUNIHIRO, T., KITAGAWA, H. et al. (2022) On the origin and evolution of the asteroid Ryugu: A comprehensive geochemical perspective. Proceedings of the Japan Academy, Series B. 6, 227– 282. https://doi.org/10.2183/pjab.98.015
- NAKAMURA, T., MATSUMOTO, M., AMANO, K., ENOKIDO, Y., ZOLENSKY, M. et al. (2022) Formation and evolution of carbonaceous asteroid Ryugu: Direct evidence from returned samples. *Science*. https://doi.org/10.1126/science.abn8671
- NIEDERER, F.R., PAPANASTASSIOU, D.A. (1984) Ca isotopes in refractory inclusions. Geochimica et Cosmochimica Acta 48, 1279–1293. https://doi.org/10.1016/ 0016-7037(84)90062-0
- PALME, H., LODDERS, K., JONES, A. (2014) 2.2 Solar System abundances of the elements. In: HOLLAND, H.D., TUREKIAN, K.K. (Exec. Eds.) Treatise on Geochemistry (Second Edition). Elsevier, Oxford, 15–36. https://doi.org/10. 1016/B978-0-08-095975-7.00118-2
- RUSSELL, W.A., PAPANASTASSIOU, D.A., TOMBRELLO, T.A. (1978) Ca isotope fractionation on the Earth and other solar system materials. *Geochimica et Cosmochimica Acta* 42, 1075–1090. https://doi.org/10.1016/0016-7037(78) 90105-9
- SAIKI, S., IMAMURA, H., ARAKAWA, M., WADA, K., TAKAGI, Y., HAYAKAWA, M., SHIRAI, K., YANO, H., OKAMOTO, C. (2017) The Small Carry-on Impactor (SCI) and the Hayabusa2 Impact Experiment. Space Science Reviews 208, 165–186. https:// doi.org/10.1007/s11214-016-0297-5
- SIMON, J.I., DEPAOLO, D.J. (2010) Stable calcium isotopic composition of meteorites and rocky planets. *Earth and Planetary Science Letters* 289, 457–466. https:// doi.org/10.1016/j.epsl.2009.11.035
- SIMON, J.I., JORDAN, M.J., TAPPA, A., SCHAUBLE, E., KOHLL, E., YOUNG, E.D. (2017) Calcium and titanium isotope fractionation in refractory inclusions: Tracers of condensation and inheritance in the early solar system. *Earth* and Planetary Science Letters 472, 277–288. https://doi.org/10.1016/j.epsl. 2017.05.002
- TACHIBANA, S., SAWADA, H., OKAZAKI, R., TAKANO, Y., SAKAMOTO, K. et al. (2022) Pebbles and sand on asteroid (162173) Ryugu: In situ observation and particles returned to Earth. Science 375, 1011–1016. https://doi.org/10.1126/ science.abj8624
- VALDES, M., MOREIRA, M., FORIEL, J., MOYNIER, F. (2014) The nature of Earth's building blocks as revealed by calcium isotopes. *Earth and Planetary Science Letters* 394, 135–145. https://doi.org/10.1016/j.epsl.2014.02.052
- YADA, T., ABE, M., OKADA, T., NAKATO, A., YOGATA, K. *et al.* (2022) Preliminary analysis of the Hayabusa2 samples returned from C-type asteroid Ryugu. *Nature* Astronomy 6, 214–220. https://doi.org/10.1038/s41550-021-01550-6

YOKOYAMA, T., NAGASHIMA, K., NAKAI, I., YOUNG, E.D., ABE, Y. et al. (2022) Samples returned from the asteroid Ryugu are similar to Ivuna-type carbonaceous meteorites. *Science* https://doi.org/10.1126/science.abn7850