

# The Solar System calcium isotopic composition inferred from Ryugu samples

**Journal Article****Author(s):**

Moynier, Frédéric; Dai, W.; Yokoyama, T.; Hu, Y.; Paquet, Matthieu; Abe, Y.; Aléon, Jérôme; Alexander, Conel M.O'D.; Amari, Sachiko; Amelin, Yuri; Bajo, Ken-ichi; Bizzarro, Martin; Bouvier, Aurelie; Carlson, Robert W.; Chaussidon, M.; Choi, Byeon-Gak; Dauphas, Nicolas; Davis, Andrew M.; Di Rocco, Tommaso; Fujiya, Wataru; Schönbächler, Maria; et al.

**Publication date:**

2022

**Permanent link:**

<https://doi.org/10.3929/ethz-b-000581252>

**Rights / license:**

[Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International](#)

**Originally published in:**

Geochemical Perspectives Letters 24, <https://doi.org/10.7185/geochemlet.2238>


## The Solar System calcium isotopic composition inferred from Ryugu samples

F. Moynier<sup>1\*</sup>, W. Dai<sup>1</sup>, T. Yokoyama<sup>2</sup>, Y. Hu<sup>1</sup>, M. Paquet<sup>1</sup>, Y. Abe<sup>3</sup>, J. Aléon<sup>4</sup>, C.M.O'D. Alexander<sup>5</sup>, S. Amari<sup>6</sup>, Y. Amelin<sup>7</sup>, K.-I. Bajo<sup>8</sup>, M. Bizzarro<sup>9,1</sup>, A. Bouvier<sup>10</sup>, R.W. Carlson<sup>5</sup>, M. Chaussidon<sup>1</sup>, B.-G. Choi<sup>11</sup>, N. Dauphas<sup>12</sup>, A.M. Davis<sup>12</sup>, T. Di Rocco<sup>13</sup>, W. Fujiya<sup>14</sup>, R. Fukai<sup>15</sup>, I. Gautam<sup>2</sup>, M.K. Haba<sup>2</sup>, Y. Hibiya<sup>16</sup>, H. Hidaka<sup>17</sup>, H. Homma<sup>18</sup>, P. Hoppe<sup>19</sup>, G.R. Huss<sup>20</sup>, K. Ichida<sup>21</sup>, T. Iizuka<sup>22</sup>, T.R. Ireland<sup>23</sup>, A. Ishikawa<sup>2</sup>, M. Ito<sup>24</sup>, S. Itoh<sup>25</sup>, N. Kawasaki<sup>8</sup>, N.T. Kita<sup>26</sup>, K. Kitajima<sup>26</sup>, T. Kleine<sup>27</sup>, S. Komatani<sup>21</sup>, A.N. Krot<sup>20</sup>, M.-C. Liu<sup>28</sup>, Y. Masuda<sup>2</sup>, K.D. McKeegan<sup>28</sup>, M. Morita<sup>21</sup>, K. Motomura<sup>29</sup>, I. Nakai<sup>29</sup>, K. Nagashima<sup>20</sup>, D. Nesvorný<sup>30</sup>, A. Nguyen<sup>31</sup>, L. Nittler<sup>5</sup>, M. Onose<sup>21</sup>, A. Pack<sup>13</sup>, C. Park<sup>32</sup>, L. Piani<sup>33</sup>, L. Qin<sup>34</sup>, S.S. Russell<sup>35</sup>, N. Sakamoto<sup>36</sup>, M. Schönbachler<sup>37</sup>, L. Tafla<sup>28</sup>, H. Tang<sup>28</sup>, K. Terada<sup>38</sup>, Y. Terada<sup>39</sup>, T. Usui<sup>15</sup>, S. Wada<sup>8</sup>, M. Wadhwa<sup>40</sup>, R.J. Walker<sup>41</sup>, K. Yamashita<sup>42</sup>, Q.-Z. Yin<sup>43</sup>, S. Yoneda<sup>44</sup>, E.D. Young<sup>28</sup>, H. Yui<sup>45</sup>, A.-C. Zhang<sup>46</sup>, T. Nakamura<sup>47</sup>, H. Naraoka<sup>48</sup>, T. Noguchi<sup>24</sup>, R. Okazaki<sup>48</sup>, K. Sakamoto<sup>15</sup>, H. Yabuta<sup>49</sup>, M. Abe<sup>15</sup>, A. Miyazaki<sup>15</sup>, A. Nakato<sup>15</sup>, M. Nishimura<sup>15</sup>, T. Okada<sup>15</sup>, T. Yada<sup>15</sup>, K. Yogata<sup>15</sup>, S. Nakazawa<sup>15</sup>, T. Saiki<sup>15</sup>, S. Tanaka<sup>15</sup>, F. Terui<sup>51</sup>, Y. Tsuda<sup>15</sup>, S.-I. Watanabe<sup>17</sup>, M. Yoshikawa<sup>15</sup>, S. Tachibana<sup>51</sup>, H. Yurimoto<sup>8</sup>



<https://doi.org/10.7185/geochemlet.2238>

### Abstract



The Hayabusa2 spacecraft has returned samples from the Cb-type asteroid (162173) Ryugu to Earth. Previous petrological and chemical analyses support a close link between Ryugu and CI chondrites that are presumed to be chemically the most primitive meteorites with a solar-like composition. However, Ryugu samples are highly enriched in Ca compared to typical CI chondrites. To identify the cause of this discrepancy, here we report stable Ca isotopic data (expressed as  $\delta^{44/40}\text{Ca}_{\text{SRM915a}}$ ) for returned Ryugu samples collected from two sites. We found that samples from both sites have similar  $\delta^{44/40}\text{Ca}_{\text{SRM915a}}$  ( $0.58 \pm 0.03 \text{ ‰}$  and  $0.55 \pm 0.08 \text{ ‰}$ , 2 s.d.) that fall within the range defined by CIs. This isotopic similarity suggests that the Ca budget of CIs and Ryugu samples is dominated by carbonates, and the variably higher Ca contents in Ryugu samples are due to the abundant carbonates. Precipitation of carbonates on Ryugu likely coincided with a major episode of aqueous activity dated to have occurred  $\sim 5$  Myr after Solar System formation. Based on the pristine Ryugu samples, the average  $\delta^{44/40}\text{Ca}_{\text{SRM915a}}$  of the Solar System is defined to be  $0.57 \pm 0.04 \text{ ‰}$  (2 s.d.).

Received 28 June 2022 | Accepted 19 September 2022 | Published 19 October 2022

### Introduction

In December 2020, the JAXA Hayabusa2 spacecraft returned to Earth with the first samples collected from a Cb-type asteroid, (162173) Ryugu (Tachibana *et al.*, 2022; Yada *et al.*, 2022). Chemical, mineralogical, petrological, and isotopic analyses of these samples suggest that they are closely related to CI

chondrites. In particular, their bulk Cr and Ti isotopic signatures and the chemical abundances of most elements are within the range of CI chondrites (Nakamura *E. et al.*, 2022; Yokoyama *et al.*, 2022). Among meteorites, CI chondrites have chemical compositions that most closely resemble the Sun; therefore, they are the most representative samples of the solar nebula composition with the exception of volatile elements (Palme *et al.*, 2014).

1. Université Paris Cité, Institut de Physique du Globe de Paris, CNRS, 75005 Paris, France  
2. Department of Earth and Planetary Sciences, Tokyo Institute of Technology, Tokyo 152-8551, Japan  
3. Graduate School of Engineering Materials Science and Engineering, Tokyo Denki University, Tokyo 120-8551, Japan  
\* Corresponding author (email: moynier@ipgp.fr)



Given that the Ryugu samples did not experience any terrestrial alteration, they are likely the chemically most pristine Solar System samples available estimating the original isotopic compositions of most elements in the Solar System (Yokoyama *et al.*, 2022).

Despite many aspects of similarities with the CIs, the Ryugu samples display an apparent excess of Ca by over 50 %, which may be related to a heterogeneous distribution of carbonates (dolomite and calcite) (Nakamura *T. et al.*, 2022; Yokoyama *et al.*, 2022) between Ryugu, Orgueil and other CIs. Calcium is a major constituent of carbonates and can be isotopically fractionated during aqueous alteration and carbonate precipitation, leading to more than 1 ‰ variations in the <sup>44</sup>Ca/<sup>40</sup>Ca ratio in terrestrial carbonates (*e.g.*, Fantle and Tipper, 2014; Blättler and Higgins, 2017). In addition, Ca exhibits large isotopic variations among bulk carbonaceous chondrites (CC), with the <sup>44</sup>Ca/<sup>40</sup>Ca ratio spanning a range of 1 ‰. This range is likely related to a combination of the variable modal abundances of refractory inclusions among CC (Hezel *et al.*, 2008) that can be enriched in the lighter Ca isotopes by several per mille (Niederer and Papanastassiou, 1984; Huang *et al.*, 2012) and the heterogeneous distribution of carbonates (Simon and DePaolo, 2010; Valdes *et al.*, 2014; Dauphas and Pourmand, 2015). Therefore, stable Ca isotopes could be useful for investigating the origin of Ca excess in Ryugu samples compared to CIs.

Here we have analysed the stable Ca isotopic compositions of Ryugu samples collected from the first and second touchdown sites, using the collision cell equipped multicollection inductively-coupled plasma mass spectrometer (CC-MC-ICP-MS), Nu Sapphire.

## Samples and Methods

The samples returned by the Hayabusa2 spacecraft consist of ~5 g of materials from the Ryugu asteroid recovered during two touchdowns (Tachibana *et al.*, 2022; Yada *et al.*, 2022). Approximately 3 g of samples representing the surface materials of Ryugu were collected during the first touchdown and stored in Chamber A. Approximately 2 g of samples likely representing a mixture of materials from the surface and subsurface were collected into Chamber C at a site that was close to the crater formed by the Small Carry-on Impactor, a kinetic impact experiment of the Hayabusa2 mission (Saiki *et al.*, 2017; Arakawa *et al.*, 2020). Two Ryugu samples, A0106-A0107 (Chamber A) and C0108 (Chamber C), were analysed in this study (for information on the mineralogy see [https://jaxa.repo.nii.ac.jp/?action=repository\\_uri&item\\_id=48255&file\\_id=31&file\\_no=1](https://jaxa.repo.nii.ac.jp/?action=repository_uri&item_id=48255&file_id=31&file_no=1)).

Sample A0106-A0107 was prepared from a mixed aggregate of A0106 (1.6 mg) and A0107 (27.3 mg). In addition to the Ryugu samples, fusion-crust free bulk samples of six CC, Orgueil (CI1), Alais (CI1), Tarda (C2-ungrouped), Tagish Lake (C2-ungrouped), Murchison (CM2), and Allende (CV3), were analysed in the same way for comparison. See Table S-1 for the weights and providers of the meteorite samples. All samples were dissolved in PFA vials with a mixture of concentrated HF and HNO<sub>3</sub> at the Tokyo Institute of Technology (Yokoyama *et al.*, 2022).

After dissolution, aliquots of ~0.15 % of the solutions containing ~5 µg of Ca were transferred and dedicated for our study. All the sample aliquots were dried and redissolved in 0.4 ml of 4 mol/L HNO<sub>3</sub> in preparation for Ca chemical purification and isotopic measurements at the Institut de Physique du Globe, following Dai *et al.* (2022) (see Supplementary Information).

4. Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Sorbonne Université, Museum National d'Histoire Naturelle, CNRS UMR 7590, IRD, 75005 Paris, France
5. Earth and Planets Laboratory, Carnegie Institution for Science, Washington, DC, 20015, USA
6. McDonnell Center for the Space Sciences and Physics Department, Washington University, St. Louis, MO 63130, USA
7. Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, GD 510640, China
8. Natural History Sciences, IIL, Hokkaido University, Sapporo 001-0021, Japan
9. Centre for Star and Planet Formation, GLOBE Institute, University of Copenhagen, Copenhagen, K 1350, Denmark
10. Bayerisches Geoinstitut, Universität Bayreuth, Bayreuth 95447, Germany
11. Department of Earth Science Education, Seoul National University, Seoul 08826, Republic of Korea
12. Department of the Geophysical Sciences and Enrico Fermi Institute, The University of Chicago, 5734 South Ellis Avenue, Chicago 60637, USA
13. Faculty of Geosciences and Geography, University of Göttingen, Göttingen, D-37077, Germany
14. Faculty of Science, Ibaraki University, Mito 310-8512, Japan
15. ISAS/JSEC, JAXA, Sagami-hara 252-5210, Japan
16. General Systems Studies, The University of Tokyo, Tokyo 153-0041, Japan
17. Earth and Planetary Sciences, Nagoya University, Nagoya 464-8601, Japan
18. Osaka Application Laboratory, SBUWDX, Rigaku Corporation, Osaka 569-1146, Japan
19. Max Planck Institute for Chemistry, Mainz 55128, Germany.
20. Hawai'i Institute of Geophysics and Planetology, University of Hawai'i at Mānoa, Honolulu, HI 96822, USA
21. Analytical Technology, Horiba Techno Service Co., Ltd., Kyoto 601-8125, Japan
22. Earth and Planetary Science, The University of Tokyo, Tokyo 113-0033, Japan
23. School of Earth and Environmental Sciences, The University of Queensland, St Lucia QLD 4072, Australia
24. Kochi Institute for Core Sample Research, JAMSTEC, Kochi 783-8502, Japan
25. Earth and Planetary Sciences, Kyoto University, Kyoto 606-8502, Japan
26. Geoscience, University of Wisconsin-Madison, Madison, WI 53706, USA
27. Max Planck Institute for Solar System Research; 37077 Göttingen, Germany
28. Earth, Planetary, and Space Sciences, UCLA, Los Angeles, CA 90095, USA
29. Thermal Analysis, Rigaku Corporation, Tokyo 196-8666, Japan
30. Department of Space Studies, Southwest Research Institute, Boulder, CO 80302, USA
31. Astromaterials Research and Exploration Science, NASA Johnson Space Center, Houston, TX 77058, USA
32. Earth-System Sciences, Korea Polar Research Institute, Incheon 21990, Korea
33. Centre de Recherches Pétrographiques et Géochimiques, CNRS - Université de Lorraine, 54500 Nancy, France
34. University of Science and Technology of China, School of Earth and Space Sciences, Anhui 230026, China
35. Department of Earth Sciences, Natural History Museum, London, SW7 5BD, UK
36. IIL, Hokkaido University, Sapporo 001-0021, Japan
37. Institute for Geochemistry and Petrology, Department of Earth Sciences, ETH Zurich, Zurich, Switzerland
38. Earth and Space Science, Osaka University, Osaka 560-0043, Japan
39. Spectroscopy and Imaging, Japan Synchrotron Radiation Research Institute, Hyogo 679-5198 Japan
40. School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85281, USA
41. Geology, University of Maryland, College Park, MD 20742, USA
42. Graduate School of Natural Science and Technology, Okayama University, Okayama 700-8530, Japan
43. Earth and Planetary Sciences, University of California, Davis, CA 95616, USA
44. Science and Engineering, National Museum of Nature and Science, Tsukuba 305-0005, Japan
45. Chemistry, Tokyo University of Science, Tokyo 162-8601, Japan
46. School of Earth Sciences and Engineering, Nanjing University, Nanjing 210023, China
47. Department of Earth Science, Tohoku University, Sendai, 980-8578, Japan
48. Department of Earth and Planetary Sciences, Kyushu University, Fukuoka 819-0395, Japan
49. Earth and Planetary Systems Science Program, Hiroshima University, Higashi-Hiroshima, 739-8526, Japan
50. Kanagawa Institute of Technology, Atsugi 243-0292, Japan
51. UTokyo Organization for Planetary and Space Science, University of Tokyo, Tokyo 113-0033, Japan



We report both the mass dependent deviation and the radiogenic ingrowth on  $^{40}\text{Ca}$  from the decay of  $^{40}\text{K}$ . For mass dependent deviation, the data are reported as  $\delta^{x/y}\text{Ca}$ :

$$\delta^{x/y}\text{Ca} = \left( \frac{(^x\text{Ca}/^y\text{Ca})_{\text{sample}}}{(^x\text{Ca}/^y\text{Ca})_{\text{SRM915b}}} - 1 \right) \times 10^3,$$

with  $x$  and  $y = 40, 42, 43$  or  $44$ . Since most of the published Ca isotope data are measured against the SRM 915a standard, the  $\delta^{44/40}\text{Ca}$  values reported here are re-normalised to SRM915a to facilitate comparison.

The radiogenic ingrowth on  $^{40}\text{Ca}$  is reported using the epsilon notation,

$$\epsilon^{40}\text{Ca} = \left( \frac{(^{40}\text{Ca}/^{44}\text{Ca})_{\text{sample}}}{(^{40}\text{Ca}/^{44}\text{Ca})_{\text{SRM915b}}} - 1 \right) \times 10^4,$$

with  $(^{40}\text{Ca}/^{44}\text{Ca})_{\text{n}}$  representing the  $^{40}\text{Ca}/^{44}\text{Ca}$  ratio corrected from the mass dependent isotopic fractionation after being normalised to the  $^{42}\text{Ca}/^{44}\text{Ca}$  ratio using the exponential law and  $^{42}\text{Ca}/^{44}\text{Ca} = 0.31221$  (Russell *et al.*, 1978).


The effect of concentration mismatch on the Sapphire is more significant than on traditional MC-ICP-MS (Moynier *et al.*, 2021), and all the samples were analysed with Ca concentrations within 1 % of the standard.

## Results and Discussion


The Ca isotopic compositions of the two Ryugu samples and the six CC are reported in Table 1, along with literature values for the chondrites where available. The radiogenic ingrowth on  $^{40}\text{Ca}$  from  $^{40}\text{K}$  decay was corrected using the K and Ca abundances of the samples (from Yokoyama *et al.*, 2022) and  $\delta^{44/40}\text{Ca}$  (age corrected) ratios are also presented in Table 1. The following discussion focuses on these corrected mass dependent isotopic variations. The  $\delta^{44/40}\text{Ca}$  difference between SRM915b and SRM915a is 0.72 ‰ (Heuser and Eisenhauer, 2008). Neither the Ryugu samples nor the meteorites analysed here show any  $^{40}\text{Ca}$  anomalies (after age corrections), which is consistent with the literature (e.g., Simon and DePaolo, 2010; Huang and Jacobsen, 2017).

In a plot of  $\delta^{44/42}\text{Ca}$  vs.  $\delta^{44/40}\text{Ca}$  (age corrected), the data fall along a mass dependent line, regardless of whether the slope for equilibrium fractionation (1/2.1 as shown in Fig. 1) or kinetic fractionation is used. Likewise, variations between  $\delta^{44/42}\text{Ca}$  and  $\delta^{44/43}\text{Ca}$  are mass dependent within error (Fig. S-2). Therefore, the Ca isotopic variations observed among the samples analysed primarily reflect mass dependent isotopic fractionation.

The meteorite data reported here are consistent with literature values (Fig. 2 and Table 1), but it should be noted that literature Ca isotopic values are variable, especially for Orgueil and Allende. The variability may reflect interlaboratory bias, but more likely it reflects isotopic heterogeneity at the sample scale analysed. This is particularly the case for Allende, which contains abundant calcium-aluminum-rich inclusions (CAIs). Since our Allende sample was obtained from the Smithsonian Museum's large batch of homogenised powder, and our measured Ca isotopic composition falls in the middle of the range previously reported, it is likely representative of the bulk (Fig. 2). For Orgueil, part of the interlaboratory variability may be controlled by the variable distribution of secondary phases produced by aqueous alteration since Ca may be isotopically fractionated during alteration and carbonate precipitation (Blättler and Higgins, 2017). However, none of the studies that report Ca isotopic data include the Ca contents of their Orgueil analyses. We report here the first  $\delta^{44/40}\text{Ca}$  values for Tarda and



**Figure 1** The  $\delta^{44/42}\text{Ca}$  values plotted against  $\delta^{44/40}\text{Ca}$  values for the various samples analysed in this study, including the Ryugu samples. All the samples fall on a mass dependent line within error. Error bars represent 2 sigma standard deviation.




**Figure 2** A comparison of age corrected  $\delta^{44/40}\text{Ca}$  values for the samples analysed here (in colour, see Fig. 1) and from the literature (grey). Ryugu samples from Chambers A and C are similar within error and fall within the range defined by the CI chondrites. The literature data are from Table 1, BSE estimate from Kang *et al.* (2017) and chondrules data from Amsellem *et al.* (2017). Error bars are 2 x standard deviation.

Tagish Lake, which are within error of one another and overlap with Orgueil.

The two Ryugu samples have Ca isotopic compositions within error of one another ( $\delta^{44/40}\text{Ca} = 0.58 \pm 0.03$  ‰ for C0108 and  $0.55 \pm 0.08$  ‰ for A0106-A0107; uncertainties represent 2 s.e. for  $n = 5$  and  $6$ , respectively). They are also within the range of published  $\delta^{44/40}\text{Ca}$  values for CIs (Fig. 2). Notably, Ryugu sample A0106-A0107 (Ca/Al  $\sim 1.9$ ) has almost twice the amount of Ca compared to the average CI (Ca/Al  $\sim 1.06$ ) (Fig. 3), and  $\sim 20$  % more than Ryugu sample C0108 (Ca/Al  $\sim 1.55$ ). If the excess Ca in A0106-A0107 is mainly stored in secondary carbonates (Yokoyama *et al.*, 2022 and our discussion below), then these carbonates must contribute significantly to the bulk Ca

**Table 1** Calcium isotopic data from this study and literature (Simon and DePaolo, 2010; Valdes *et al.*, 2014; Amsellem *et al.*, 2017; Huang and Jacobsen, 2017). 2 s.d. = 2 x standard deviation and 2 s.e. = 2 x standard error (2sd/ $\sqrt{n}$ ). n is number of measurements.

Sample names	$\delta^{40/44}\text{Ca}_{\text{SRM915b}}$	2 s.d.	$\delta^{42/44}\text{Ca}_{\text{SRM915b}}$	2 s.d.	$\delta^{43/44}\text{Ca}_{\text{SRM915b}}$	2 s.d.	$\epsilon^{40}\text{Ca}$	2 s.e.	n	$\delta^{44/40}\text{Ca}_{\text{SRM915a}}$	$\delta^{44/40}\text{Ca}_{\text{SRM915a}}$ (age corrected)
Ryugu C0108	0.22	0.03	0.12	0.03	0.04	0.07	-0.17	0.25	5	0.50	0.58
Ryugu A0106-A0107	0.23	0.08	0.11	0.05	0.06	0.07	0.09	0.22	6	0.49	0.55
Murchison	0.12	0.07	0.02	0.07	0.01	0.10	0.85	0.46	5	0.60	0.67
Murchison (Valdes+)											0.84
Murchison (Huang+)										0.72	
Allende	0.36	0.05	0.10	0.08	0.06	0.13	1.47	0.79	4	0.36	0.39
Allende (Simon+)							0.52	0.58	2	0.49	0.54
Allende (Valdes+)											0.55
Allende (Amsellem+)											0.26
Allende (Amsellem+)											0.10
Allende (Amsellem+)											0.44
Allende (Huang+)										0.28	
Alais	0.41	0.06	0.17	0.06	0.10	0.08	0.56	0.34	6	0.31	0.42
Tarda	0.34	0.06	0.15	0.07	0.05	0.07	0.25	0.37	6	0.38	0.45
Tagish Lake	0.30	0.03	0.13	0.03	0.06	0.07	0.33	0.26	6	0.42	0.48
Orgueil	0.30	0.04	0.13	0.02	0.06	0.08	0.44	0.40	4	0.42	0.48
Orgueil (Amsellem+)											0.45
Orgueil (Valdes+)											0.65
Orgueil (Huang+)										0.75	

**Figure 3**  $\delta^{44/40}\text{Ca}$  plotted against the Ca/Al ratio of the samples. The Ca/Al ratios of bulk Chamber A and Chamber C samples from Yokoyama *et al.* (2022) are used for Ryugu samples in this work and are taken from the average value in Barrat *et al.* (2012) for Orgueil, as it was not available for specific samples used here.

isotopic composition. Therefore, our results imply that the Ca isotopic composition of the Ryugu samples and CIs are not significantly modified by the dissolution of primary Ca-bearing phases and precipitation as carbonates during aqueous alteration. One reason for the similar Ca isotopic compositions in Ryugu samples and CIs is that their Ca isotopic compositions are mostly dominated by the high abundance and the composition of the carbonates.

Several studies have quantified the modal abundances of minerals in CIs, but there is no systematic study and consensus

on the Ca carriers. Calcium sulfates are usually not detected in CIs but can be present at up to ~1 vol. % (*e.g.*, Endress and Bischoff, 1996; Morlok *et al.*, 2006). Even when present, they were suggested to have formed during terrestrial alteration (Gounelle and Zolensky, 2001). Ca-rich phosphates exist in CI chondrites (Morlok *et al.*, 2006), but they appear to be quite rare (0–0.05 vol. %; King *et al.*, 2015; Alving *et al.*, 2019) and therefore are unlikely to have a strong control on the total Ca budget. Thus, carbonates are the most likely major carriers of Ca in CIs (Endress and Bischoff, 1996; Morlok *et al.*, 2006, Alving *et al.*, 2019). Scanning electron microscopic (SEM) analyses of 18 sections of CIs (including Orgueil) point to an average carbonate abundance of ~5 vol. % and the carbonates are dominated by dolomite (Endress and Bischoff, 1996). However, analyses of  $\text{CO}_2$  released by phosphoric acid dissolution of ~100 mg of Orgueil only returned ~0.1 wt. % of carbonate C (Alexander *et al.*, 2015), which is equivalent to ~0.8 wt. % carbonate (although carbonate abundance in Ivuna estimated by a similar method is three times higher). X-ray diffraction (detection limit ~1 vol. %) did not reveal carbonates in three Orgueil samples (from 50 to 200 mg), but 2 vol. % in Alais (200 mg) and 3 vol. % in Ivuna (50 mg) (King *et al.*, 2015), while Bland *et al.* (2004) detected no carbonates in Orgueil (200–300 mg samples). Given the variability in the modal mineralogy in the literature, Alving *et al.* (2009) focused on phases >5  $\mu\text{m}$  (which only represent ~6 vol. % of CIs) and found ~0.5 wt. % of carbonates in CIs. A variability in the abundance of carbonates in CIs is consistent with variable Ca concentrations (from 0.77 to 0.96 wt. %) measured even in large (0.5–1 g) Orgueil bulk samples (Barrat *et al.*, 2012). Considering that the most abundant carbonates in Orgueil are dolomites with ~20 wt. % Ca (Endress and Bischoff, 1996), the presence of ~4 wt. % carbonates in CIs would be sufficient to dominate their Ca budget, less if calcites or aragonites are involved. Despite the variability in carbonate abundances of CIs reported in the literature, we suggest that the major fraction of Ca in CIs is stored in carbonates.

Nakamura, T. *et al.* (2022) estimated the mineral abundances and compositions of the main phases of the Ryugu samples by SEM observations of two ~10 mm<sup>2</sup> sections from a sample from the second touchdown site (sample C0002) (see their Tables S6 and S7). No Ca sulfates were found in these sections, and a simple mass balance using their data shows that carbonates account for 75–80 % of the Ca budget, with apatite and phyllosilicates accommodating the remaining Ca more or less equally. This calculation may underestimate the Ca fraction in carbonates because the samples also contain small Ca carbonate grains (*e.g.*, Table S7 of Nakamura T. *et al.*, 2022), the abundances of which could not be quantified here. C0002 is the third largest sample among all returned grains containing the major lithology (Nakamura T. *et al.*, 2022), suggesting that the Ca budget in the Ryugu samples is dominated by carbonates. It should be noted that another study also found several vol. % of carbonate minerals in samples from both touchdown sites, with a large variability in Ca content between ~1 mg grains (Nakamura E. *et al.*, 2022) and a correlation between Ca content and the dolomite abundances of the grains (Fig. S-1). The two heaviest Ca isotopic compositions for Orgueil samples from Valdes *et al.* (2014) ( $0.65 \pm 0.17$  ‰, 2 s.d.) and Huang and Jacobsen (2017) ( $0.75 \pm 0.11$  ‰, 2 s.d.) may reflect different proportions of carbonates. Unfortunately, these two studies did not report the Ca contents of their Orgueil fractions, so it is not possible to test this hypothesis.

The similar Ca isotopic compositions between the two Ryugu samples and CIs are most simply explained if the Ca excesses observed in the bulk Ryugu samples are due to the heterogeneous distribution of carbonates, and if these carbonates have similar Ca isotopic compositions to the bulk samples. This explanation is consistent with an episode of fluid circulation and carbonate precipitation in the Ryugu samples that occurred 2.5 to 5 Myr after CAIs formation, as dated using <sup>53</sup>Mn-<sup>53</sup>Cr chronometry in carbonate phases (Nakamura E. *et al.*, 2022; Yokoyama *et al.*, 2022). Hence, at present the average of the two Ryugu samples ( $0.57 \pm 0.04$  ‰, 2 s.d.) represents the best estimate of Ryugu's and Solar System Ca isotopic composition. Future work should test whether this value is representative of the whole body by analysing other Ryugu fragments containing fewer carbonate phases and less total Ca, such as the Ryugu material in section 5 from C0002 (Nakamura T. *et al.*, 2022), which only contains ~75 % of its Ca in carbonates.

## Acknowledgements

This work was partly supported by the IPGP analytical platform PARI, Ile-de-France SESAME Grants 12015908, DIM ACAV+, the ERC grant 101001282 (METAL), Labex UnivEarth (FM), JSPS Kaken-hi grants (ST, HY, TY) and the CNES. We thank Dominik Hezel and one anonymous reviewer for constructive comments that helped improve the manuscript.

Editor: Anat Shahar

## Additional Information

Supplementary Information accompanies this letter at <https://www.geochemicalperspectivesletters.org/article2238>.



© 2022 The Authors. This work is distributed under the Creative Commons Attribution Non-Commercial No-Derivatives 4.0

License, which permits unrestricted distribution provided the original author and source are credited. The material may not be adapted (remixed, transformed or built upon) or used for

commercial purposes without written permission from the author. Additional information is available at <https://www.geochemicalperspectivesletters.org/copyright-and-permissions>.

**Cite this letter as:** Moynier, F., Dai, W., Yokoyama, T., Hu, Y., Paquet, M., Abe, Y., Aléon, J., Alexander, C.M.O'D., Amari, S., Amelin, Y., Bajo, K.-I., Bizzarro, M., Bouvier, A., Carlson, R.W., Chaussidon, M., Choi, B.-G., Dauphas, N., Davis, A.M., Di Rocco, T., Fujiya, W., Fukai, R., Gautam, I., Haba, M.K., Hibiy, Y., Hidaka, H., Homma, H., Hoppe, P., Huss, G.R., Ichida, K., Iizuka, T., Ireland, T.R., Ishikawa, A., Ito, M., Itoh, S., Kawasaki, N., Kita, N.T., Kitajima, K., Kleine, T., Komatani, S., Krot, A.N., Liu, M.-C., Masuda, Y., McKeegan, K.D., Morita, M., Motomura, K., Nakai, I., Nagashima, K., Nesvorný, D., Nguyen, A., Nittler, L., Onose, M., Pack, A., Park, C., Piani, L., Qin, L., Russell, S.S., Sakamoto, N., Schönbachler, M., Tafla, L., Tang, H., Terada, K., Terada, Y., Usui, T., Wada, S., Wadhwa, M., Walker, R.J., Yamashita, K., Yin, Q.-Z., Yoneda, S., Young, E.D., Yui, H., Zhang, A.-C., Nakamura, T., Naraoka, H., Noguchi, T., Okazaki, R., Sakamoto, K., Yabuta, H., Abe, M., Miyazaki, A., Nakato, A., Nishimura, M., Okada, T., Yada, T., Yogata, K., Nakazawa, S., Saiki, T., Tanaka, S., Terui, F., Tsuda, Y., Watanabe, S.-I., Yoshikawa, M., Tachibana, S., Yurimoto, H. (2022) The Solar System calcium isotopic composition inferred from Ryugu samples. *Geochem. Persp. Let.* 24, 1–6. <https://doi.org/10.7185/geochemlet.2238>

## References

- ALFING, J., PATSEK, M., BISCHOFF, A. (2019) Geochemistry Modal abundances of coarse-grained (>5µm) components within CI-chondrites and their individual clasts - Mixing of various lithologies on the CI parent body(ies). *Geochemistry* 39, 3–16. <https://doi.org/10.1016/j.chemer.2019.08.004>
- ALEXANDER, C.M.O'D., BOWDEN, R., FOGEL, M.L., HOWARD, K. (2015) Carbonate abundances and isotopic compositions in chondrites. *Meteoritics & Planetary Science* 50, 810–833. <https://doi.org/10.1111/maps.12410>
- AMSELEM, E., MOYNIER, F., PRINGLE, E., BOUVIER, A., CHEN, H., DAY, J.M.D. (2017) Testing the chondrule-rich accretion model for planetary embryos using calcium isotopes. *Earth and Planetary Science Letters* 469, 75–83. <https://doi.org/10.1016/j.epsl.2017.04.022>
- ARAKAWA, M., SAIKI, T., WADA, K., OGAWA, K., KADONO, T. *et al.* (2020) An artificial impact on the asteroid (162173) Ryugu formed a crater in the gravity-dominated regime. *Science* 368, 67–71. <https://doi.org/10.1126/science.aaz1701>
- BARRAT, J.A., ZANDA, B., MOYNIER, F., BOLLINGER, C., LIORZOU, C. *et al.* (2012) Geochemistry of CI chondrites: Major and trace elements, and Cu and Zn Isotopes. *Geochimica et Cosmochimica Acta* 83, 79–92. <https://doi.org/10.1016/j.gca.2011.12.011>
- BLAND, P., CRESSEY, G., MENZIES, O. (2010) Modal mineralogy of carbonaceous chondrites by X-ray diffraction and Mossbauer spectroscopy. *Meteoritics and Planetary Science* 39, 3–16. <https://doi.org/10.1111/j.1945-5100.2004.tb00046.x>
- BLÄTTLER, C., HIGGINS, J. (2017) Testing Urey's carbonate-silicate cycle using the calcium isotopic composition of sedimentary carbonates. *Earth and Planetary Science Letters* 479, 241–251. <https://doi.org/10.1016/j.epsl.2017.09.033>
- DAI, W., MOYNIER, F., PAQUET, M., MOUREAU, J., DEBRET, B., SIEBERT, J., GERARD, Y., ZHAO, Y. (2021) Calcium isotope measurements using a collision cell (CC)-MC-ICP-MS. *Chemical Geology* 590, 120688. <https://doi.org/10.1016/j.chemgeo.2021.120688>
- DAUPHAS, N., POURMAND, A. (2015) Thulium anomalies and rare earth element patterns in meteorites and Earth: Nebular fractionation and the nugget effect. *Geochimica et Cosmochimica Acta* 163, 234–261. <https://doi.org/10.1016/j.gca.2015.03.037>
- ENDRESS, M., BISHOFF, A. (1996) Carbonates in CI chondrites: Clues to parent body evolution. *Geochimica et Cosmochimica Acta* 60, 489–507. [https://doi.org/10.1016/0016-7037\(95\)00399-1](https://doi.org/10.1016/0016-7037(95)00399-1)
- FANTLE, M.S., TIPPER, E.T. (2014) Calcium isotopes in the global biogeochemical Ca cycle: Implications for development of a Ca isotope proxy. *Earth-Science Reviews* 129, 148–177. <https://doi.org/10.1016/j.earscirev.2013.10.004>



- GOUNELLE, M., ZOLENSKY, M. (2001) A terrestrial origin for sulfate veins in CI1 chondrites. *Meteoritics & Planetary Science* 36, 1321–1329. <https://doi.org/10.1111/j.1945-5100.2001.tb01827.x>
- HEUSER, A., EISENHAEUER, A. (2008) The calcium isotope composition ( $\delta\text{Ca-44/40}$ ) of NIST SRM 915b and NIST SRM 1486. *Geostandards and Geoanalytical Research* 32, 311–315. <https://doi.org/10.1111/j.1751-908X.2008.00877.x>
- HEZEL, DC., RUSSELL, S.S., ROSS, A.J., KEARSLEY, A.T. (2008) Modal Abundances of CAIs: Implications for bulk chondrite element abundances and fractionations. *Meteoritics & Planetary Science* 43, 1879–1894. <https://doi.org/10.1111/j.1945-5100.2008.tb00649.x>
- HUANG, S., JACOBSEN, S.B. (2017) Calcium isotopic compositions of chondrites. *Geochimica et Cosmochimica Acta* 201, 364–376. <https://doi.org/10.1016/j.gca.2016.09.039>
- HUANG, S., FARKAS, J., YU, G., PETAEV, M.I., JACOBSEN, S.B. (2012) Calcium isotopic ratios and rare earth elements abundances from refractory inclusions from the Allende CV3 chondrite. *Geochimica et Cosmochimica Acta* 77, 252–265. <https://doi.org/10.1016/j.gca.2011.11.002>
- KANG, J.T., IONOV, D.A., LIU, F., ZHANG, C.L., GOLOVIN, A.V., QIN, L.-P., ZHANG, Z.-F., HUANG, F. (2017) Calcium isotopic fractionation in mantle peridotites by melting and metasomatism and Ca isotope composition of the Bulk Silicate Earth. *Earth and Planetary Science Letters* 474, 128–137. <https://doi.org/10.1016/j.epsl.2017.05.035>
- KING, A., SCHOFIELD, P.F., HOWARD, K., RUSSELL, S.S. (2015) Modal mineralogy of CI and CI-like chondrites by X-ray diffraction. *Geochimica et Cosmochimica Acta* 165, 148–160. <https://doi.org/10.1016/j.gca.2015.05.038>
- MORLOK, A., BISCHOFF, A., STEPHAN, T., FLOSS, C., ZINNER, E., JESSBERGER, E.K. (2006) Brecciation and chemical heterogeneities of CI chondrites. *Geochimica et Cosmochimica Acta* 70, 5371–5394. <https://doi.org/10.1016/j.gca.2006.08.007>
- MOYNIER, F., HU, Y., WANG, K., ZHAO, Y., GÉRARD, Y., DENG, Z., MOUREAU, J., LI, W., SIMON, J.I., TENG, F.-Z. (2021) Potassium isotopic composition of various samples using a dual-path collision-cell-capable multiple-collector inductively coupled plasma mass spectrometer. *Chemical Geology* 571, 120144. <https://doi.org/10.1016/j.chemgeo.2021.120144>
- NAKAMURA, E., KOBAYASHI, K., TANAKA, R., KUNIHIRO, T., KITAGAWA, H. *et al.* (2022) On the origin and evolution of the asteroid Ryugu: A comprehensive geochemical perspective. *Proceedings of the Japan Academy, Series B*, 6, 227–282. <https://doi.org/10.2183/pjab.98.015>
- NAKAMURA, T., MATSUMOTO, M., AMANO, K., ENOKIDO, Y., ZOLENSKY, M. *et al.* (2022) Formation and evolution of carbonaceous asteroid Ryugu: Direct evidence from returned samples. *Science*. <https://doi.org/10.1126/science.abn8671>
- NIEDERER, F.R., PAPANASTASSIOU, D.A. (1984) Ca isotopes in refractory inclusions. *Geochimica et Cosmochimica Acta* 48, 1279–1293. [https://doi.org/10.1016/0016-7037\(84\)90062-0](https://doi.org/10.1016/0016-7037(84)90062-0)
- PALME, H., LODDERS, K., JONES, A. (2014) 2.2 - Solar System abundances of the elements. In: HOLLAND, H.D., TUREKIAN, K.K. (Exec. Eds.) *Treatise on Geochemistry (Second Edition)*. Elsevier, Oxford, 15–36. <https://doi.org/10.1016/B978-0-08-095975-7.00118-2>
- RUSSELL, W.A., PAPANASTASSIOU, D.A., TOMBRELLO, T.A. (1978) Ca isotope fractionation on the Earth and other solar system materials. *Geochimica et Cosmochimica Acta* 42, 1075–1090. [https://doi.org/10.1016/0016-7037\(78\)90105-9](https://doi.org/10.1016/0016-7037(78)90105-9)
- SAIKI, S., IMAMURA, H., ARAKAWA, M., WADA, K., TAKAGI, Y., HAYAKAWA, M., SHIRAI, K., YANO, H., OKAMOTO, C. (2017) The Small Carry-on Impactor (SCI) and the Hayabusa2 Impact Experiment. *Space Science Reviews* 208, 165–186. <https://doi.org/10.1007/s11214-016-0297-5>
- SIMON, J.I., DePAOLO, D.J. (2010) Stable calcium isotopic composition of meteorites and rocky planets. *Earth and Planetary Science Letters* 289, 457–466. <https://doi.org/10.1016/j.epsl.2009.11.035>
- SIMON, J.I., JORDAN, M.J., TAPPA, A., SCHAUBLE, E., KOHLL, E., YOUNG, E.D. (2017) Calcium and titanium isotope fractionation in refractory inclusions: Tracers of condensation and inheritance in the early solar system. *Earth and Planetary Science Letters* 472, 277–288. <https://doi.org/10.1016/j.epsl.2017.05.002>
- TACHIBANA, S., SAWADA, H., OKAZAKI, R., TAKANO, Y., SAKAMOTO, K. *et al.* (2022) Pebbles and sand on asteroid (162173) Ryugu: In situ observation and particles returned to Earth. *Science* 375, 1011–1016. <https://doi.org/10.1126/science.abj8624>
- VALDES, M., MOREIRA, M., FORIEL, J., MOYNIER, F. (2014) The nature of Earth's building blocks as revealed by calcium isotopes. *Earth and Planetary Science Letters* 394, 135–145. <https://doi.org/10.1016/j.epsl.2014.02.052>
- YADA, T., ABE, M., OKADA, T., NAKATO, A., YOGATA, K. *et al.* (2022) Preliminary analysis of the Hayabusa2 samples returned from C-type asteroid Ryugu. *Nature Astronomy* 6, 214–220. <https://doi.org/10.1038/s41550-021-01550-6>
- YOKOYAMA, T., NAGASHIMA, K., NAKAI, I., YOUNG, E.D., ABE, Y. *et al.* (2022) Samples returned from the asteroid Ryugu are similar to Ivuna-type carbonaceous meteorites. *Science* <https://doi.org/10.1126/science.abn7850>

