

Assembly and nuclear export of pre-ribosomal particles in budding yeast

Journal Article**Author(s):**

Gerhardy, Stefan; Menet, Anna M.; Peña, Cohue; Petkowski, Janusz J.; Panse, Vikram G.

Publication date:

2014-08

Permanent link:

<https://doi.org/10.3929/ethz-b-000088112>

Rights / license:

[In Copyright - Non-Commercial Use Permitted](#)

Originally published in:

Chromosoma 123(4), <https://doi.org/10.1007/s00412-014-0463-z>

Funding acknowledgement:

260676 - Dissecting the biogenesis of eukaryotic ribosomal subunits (EC)

Assembly and nuclear export of pre-ribosomal particles in budding yeast

Stefan Gerhardy · Anna Maria Menet · Cohue Peña ·
Janusz Jurand Petkowski · Vikram Govind Panse

Received: 18 November 2013 / Revised: 18 March 2014 / Accepted: 7 April 2014 / Published online: 11 May 2014
© Springer-Verlag Berlin Heidelberg 2014

Abstract The ribosome is responsible for the final step of decoding genetic information into proteins. Therefore, correct assembly of ribosomes is a fundamental task for all living cells. In eukaryotes, the construction of the ribosome which begins in the nucleolus requires coordinated efforts of >350 specialized factors that associate with pre-ribosomal particles at distinct stages to perform specific assembly steps. On their way through the nucleus, diverse energy-consuming enzymes are thought to release assembly factors from maturing pre-ribosomal particles after accomplishing their task(s). Subsequently, recruitment of export factors prepares pre-ribosomal particles for transport through nuclear pore complexes. Pre-ribosomes are exported into the cytoplasm in a functionally inactive state, where they undergo final maturation before initiating translation. Accumulating evidence indicates a tight coupling between nuclear export, cytoplasmic maturation, and final proofreading of the ribosome. In this review, we summarize our current understanding of nuclear export of pre-ribosomal subunits and cytoplasmic maturation steps that render pre-ribosomal subunits translation-competent.

Introduction

Error-free protein synthesis is vital for optimal cellular growth and proliferation, a fundamental task carried out by the

ribosome. This universal molecular machine consists of a large and small subunit comprising 60 % ribosomal RNA (rRNA) and 40 % ribosomal proteins (r-proteins) (Melnikov et al. 2012). The small subunit decodes the genetic information by bringing together the messenger RNA (mRNA) template and cognate transfer RNAs (tRNAs). The large subunit catalyzes the peptidyl transfer reaction to synthesize the nascent polypeptide chain. Despite a similar core, eukaryotic ribosomes are significantly larger than their prokaryotic counterparts. The large ribosomal subunit (60S) in yeast contains three rRNAs (25S, 5.8S, 5S) and 46 r-proteins, whereas the small subunit (40S) contains one single rRNA (18S) and 33 r-proteins (Ben-Shem et al. 2011; Klinge et al. 2011; Rabl et al. 2011). While the structure of the mature yeast ribosome is described at the molecular level, our knowledge regarding its assembly is emerging.

The assembly of the eukaryotic ribosome is a highly dynamic process that occurs in different cellular compartments: the nucleolus, the nucleoplasm, and the cytoplasm. In contrast to prokaryotes, eukaryotic ribosome assembly requires coordinated efforts of the intracellular transport machinery as well as numerous transiently interacting nonribosomal assembly factors. Pre-ribosomal particles released from the nucleolus undergo sequential maturation in the nucleoplasm and cytoplasm before they acquire translation competence.

Pioneering work from the Planta and Warner laboratories, in the early 1970s, led to the identification of the earliest pre-ribosome: the 90S that is the common precursor of the mature 60S and 40S subunits (Trapman et al. 1975; Udem and Warner 1972). The 90S was thought to contain pre-rRNAs, r-proteins, and numerous assembly factors. In the early 1990s, genetic screens in budding yeast permitted the identification of ~30 assembly factors. Analyses of these mutant strains primarily performed at the rRNA level led to the ordering and positioning of the cleavage sites within maturing pre-rRNA (Kressler et al. 1999; Venema and Tollervy 1995). In the late 1990s, visual approaches were developed to identify factors involved

Stefan Gerhardy, Anna Maria Menet, Cohue Peña, and Janusz Jurand Petkowski contributed equally to this work.

S. Gerhardy · A. M. Menet · C. Peña · J. J. Petkowski ·
V. G. Panse (✉)

Institute of Biochemistry (IBC), ETH Zürich, Otto-Stern-Weg 3,
8093 Zürich, Switzerland
e-mail: vikram.panse@bc.biol.ethz.ch

S. Gerhardy
BSM Program, Life Science Zürich Graduate School,
Winterthurerstrasse 190, 8057 Zürich, Switzerland

in the transport of pre-ribosomal subunits (Hurt et al. 1999; Milkereit et al. 2003; Moy and Silver 2002; Stage-Zimmermann et al. 2000). Screening of temperature-sensitive mutant libraries by visual approaches revealed factors involved in early ribosome maturation indicating a tight coupling between assembly and transport of pre-ribosomes (Gadal et al. 2001a, b; Milkereit et al. 2001). Despite these advances, the ribosome assembly pathway remained refractory to biochemical analyses. Only during the last decade, tandem affinity purification (TAP) protocols and sensitive mass spectrometry (MS) permitted the isolation and compositional analyses of pre-ribosomal particles (Bassler et al. 2001; Harnpicharnchai et al. 2001). These approaches have facilitated the general ordering of assembly events on the 60S and 40S maturation pathways and provided a framework to dissect this highly dynamic process (Grandi et al. 2002; Nissan et al. 2002; Schafer et al. 2003). Structural approaches are now beginning to facilitate high-resolution analyses of this dynamic pathway (Armache et al. 2010; Ben-Shem et al. 2011; Bradatsch et al. 2012; Greber et al. 2012; Klinge et al. 2011; Rabl et al. 2011). These studies are also guiding mechanistic analyses in higher eukaryotes (Tafforeau et al. 2013; Wild et al. 2010) and uncovering the molecular basis of ribosomopathies: diseases that are associated with impaired ribosome assembly and function.

Here, we summarize our current understanding of the eukaryotic ribosome assembly pathway in budding yeast with a focus on the nuclear export and cytoplasmic maturation of pre-ribosomal particles. For excellent reviews on bacterial ribosome assembly, the readers are referred to Shajani et al. (2011) and Britton (2009).

Assembly of the earliest ribosomal precursor, the 90S pre-ribosome

The process of ribosome assembly in budding yeast begins with RNA polymerase I-driven transcription of ribosomal DNA (rDNA) repeats on chromosome XII in the nucleolus to produce 35S pre-rRNAs (Fig. 1). Co-transcriptional association with small nucleolar RNAs (snoRNAs), assembly factors, and r-proteins mainly of the 40S drives the formation of the earliest ribosomal precursor, the 90S pre-ribosome (Grandi et al. 2002). The emerging 35S pre-rRNA can be cleaved co-transcriptionally in the internal transcribed spacer 1 (ITS1), thereby releasing the pre-40S subunit (Kos and Tollervey 2010; Osheim et al. 2004).

Two independent studies have revealed the composition of the 90S that remained refractory to biochemical analyses for nearly 30 years. By isolating Mpp10-TAP, a factor associated with the box C/D U3 snoRNA and a crucial component of the 90S, the Baserga group discovered 17 novel assembly factors, called UTPs (1–17) (Dragon et al. 2002) (Fig. 1). They

designated this large >2.2 MDa U3 snoRNA-containing particle responsible for processing of the small subunit (SSU) processome. The composition of the Mpp10-TAP particle significantly overlaps with several nucleolar pre-ribosomal particles isolated and characterized by the Hurt laboratory (Grandi et al. 2002).

Assembly of the 90S appears to be a hierarchical addition of pre-formed protein sub-complexes (Fig. 1a). The stepwise assembly of UTP complexes and the U3 snoRNP is closely coupled to the pre-rRNA modification/folding and possibly drives compaction of the 90S (Perez-Fernandez et al. 2011). Several components of the 90S contain motifs involved in RNA-binding and/or protein-protein interaction (Table 1 and Fig. 1b). A protein-protein interaction map of the SSU processome, compiled by the Baserga group, provides an important framework for the elucidation of the architecture of the 90S (Lim et al. 2011). A detailed understanding of protein-protein and RNA-protein interactions will also be crucial to uncover the spatial-temporal assembly of the 90S and its subsequent disassembly.

More than 60 different snoRNPs mediate >100 covalent modifications of the 35S pre-rRNA during the assembly of the 90S (Decatur and Fournier 2002, 2003; Decatur et al. 2007; Hughes and Ares 1991; Hughes 1996). There are two types of modifications: methylation of the 2'-hydroxyl group of the ribose sugars (2'-O-methylation) carried out by C/D box containing snoRNAs and conversion of uridine to pseudouridine carried out by H/ACA snoRNAs (Kiss 2001). These pre-rRNA modifications by snoRNAs were suggested to be important for correct folding and possibly compaction of the 90S structure (Watkins and Bohnsack 2012). Modifications of the 35S pre-rRNA are accompanied by early pre-rRNA cleavages within the developing 90S pre-ribosome (Wehner et al. 2002). Binding of U3 snoRNP to pre-rRNA is necessary for the earliest pre-rRNA cleavages at the A₀ and A₁ sites and crucial for the assembly of the early pre-40S subunit (Borovjagin and Gerbi 1999, 2001; Marmier-Gourrier et al. 2011) (reviewed in Phipps et al. (2011), Watkins and Bohnsack (2012), and Yip et al. (2013)).

Cleavage at the A₂ site, by a yet unidentified endonuclease, releases the earliest pre-40S and pre-60S particles (Fig. 2). Pre-40S subunits containing immature 20S pre-rRNAs are rapidly exported into the cytoplasm where they undergo final pre-rRNA processing to 18S rRNA. In contrast, the 27SA₂ pre-rRNA precursor within a pre-60S subunit undergoes a series of sequential processing steps to yield the mature 25S rRNA and 5.8S rRNA (Fig. 2b). A third rRNA species of the 60S subunit, the 5S rRNA, is generated by RNA polymerase III (Fig. 2). Processing of the 5S rRNA 3' end can be carried out by multiple redundant exonucleases Rex1, Rex2, and Rex3 (van Hoof et al. 2000). A rationally guided screen for ribosome biogenesis factors uncovered a requirement of Snu66, a component of the splicing machinery, for proper




Fig. 1 Model for the hierarchical assembly of the 90S pre-ribosome. **a** The assembly of the tUTP sub-complex is responsible for the initial formation of the 90S. This step allows subsequent incorporation of the indicated sub-complexes. Two independent assembly steps guide 90S formation: recruitment of the U3 snoRNP and UTP-B sub-complexes (*top*). These primary steps are necessary for the assembly of at least 20 components of the particle (Dunbar et al. 1997; Granneman et al. 2003; Lee and Baserga 1999; Wehner et al. 2002). GTPase Bms1 is necessary

for a secondary assembly step that promotes the subsequent incorporation of numerous proteins and the Mpp10 sub-complex. A second assembly step involves the incorporation of Rrp5 (*bottom*), which is crucial for the recruitment of the UTP-C sub-complex but not the U3 snoRNP, the tUTP, or the UTP-B complexes (Perez-Fernandez et al. 2007; Vos et al. 2004). **b** Schematic representation of the protein constituents of the 90S sub-complexes. The domain identification was performed using the online tool Pfam, as described by Punta et al. (2012)

processing of the 5S rRNA precursor (Li et al. 2009). However, its precise contribution to this processing step remains unclear.

Compositional dynamics of pre-ribosomal particles before nuclear export

After separation of the 90S into pre-40S and pre-60S particles, the two precursors follow independent maturation pathways. Pre-40S subunits undergo few compositional changes as they travel through the nucleoplasm and are rapidly exported into the cytoplasm. During their transit through the nucleoplasm, pre-40S subunits associate with protein kinases (Rio1 and Hrr25) and the ATPase Rio2. These energy-consuming steps possibly prepare the pre-40S subunit for nuclear export and/or

final cytoplasmic maturation (Geerlings et al. 2003; Vanrobays et al. 2003). However, the relevant substrates for Rio1, Rio2, and Hrr25 remain to be uncovered. The assembly factors Enp1, Ltv1, and the r-protein uS3 (Rps3, new nomenclature as proposed by Ban et al. (2014)) form a sub-complex at the landmark beak structure of the 40S subunit (Schafer et al. 2006). This complex can be dissociated in vitro from pre-40S subunits by the activity of the conserved kinase Hrr25. It was proposed that phosphorylation of the Enp1-Ltv1-uS3 (Enp1-Ltv1-Rps3) sub-complex increases the conformational flexibility in the head region of the pre-40S subunit in vivo. Hrr25 depletion impairs nuclear export of pre-40S subunits supporting the notion that its kinase activity makes the rigid head region more flexible during transport through the nuclear pore complex (NPC). Dephosphorylation by an unknown phosphatase in the cytoplasm could permit the stable

Table 1 Composition of sub-complexes within the earliest ribosome precursor, the 90S. The domain identification was performed using the online tool Pfam, as described by Punta et al. (2012)

Sub-complex	Component	Activity/function/[predicted domain composition]	References
tUTP/UTP-A	tUTP4	[WD40 repeats]	Dragon et al. (2002), Gallagher et al. (2004), Krogan et al. (2002)
	tUTP5	[UTP12 domain]	Dragon et al. (2002), Gallagher et al. (2004)
	tUTP8	Participates also in nuclear export of tRNA; [UTP8 family domain]	Dragon et al. (2002), Gallagher et al. (2004), Strub et al. (2007)
	tUTP9	[Dip2/UTP12 domain]	Dragon et al. (2002), Gallagher et al. (2004), Krogan et al. (2002)
	tUTP10	HEAT repeats; [snoRNA BD]; [BP28CT domain]	Dragon et al. (2002), Gallagher et al. (2004), Krogan et al. (2002)
	tUTP15	[WD40 repeats]; [UTP15C domain]	Dragon et al. (2002), Gallagher et al. (2004), Krogan et al. (2002)
	tUTP17 (Nan1)	Member of the RENT complex; [WD40 repeats]	Dragon et al. (2002), Gallagher et al. (2004), Krogan et al. (2002)
	Pol5	Required for rRNA synthesis; [DNA pol phi domain]	Gallagher et al. (2004), Krogan et al. (2002), Shimizu et al. (2002)
U3 snoRNP associated	Nop1	2'-O-methyltransferase; [Fibrilrin domain]	Lischwe et al. (1985), Tollervey et al. (1991)
	Nop56	KKE/D repeats; coiled coils; structural scaffold for the snoRNP complex; [Nop5 RBD]; [Nopic domain]; [snoRNA BD]	Lafontaine and Tollervey (2000), Watkins and Bohnsack (2012)
	Nop58	KKE/D repeats; [Nop5 RBD]; [Nopic domain]; [snoRNA BD]	Wu et al. (1998)
	Snu13	Binds to K-turn motifs of U3 snoRNA; member of the U4/U6-U5 tri-snoRNP; [r-protein L7-like domain]	Watkins et al. (2000)
	Rrp9	Binds to the B/C motif of U3 snoRNA; essential for recruitment of U3 snoRNP to SSU processome; [WD40 repeats]	Lukowiak et al. (2000), Venema et al. (2000)
UTP-B	UTP1	β Transducin family; [WD40 repeats]; [Dip2/UTP12 domain]	Champion et al. (2008), Dosil and Bustelo (2004), Dragon et al. (2002), Krogan et al. (2002)
	UTP6	HAT motif; [U3 assoc. domain]	Champion et al. (2008), Dosil and Bustelo (2004), Dragon et al. (2002), Krogan et al. (2002)
	UTP12	[WD40 repeats]; [Dip2/UTP12 domain]	Champion et al. (2008), Dosil and Bustelo (2004), Dragon et al. (2002), Krogan et al. (2002)
	UTP13	[WD40 repeats]; [UTP-like3/WD40 assoc. domain]	Champion et al. (2008), Dosil and Bustelo (2004), Dragon et al. (2002), Krogan et al. (2002)
	UTP18	[WD repeats]	Bernstein et al. (2004), Champion et al. (2008), Dosil and Bustelo (2004), Krogan et al. (2002)
	UTP21	Coiled coils; [WD40 repeats]; [UTP21 domain]	Bernstein et al. (2004), Champion et al. (2008), Dosil and Bustelo (2004), Krogan et al. (2002), Samanta and Liang (2003)
Bms1/Rcl1	Bms1	GTPase stimulated by Rcl1; [Nuc121 domain]; [DUF633 domain]	Karbstein et al. (2005), Wegierski et al. (2001)
	Rcl1	Stimulates Bms1; [RNA 3'-terminal phosphate cyclase-like domain]	Billy et al. (2000), Tanaka et al. (2011), Wegierski et al. (2001)
UTP-C	Rrp7	Also part of the CURI complex; [Rrp7 domain]	Baudin-Baillieu et al. (1997), Dalley et al. (2008), Krogan et al. (2002), Rudra et al. (2007)
	UTP22	Also part of the CURI complex; [nucleolar RNA assoc. protein domain]	Bernstein et al. (2004), Krogan et al. (2002), Peng et al. (2003), Rudra et al. (2007), Samanta and Liang (2003)
	Cka1	α Catalytic subunit casein kinase 2; nonessential; also part of the CURI complex; [protein kinase domain]	Krogan et al. (2002), Rudra et al. (2007)
	Cka2	α' Catalytic subunit casein kinase 2; nonessential; also part of the CURI complex; [protein kinase domain]	Krogan et al. (2002), Rudra et al. (2007)
	Ckb1	β Regulatory subunit casein kinase 2; nonessential; also part of the CURI complex; [casein kinase II reg. subunit domain]	Krogan et al. (2002), Rudra et al. (2007)
	Ckb2		Krogan et al. (2002), Rudra et al. (2007)

(continued)

Sub-complex	Component	Activity/function/[predicted domain composition]	References
Mpp10	Imp3	β' Regulatory subunit casein kinase 2; nonessential; also part of the CUR1 complex; [casein kinase II reg. subunit domain]	Lee and Baserga (1999)
	Imp4	[S4 RNA-binding domain]	Ng et al. (2005)
	Mpp10	Coiled coils; interacts with the hinge region of U3 snoRNA; [Mpp10 domain]	Dunbar et al. (1997)

incorporation of uS3 (Rps3) into the mature 40S subunit and formation of the final beak structure (Schafer et al. 2006).

In the 60S assembly pathway, incorporation of 5S rRNA in complex with r-proteins uL18 (Rpl5) and uL5 (Rpl11) is an important early nucleolar/nuclear event. Recent work from the Hurt laboratory suggested that the r-proteins uL18 (Rpl5) and uL5 (Rpl11) are co-imported by Syo1, an adaptor for the importin Kap104 (Kressler et al. 2012a). In the nucleus, the uL18-uL5-Syo1 (Rpl5-Rpl11-Syo1) complex is released from Kap104 in a RanGTP-dependent manner and then loaded on the 5S rRNA (Kressler et al. 2012a). Incorporation of uL18-uL5-5S rRNA complex into the pre-60S particle is facilitated by the assembly factors Rpf2 and Rpf1 (Ciganda and Williams 2011; Zhang et al. 2007). However, the exact timing of incorporation into pre-60S subunits remains to be investigated.

During the journey through the nucleoplasm, pre-60S particles associate with >150 assembly factors as they travel through the nucleoplasm toward the nuclear periphery. At distinct maturation stages, assembly factors are released from pre-ribosomal particles and recycled back to participate in new rounds of biogenesis steps. This sequential reduction in complexity of the pre-60S subunits is very likely triggered by a multitude of energy-consuming enzymes. ATP-dependent RNA helicases, AAA-ATPases, ABC-ATPases, and GTPases associate with maturing pre-ribosomal particles and confer directionality to the assembly and maturation process. The binding site(s) of these energy-consuming enzymes on maturing pre-60S particles are beginning to be uncovered (Kressler et al. 2012b; Strunk et al. 2012).

Among the diverse energy-consuming enzymes that drive ribosome assembly, the molecular roles of AAA-ATPases are better understood. Three essential AAA-ATPases contribute to pre-60S subunit maturation. The AAA-ATPases Rix7 and Drg1 are closely related to the well-characterized Cdc48 (p97 in mammals) and contribute to early nucleolar and cytoplasmic pre-60S subunit maturation, respectively (Kappel et al. 2012; Pertschy et al. 2007). Rea1, which is the largest protein in yeast, shares similarity to the microtubule motor protein dynein heavy chain and functions at different nuclear steps during 60S maturation (Bassler et al. 2010; Kressler et al. 2012b; Ulbrich et al. 2009).

Rix7 is the first AAA-ATPase that is involved in the maturation of the pre-60S subunit. Like the archetypical AAA-ATPase Cdc48, which uses cycles of ATP hydrolysis to remodel protein complexes (Meyer et al. 2012), Rix7 was suggested to strip off the assembly factor Nsa1 and facilitate the nucleolar to nucleoplasmic transition of pre-60S subunits along the assembly pathway (Kressler et al. 2008). How Rix7 recognizes Nsa1 and extracts it from maturing pre-60S subunits remains to be investigated. It was suggested that Rix7 might recognize posttranslational modifications like ubiquitin and SUMO, either directly or via adaptor proteins, to release Nsa1 and further assembly factors (Kressler et al. 2008; Panse et al. 2006).

Rea1 consists of six ATPase modules, forming a binding platform on the ribosome and binds in close proximity to the Rix1-Ipi3-Ipi1 sub-complex. A long α -helical linker domain, a D/E-rich region, and a functionally important metal-ion-dependent adhesion site (MIDAS) domain follow the ATPase modules. By employing the MIDAS domain, Rea1 can directly contact the MIDAS interacting domains (MIDO) of Ytm1 and Rsa4 and triggers the release of both assembly factors via ATP hydrolysis cycles (Bassler et al. 2010; Ulbrich et al. 2009). Removal of the Ytm1-Erb1-Nop7 sub-complex may trigger the release of neighboring biogenesis factors on the pre-ribosomal particles, as well as recruitment of further assembly factors involved in later biogenesis steps. The release of Rsa4 occurs at a later stage of the assembly pathway and signals the progression toward export competence (Matsuo et al. 2014; Nissan et al. 2002; Ulbrich et al. 2009).

Recently, Rea1 and the GTPase Nug2 were implicated in a nuclear checkpoint step that prevents premature formation of an export competent pre-60S subunit (Matsuo et al. 2014). The GTPase Nug2 (also known as Nog2) binds the pre-60S subunits in the nucleus at a site, which clashes with the binding site for the export adaptor Nmd3 (Sengupta et al. 2010). Nug2 bound to the pre-60S subunit allows nucleoplasmic maturation to occur. Release of Nug2 depends on its GTPase activity as well as the ATPase activity of Rea1. Only after the release of Nug2, Nmd3 can bind pre-60S subunits and they can be exported. This mechanism provides a timed acquisition of export competence for the large ribosomal subunit (Matsuo et al. 2014).

Nuclear export of pre-ribosomal particles



Assembly of eukaryotic ribosomes begins in the nucleolus, but translation of mRNA into proteins by the mature ribosome occurs in the cytoplasm. Inevitably, pre-ribosomal particles need to be transported through the NPCs into the cytoplasm. NPCs are huge protein assemblies embedded within the double lipid bilayer of the nuclear envelope and serve as ports to exchange macromolecules between the nucleus and cytoplasm. For excellent reviews on the NPC structure and assembly, readers are referred to Hoelz et al. (2011), Wentz and Rout (2010), and Zwerger and Medalia (2012). The NPC transport channel permits free diffusion of molecules <40 kDa (Raices and D'Angelo 2012). Translocation of complex cargos such as the charged >2 MDa pre-ribosome through the hydrophobic phenylalanine-glycine (FG)-repeat meshwork of the NPC channel poses a major challenge. Transport of pre-ribosomal particles is facilitated by multiple factors that interact with the FG-meshwork of the NPC channel (Fig. 3). In actively growing budding yeast cells, it is estimated that each NPC contributes to the export of ~25 pre-ribosomal particles per minute (Warner 1999). Such a rapid process requires an efficient transport machinery that ensures rapid translocation of pre-ribosomal cargos through the NPC channel. Cell biological tools that monitored the intracellular localization of ribosomal subunits revealed the requirement of several components of the NPC, the Ran GTPase cycle, and the export receptor Xpo1, in the nuclear export of pre-ribosomes (Hurt et al. 1999; Stage-Zimmermann et al. 2000). Subsequently, a visual screen and an independent genetic approach uncovered an essential nuclear export signal (NES) containing adaptor Nmd3. Nmd3 forms a complex with Xpo1 in the presence of the GTPase Ran and facilitates export of the bound pre-60S subunit (Gadal et al. 2001a, b; Ho et al. 2000).

Efficient translocation of pre-ribosomes through the NPC requires multiple export factors (Oeffinger et al. 2004). Notably, the essential general mRNA transport receptor Mex67-Mtr2 (Santos-Rosa et al. 1998; Segref et al. 1997) and the HEAT-repeat containing protein Rrp12 contribute to export of both pre-ribosomal subunits (Faza et al. 2012; Oeffinger et al. 2004; Yao et al. 2007) (Fig. 3). Mex67, the large subunit of the heterodimer, is composed of an amino-terminal (N) domain, a leucine-rich repeat (LRR) domain, a nuclear transport factor 2 (NTF2)-like middle domain, and a C-terminal ubiquitin associated (UBA)-like domain (Strasser et al. 2000). Mtr2 shares structural features with NTF2 (Bayliss et al. 2002) and heterodimerizes with the NTF2-like middle domain of Mex67 (Santos-Rosa et al. 1998; Strasser et al. 2000). Loops emanating from the NTF2-like domains contribute to pre-60S and pre-40S subunit binding (Fribourg and Conti 2003; Senay et al. 2003; Yao et al. 2007) suggesting a versatile common interaction platform on Mex67-Mtr2. A recent study from the Tollervey laboratory revealed crosslinks

Fig. 2 The assembly pathway of the eukaryotic ribosome. **a** Co-transcriptional recruitment of small subunit r-proteins and assembly factors to the 35S pre-rRNA yields the earliest ribosomal precursor, the 90S (orange). Cleavage at the A₂ site separates the 90S into a pre-40S subunit (green) and a pre-60S subunit (blue), which undergo independent maturation. Transiently associating assembly factors drive maturation of the pre-ribosomal subunits as they travel through the nucleoplasm toward the NPCs. Final maturation in the cytoplasm yields translation competent ribosomal subunits. **b** The pre-rRNA processing pathway of pre-ribosomal particles. The common 35S rRNA precursor is trimmed at both ends (Kufel et al. 1999) and cleaved at the A₂ site. The cleavage yields 20S and 27SA₂ rRNAs which mature independently: The 20S rRNA is processed to 18S rRNA in the cytoplasm (Lamanna and Karbstein 2009; Pertschy et al. 2009). The 27SA₂ pre-rRNA is processed in two ways generating two different 5.8S rRNAs (Lygerou et al. 1996). Following the cleavage at the A₃ site, the 5' end of 5.8S is rapidly processed to site B_{1S}, by exonucleases Rrp17 (Oeffinger et al. 2009) and the Rat1-Rai1 heterodimer (Henry et al. 1994). 27SB_{1L} and the 27SB_{1S} pre-rRNAs are cleaved at the C₂ site, to produce the 7S_{L/S} and the 25.5S pre-rRNA. The latter is converted into the 25S rRNA by Rat1-Rai1 in the nucleus (Geerlings et al. 2000). The 7S pre-rRNAs are processed at the 3' ends to shorter intermediates (Allmang et al. 1999; Mitchell et al. 1997) and then to 6S_{1S/1L} by the nuclear exosome (Briggs et al. 1998). Processing of 6S pre-rRNA to mature 5.8S rRNA takes place in the cytoplasm and requires the nucleases Rex1, Rex2, and Ngl2 (Faber et al. 2002; Thomson and Tollervey 2010; van Hoof et al. 2000). The 5S rRNA is processed in by the nucleases Rex1, Rex2, and Rex3 (van Hoof et al. 2000). For detailed information on pre-rRNA processing, readers are referred to Woolford and Baserga (2013) and Fromont-Racine et al. (2003)

between Mex67 and the 3' end of 20S pre-rRNA transcript as well as 5.8S rRNA and multiple regions along the 25S rRNA in the vicinity of 5.8S rRNA supporting the idea that Mex67 interacts with both pre-40S and pre-60S subunits (Tuck and Tollervey 2013). The NTF2-like domains of Mex67-Mtr2 and the UBA-like domain of Mex67 interact directly with the FG-meshwork and therefore facilitate nuclear export of the bound cargo (Strasser et al. 2000; Strawn et al. 2001). Mex67-Mtr2 does not directly rely on the Ran cycle for export (Yao et al. 2007) and is the only known transport factor that contributes to the export of three major cargos: mRNA, pre-60S, and pre-40S subunits. It is not understood how the Mex67-Mtr2-pools are split to export all three substrates. Understanding the molecular basis of this allocation will reveal how the three export pathways cross talk to deliver required levels of mRNA and ribosomal subunits in the cytoplasm.

Rrp12 is the second factor that binds both pre-60S and pre-40S subunits and is required for their nuclear export (Oeffinger et al. 2004). Rrp12 contains secondary structural elements called HEAT (Huntingtin, elongation factor 3, protein phosphatase 2A, and TOR1) repeats, which are found in other RanGTP-dependent export factors, and shown to interact with the FG-meshwork. How Rrp12 binds FXFG-repeat nucleoporins remains to be investigated. Whether RanGTP regulates the interaction of Rrp12 with pre-ribosomal subunits and/or FG-nucleoporins remains to be determined. Identifying mutants of Rrp12 that are impaired in binding to pre-






Fig. 3 Transport of pre-ribosomal particles through the NPC channel. Export of the pre-ribosomal particles is facilitated by indicated export factors (yellow) that interact with the FG meshwork of the NPC channel. Cryo-EM maps of late pre-ribosomal particles isolated by Rio2-TAP and

Alb1-TAP. The pre-40S model is adapted from Figure 2 from Strunk et al. (2011) and the pre-60S model is adapted from Figure 2 from Bradatsch et al. (2012)

ribosomal particles, FXFG-repeat nucleoporins and RanGTP will clarify the contribution of Rrp12 to pre-ribosome export.

Additionally, pre-ribosomal particles employ multiple non-essential, auxiliary factors that can directly bind FG-rich nucleoporins and directly facilitate translocation of pre-ribosomal particles through the NPC channel (Fig. 3). The *trans*-acting factor Arx1 that localizes to the exit tunnel plays an auxiliary role in pre-60S subunit nuclear export (Bradatsch et al. 2007; Hung et al. 2008). Arx1 contains a methionine aminopeptidase (MetAP)-like fold, which is present in a family of proteins that remove the N-terminal methionine from nascent polypeptides as they emerge from the ribosome. However, Arx1 lacks a methionine aminopeptidase activity. Mutations in the methionine-binding pocket impair the function of Arx1 in pre-60S subunit export, which led to the proposal that this fold has evolved to interact with FG-nucleoporins (Bradatsch et al. 2007). Two recent studies revealed that the proposed FG-interacting pocket of Arx1 points toward the exit tunnel of the 60S subunit and hence does not appear to be exposed to the solvent. Further, Arx1 not only

covers the exit tunnel of the 60S subunit, but it also arrests a conserved rRNA expansion segment 27 (ES27) in a so-called tunnel conformation (Bradatsch et al. 2012; Greber et al. 2012). It could be that Arx1 gets detached from the main body of the pre-60S subunit, while it remains bound to the pre-60S subunit via its interaction with ES27, during translocation of pre-60S subunits through the NPC. Such a scenario may reconcile the apparent paradox, as to how Arx1 simultaneously interacts with the pre-60S subunit as well as FG-rich nucleoporins. Nevertheless, the functional significance of the interaction between ES27 and Arx1 remains to be determined.

Functional screens in yeast have identified factors that directly promote nuclear export of pre-ribosomes. These approaches uncovered the shuttling assembly factors (Ecm1 and Bud20) and mRNA export factor (Npl3) involved in pre-60S subunit nuclear export (Altwater et al. 2012; Bassler et al. 2001, 2012; Hackmann et al. 2011; Yao et al. 2010). Whether pre-ribosomal subunits employ all export factors for their translocation through the NPC is currently unclear. It could be that a minimal set might be sufficient to facilitate

rapid export. A particular challenge is to localize export factors on pre-ribosomal particles. These analyses are expected to provide insight into how pre-ribosomal particles are oriented during the translocation through the NPC channel.

All export factors described to date utilize the FG-meshwork to translocate pre-ribosomal subunits through the NPCs. Recently, we have uncovered a role for the non-FG-interacting transport factor Gle2 in pre-60S subunit export (Occhipinti et al. 2013). Gle2 interacts with pre-60S through a conserved basic patch and utilizes a second interaction surface to simultaneously bind the GLEBS (Gle2-binding sequence) motif of Nup116 (Fig. 3). These interactions together facilitate the transit of the pre-60S particle through the NPC. Mutations that impair the function of Gle2 in pre-60S nuclear export do not affect mRNA export (Occhipinti et al. 2013). Curiously, the recruitment of Gle2 to pre-60S subunits requires its prior tethering to the GLEBS motif of nucleoporin Nup116. Thus, Gle2 could utilize distinct interaction surfaces to prevent kinetic delays experienced by mRNPs and pre-60S subunits during translocation through the NPC channel, especially in the case when cargos have failed to recruit its optimal complement of export factors.

In contrast to the pre-60S, fewer export factors are described for the pre-40S subunit. Studies in mammalian cells have revealed multiple NES containing assembly factors (Ltv1, Dim2, and Rio2) that recruit the export factor Crm1 (Xpo1) (Zemp et al. 2009). An essential NES containing adaptor for pre-40S subunits that recruits Xpo1 remains elusive. One possibility could be that multiple NES containing adaptors play redundant roles to recruit the essential export receptor Xpo1 and guarantee efficient nuclear export of pre-40S subunits.

Another conserved factor that specifically functions in the nuclear export of pre-40S subunits is the conserved RanGTP-binding protein Yrb2 (Moy and Silver 2002; Taura et al. 1998). The *yrb2*Δ mutant exhibits a marked decrease in the 40S subunit levels and strong nucleoplasmic accumulation of the small subunit reporters (Faza et al. 2012; Moy and Silver 2002). Notably, in vitro studies showed that the human homolog of Yrb2 (RanBP3) triggers the loading of Xpo1 (Crm1) and RanGTP on certain cargoes that are exported into the cytoplasm (Englmeier et al. 2001). Therefore, one possibility could be that Yrb2 delivers Xpo1 and RanGTP to certain yet unknown NES containing adaptor(s) to promote nuclear export of pre-40S subunits.

Cytoplasmic maturation pathway of pre-ribosomes

Prior to nuclear export, the majority of assembly factors, which associate with pre-ribosomal particles during early biogenesis, are released after fulfilling their function. Only a handful of assembly factors travel with pre-ribosomal particles to the cytoplasm. The release of these factors, the

incorporation of the remaining r-proteins, and final pre-rRNA processing events constitute cytoplasmic maturation in the ribosome biogenesis pathway (Fig. 4). These steps are not only crucial for completing ribosome maturation, but are also crucial for new rounds of ribosome biogenesis. A failure to release and recycle assembly and export factors back to the nucleus induces pre-rRNA processing delays, assembly defects, and impaired nuclear export.

Cytoplasmic maturation of pre-40S subunits

The pre-40S subunits are accompanied to the cytoplasm by a handful of proteins (Enp1, Tsr1, Ltv1, Dim1, Dim2, Nob1, Rio2, and Hrr25) that contribute to their export as well as subsequent pre-rRNA processing (Strunk et al. 2011). How these shuttling assembly factors are released from pre-40S subunits remains unclear. Cytoplasmic processing of immature 20S pre-rRNA within pre-40S subunits involves two conserved events: dimethylation of the 20S pre-rRNA and the endonucleolytic cleavage (site D in Fig. 2b) of 20S pre-rRNA to 18S rRNA.

A late pre-rRNA modification step in 40S biogenesis is the dimethylation of two adenine bases near the 3' end of the 18S rRNA. The enzyme responsible for catalyzing this modification is the essential factor Dim1 that is loaded on the pre-40S subunits (Lafontaine et al. 1994). Dimethylation is first detected on the 20S rRNA precursor and was suggested to take place once the pre-40S particle reaches the cytoplasm. Although dimethylation occurs late during subunit maturation, Dim1 associates already with the 90S. Dim1 depletion causes an early nucleolar pre-rRNA processing defect, which can be rescued by a catalytically inactive *dim1* mutant. Intriguingly, the catalytically inactive *dim1* mutant rescues the lethality of the *dim1*Δ strain suggesting that the dimethylating activity of Dim1 is not essential and can be separated from its essential role in early pre-rRNA processing. Dimethylation was suggested to play a role in fine-tuning translation as the *dim1* mutant displays increased antibiotic sensitivity (Lafontaine et al. 1998).

An essential cytoplasmic maturation event that renders pre-40S particles translation-competent is the endonucleolytic cleavage of the immature 20S rRNA into mature 18S rRNA. Multiple energy-consuming ATPases (Prp43, Rio2, and Fap7) and the PIN-domain endonuclease Nob1 were implicated in this late maturation step (Clissold and Ponting 2000; Geerlings et al. 2003; Jakovljevic et al. 2004; Lamanna and Karbstein 2009; Pertschy et al. 2009; Vanrobays et al. 2003). Nob1 is recruited to 40S pre-ribosomes already in the nucleus, suggesting that there must be an activating mechanism for Nob1 in the cytoplasmic compartment. Studies from the Tollervey and Karbstein laboratories revealed that pre-40S subunits interact with mature 60S subunits to form an 80S-like particle in vitro




Fig. 4 Cytoplasmic maturation of a large pre-ribosomal subunit prior to initiating translation. Exported pre-60S subunits are bound by export factors (yellow) and shuttling factors (green) which are released in the cytoplasm. The ATPase Drg1 releases Rlp24 from the pre-ribosomal particles. This event triggers the subsequent maturation events. Arx1 and Alb1 require Rei1, Jjj1, and Ssa1/Ssa2 for their release, whereas the stalk assembly can only take place after the release of the shuttling ribosomal-like protein Mrt4 by the cytoplasmic release factor Yvh1.

Recruitment of uL10 (Rpp0) releases Yvh1, which allows further assembly of the P1 (Rpp1) and P2 (Rpp2) heterodimer onto the stalk. Loading of uL16 (Rpl10) triggers the final maturation steps. The GTPase Efl1 and its co-factor Sdo1 release Tif6 and another GTPase Kre35 (Lsg1) removes Nmd3. The release mechanisms/factors of shuttling assembly factors like Nog1, Nug1, and Nsa2 and the transport factors Mex67-Mtr2, Bud20, Ecm1, and Gle2 (depicted in grey) remain to be uncovered

and in vivo (Lebaron et al. 2012; Strunk et al. 2012). The interaction with the 60S subunit triggers the activity of Nob1 to cleave 20S pre-rRNA to mature 18S rRNA in vitro. Both studies implicated that the conserved cytoplasmic GTPase and translation initiation factor eIF5b/Fun12 might be important for the formation of the 80S-like particle, which triggers this final cleavage step in the cytoplasm. Subsequently, Nob1 is released and recycled due to the action of the ABC-ATPase Rli1 that dissociates the pre-40S subunit from the 60S subunit, after 20S pre-rRNA cleavage (Strunk et al. 2012). Fun12 is not an essential gene in budding yeast, whereas the endonucleolytic cleavage of 20S pre-rRNA is essential. Considering the high abundance of pre-40S subunits, it is intriguing that the *fun12*Δ mutant shows only minor accumulation of 20S pre-rRNAs in the cytoplasm (Lebaron et al. 2012). It could be that in a *fun12*Δ mutant, pre-40S subunits containing immature 20S pre-rRNA fail to undergo cytoplasmic maturation and are then rapidly degraded. In support of an 80S-like translation cycle, mutations in the 60S subunit r-protein uL3 (Rpl3), which affect the affinity to translation elongation factors, were shown to specifically impair cytoplasmic 20S pre-rRNA processing (Garcia-Gomez et al. 2014).

Cytoplasmic maturation of pre-60S subunits

In addition to shuttling export factors (Nmd3, Arx1, Ecm1, and Mex67-Mtr2), genetic approaches identified few assembly factors (Rlp24, Tif6, Nog1, and Alb1) that associate with pre-60S particles in the nucleus and are transported to the cytoplasm. Release of these factors is catalyzed by conserved

energy-consuming GTPases (Kre35, Efl1), ATPases (Drg1, Hsp70), and their cofactors (Sdo1, Rei1, Jjj1) that transiently associate with pre-60S particles exclusively in the cytoplasm (Panse and Johnson 2010).

The Johnson laboratory established the order of known release events that provided an initial framework of the cytoplasmic maturation pathway (Fig. 4) (Lo et al. 2010). A characteristic feature of the pre-60S cytoplasmic maturation is the sequential release of assembly factors and transport factors (Fig. 4 and Table 2). The AAA-ATPase Drg1 triggers the earliest cytoplasmic maturation step on pre-60S particles, which is a prerequisite for subsequent cytoplasmic maturation steps of 60S pre-ribosomes. Catalytically impaired *drg1* mutants accumulate the nuclear assembly factors Rlp24, Nog1, Arx1, and Tif6 in the cytoplasm where they remain bound to pre-60S subunits (Pertschy et al. 2007). Thus, the activity of Drg1 is required for their release from pre-60S particles, thereby also allowing their re-import into the nucleus. AAA-ATPases typically do not exhibit broad substrate specificities (Lupas and Martin 2002), and therefore, it is unlikely that Drg1 releases each of these factors from pre-60S subunits. The Bergler laboratory demonstrated that Drg1 catalyzes the release of Rlp24 that is essential for subsequent release events (Kappel et al. 2012). A C-terminal region within Rlp24 acts as a recruiting site for Drg1 to stimulate its ATPase activity. The release of Rlp24 from the pre-60S subunit appears to require the nonessential FG-nucleoporin Nup116. Thus, the mechanochemical activity of Drg1 makes the export step irreversible and simultaneously initiates the cytoplasmic maturation cascade.

Table 2 Cytoplasmic release of shuttling assembly and transport factors in the 60S biogenesis pathway

Release factor/co-factor	Activity/domain	Target	References
Drg1	AAA-ATPase	Rlp24	Kappel et al. (2012), Pertschy et al. (2007)
Rei1	Zn-finger protein	Arx1/Alb1	Hung and Johnson (2006), Lebreton et al. (2006), Meyer et al. (2007)
Jjj1-Ssa1/Ssa2	DnaJ domain stimulates the ATPase Ssa1/Ssa2	Arx1/Alb1	Demoinet et al. (2007), Meyer et al. (2007, 2010)
Yvh1	Dual specificity phosphatase	Mrt4	Kemmler et al. (2009), Lo et al. (2009)
Efl1	GTPase	Tif6	Becam et al. (2001), Senger et al. (2001)
Sdo1	–	Tif6	Menne et al. (2007)
Kre35	GTPase	Nmd3	Hedges et al. (2005)
–	–	Bud20	
–	–	Nsa2	
–	–	Nug1	
–	–	Np13	
–	–	Mex67/Mtr2	
–	–	Nog1	
–	–	Gle2	
–	–	Ecm1	

Release of the ribosomal-like protein Rlp24 is necessary to allow the r-protein eL24 (Rpl24) to assemble into the pre-60S subunit. This exchange event triggers recruitment of the zinc-finger proteins Rei1 and Yvh1 (Altwater et al. 2012; Lo et al. 2010). Rei1 is nonessential, but needed for the recycling of Arx1 and its interacting partner Alb1 (Hung and Johnson 2006; Lebreton et al. 2006). Rei1 works in conjunction with the DnaJ domain-containing Jjj1 and the ATPase Ssa1/Ssa2 (Hsp70) to release Arx1 (Demoinet et al. 2007; Lo et al. 2010; Meyer et al. 2007, 2010). This data also indicates that Arx1 appears to have an inhibitory role in driving cytoplasmic maturation pathway of pre-60S subunits. Based on similarity of Arx1 to MetAPs, a prediction is that they bind to the same site on the ribosome and that Arx1 prevents the binding of MetAP. Further, Arx1 binds in the vicinity of the r-protein uL23 (Rpl25) at the polypeptide exit tunnel, which is an important functional site on the ribosome, as uL23 (Rpl25) interacts with the signal recognition particle (SRP) as well as the translocon in the endoplasmic reticulum (ER) (Bradatsch et al. 2012; Dalley et al. 2008; Greber et al. 2012).

The Johnson and Panse laboratories have uncovered a cytoplasmic maturation event that is crucial for assembly of the ribosome stalk, a structural landmark of the 60S subunit (Kemmler et al. 2009; Lo et al. 2009). Assembly of the stalk is a major step in acquisition of functionality of the ribosome, since it is essential for recruitment and activation of translation factors, in particular the elongation factors. In yeast, the stalk is composed of uL10 (Rpp0) and two heterodimers of P1 and P2 (Rpp1/Rpp2). uL10 (Rpp0) anchors the stalk to the ribosome by binding to the rRNA of helices 43 and 44. However, pre-60S subunits are first assembled in the nucleus with the ribosomal-like protein Mrt4 in place of uL10 (Rpp0). Mrt4

lacks the domains that recruit translation factors, necessitating the exchange of Mrt4 for uL10 (Rpp0). The dual specificity phosphatase Yvh1 is required for the removal of Mrt4 and uses its zinc-binding domain but not its phosphatase activity to release of Mrt4 from pre-60S subunits. While the key players are identified, the precise mechanism of Mrt4 release and the molecular events that lead to the assembly of the stalk remain elusive.

Following the assembly of the stalk and the removal of Arx1, Tif6 is released. Tif6 is a shuttling assembly factor that prevents the joining of immature 60S to 40S subunits (Russell and Spemulli 1979; Valenzuela et al. 1982). The release of Tif6 depends on previous events as it mislocalizes in *yvh1* mutants in which stalk assembly is blocked (Kemmler et al. 2009). During translation, the stalk functions in recruitment and activation of the GTPase eEF2 (Bargis-Surgey et al. 1999). Given that Efl1 is closely related to eEF2, stalk assembly might play a similar role in biogenesis, thus recruiting Efl1 for the release of Tif6. This also suggests that the cytoplasmic maturation events in the 60S biogenesis are coupled and ordered sequentially from Drg1-dependent release of Rlp24 to Efl1-mediated release of Tif6 (Fig. 4).

The GTPase Efl1 and the Shwachman-Bodian-Diamond syndrome protein Sdo1 are required to release Tif6 (Becam et al. 2001; Senger et al. 2001). In *efl1* and *sdo1* mutants, Tif6 accumulates on late pre-60S subunits and is mislocalized to the cytoplasm (Menne et al. 2007; Senger et al. 2001). Mutations in Tif6 that weaken its affinity for the 60S subunit suppress the growth defects of *efl1* and *sdo1* mutants, providing strong genetic evidence that Tif6 is the primary substrate of Efl1 and Sdo1. Efl1 bears a significant sequence similarity to the translation elongation factor 2, a GTPase that facilitates

translocation of the ribosome following the action of the peptidyl transferase.

The essential NES containing adaptor Nmd3 must be released from pre-60S subunits and recycled back to the nucleus. The r-protein uL16 (Rpl10) and the GTPase Kre35 (Lsg1) were implicated in the release of Nmd3 (Hedges et al. 2005; West et al. 2005). Mutations in uL16 (Rpl10) prevent the release of Nmd3 from pre-60S subunits. Moreover, mutations in Kre35 that are predicted to disrupt its GTPase activity also block Nmd3 release in the cytoplasm (Hedges et al. 2005). These results suggest that Kre35 triggers the binding of uL16 (Rpl10) to the 60S, an event that is coupled to the release of Nmd3 (Hedges et al. 2005; Karl et al. 1999; West et al. 2005). Interestingly, recent work from the Johnson laboratory indicates that loading of uL16 (Rpl10) on late pre-60S subunits is a prerequisite for the release of Tif6 (Bussiere et al. 2012).

Recently, we have employed a combination of genetic trapping, affinity purification, and a targeted proteomic approach based on selected reaction monitoring mass spectrometry (SRM-MS) to interrogate the proteome of 60S pre-ribosomes after nuclear export (Altwater et al. 2012). Using a resource of SRM assays, we uncovered unanticipated assembly factors (Bud20, Nug1, Nsa2, and Rli1). They are exported to the cytoplasm and are only released after Drg1-mediated release of Rlp24. The functional significance of shuttling behavior of the identified assembly factors is unknown. It could be that they participate directly in their transport and/or final functional proofreading of pre-60S subunits. Mechanisms that trigger the release of these shuttling assembly factors from pre-60S particles in the cytoplasm remain to be discovered.

Cytoplasmic proofreading systems for the ribosome

Given the importance to correctly translate proteins, an efficient quality control system must ensure that only functional ribosomes enter translation. In the nucleus, the TRAMP complex marks and targets aberrant pre-rRNAs for degradation by the nuclear exosome (Allmang et al. 2000; Dez et al. 2006; LaCava et al. 2005; Mitchell et al. 1997). The precise mechanism(s) of recognition of incorrectly assembled pre-ribosomal particles by the TRAMP complex in the nucleus remains unclear. Improperly assembled pre-ribosomal subunits that escape nuclear surveillance mechanisms are segregated and targeted for degradation in the cytoplasm by non-functional RNA decay (NRD) (Cole et al. 2009; Fujii et al. 2009, 2012; Graille and Seraphin 2012; LaRiviere et al. 2006). For a more detailed review on nuclear and cytoplasmic surveillance mechanisms for eukaryotic ribosome assembly, the readers are referred to Lafontaine (2010).

In the cytoplasm, two antagonistic mechanisms appear to proofread pre-ribosomes: (1) Assembly factors might either

prevent and/or delay pre-ribosomes from prematurely interacting or initiating translation. (2) They could actively check pre-ribosomal subunits for functionality. The ribosomal-like proteins (Rlp24 and Mrt4) act as placeholders for r-proteins. Therefore they delay maturation of the subunit and perhaps provide a time window for the functional proofreading of the 60S subunits. The assembly factor Tif6 is another example, whose binding to the 60S subunit interface prevents premature interactions with the 40S subunit. Tif6 is released from pre-60S subunits in the cytoplasm only after the formation of the acidic ribosomal stalk. Efl1 and Sdo1 promote the release of Tif6 from pre-60S ribosomes (Lo et al. 2010; Menne et al. 2007; Senger et al. 2001). Efl1 shares sequence similarity with the GTPase elongation factor eEF2. Additionally, Efl1 was proposed to check the integrity of the GTPase activating center, the P-site of the ribosome for functionality (Bussiere et al. 2012). Thus, cytoplasmic release factors may couple the recycling of shuttling assembly factors and simultaneously check ribosome function.

In the case of the pre-40S subunit, a number of assembly factors were proposed to block premature binding of initiation factors, mRNA, tRNA, and the 60S subunit. The cryo-EM structure of a late cytoplasmic pre-40S particle assigned the binding sites for Ltv1, Enp1, Rio2, Tsr1, Dim1, Pno1, and Nob1 on the small subunit (Strunk et al. 2011, Fig. 3). Their binding sites imply possible functions in preventing premature translation initiation by blocking access of translation factors. Ltv1 and Enp1 directly bind uS3 (Rps3) on its solvent exposed side thereby blocking the mRNA channel opening. Rio2, Tsr1, and Dim1 bind the subunit interface, thus preventing joining of the mature 60S subunit and translation initiation factor eIF1A. Nob1 and Pno1 block the binding of eIF3 and thereby interfere with translation initiation. After the release of Rio2, Tsr1, and Dim1, such a pre-40S particle may structurally mimic the translation initiation surface of a mature 40S subunit and hence can interact with a mature 60S subunit to form an 80S-like particle (Lebaron et al. 2012). This translation-like interaction could test the ability of a pre-40S subunit to bind 60S subunits. Additionally, by constantly interacting with each other in the cytoplasm, ribosomal subunits may sense their decoding ability to segregate and target aberrant particles for disassembly and degradation. Interestingly, Nob1 activity is stimulated by the translation initiation factor Fun12 (eIF5B) (Lebaron et al. 2012; Strunk et al. 2012). Thus, processing of the 20S pre-rRNA may represent a quality control mechanism that simultaneously triggers subunit maturation and senses translation competence.

Concluding remarks

Despite the initial description of eukaryotic ribosome assembly, which now dates back nearly 40 years, our understanding of this fundamental pathway still remains rudimentary. During

the last decades genetic screens, the development of visual reporters and proteomic approaches in budding yeast has provided crucial breakthroughs to further dissect eukaryotic ribosome biogenesis. These methodologies have expanded the inventory of assembly and transport factors that aid ribosome production. Deciphering the contribution of assembly factors during ribosome formation remains an important challenge. Structural approaches are beginning to facilitate these functional analyses of this highly dynamic pathway (Armache et al. 2010; Ben-Shem et al. 2011; Bradatsch et al. 2012; Greber et al. 2012; Klinge et al. 2011; Rabl et al. 2011). Moreover, these studies are guiding mechanistic analyses of this conserved pathway in higher eukaryotes (Tafforeau et al. 2013; Wild et al. 2010). Large-scale genome-wide visual screens and advanced imaging technologies will provide new insights into the assembly pathway of the human ribosome.

The importance of producing translation-competent ribosomes is reflected by the growing list of diseases that are linked to defects in ribosome assembly. Diamond-Blackfan anemia (DBA) is a rare human genetic disease. It causes bone marrow failure and severe anemia (Ellis and Lipton 2008; Ellis and Gleizes 2011). DBA is characterized by mutations in multiple proteins of the small and the large subunit and results in reduced translation capacity of the cell. Recently, mutations in the r-protein uL16 (Rpl10) and uL18 (Rpl5) were associated with T-cell acute lymphoblastic leukemia (De Keersmaecker et al. 2013). Mutations in the SSU component hUTP4 might be the basis of North American Indian childhood cirrhosis (Freed et al. 2012). The Shwachman-Bodian-Diamond syndrome arises from the inability to release Tif6 from pre-60S subunits (Finch et al. 2011; Menne et al. 2007; Wong et al. 2011).

Robust ribosome production allows cells to increase their mass and to proliferate, thus making ribosome assembly an important target for cancer treatments. Increased transcription of rDNA by RNA polymerase I is a characteristic feature of malignant cancers and to boost ribosome synthesis, emphasizing the importance of targeting the ribosome biogenesis pathway. Indeed, a recently developed small molecule inhibitor CX-5461 that targets rDNA transcription is able to selectively kill B-lymphoma cells (Bywater et al. 2012; Hannan et al. 2013). Unraveling the pathways and mechanisms by which eukaryotes build ribosomes will provide fundamental knowledge that will facilitate the rational design of therapeutics in the treatment of malignant cancers.

Acknowledgments We thank M. Altwater and U. Rothenbusch for comments and suggestions on the manuscript. J. J. Petkowski is supported by an ETH Zürich Postdoctoral Fellowship and the Marie Curie Actions for People COFUND program. V. G. Panse is supported by

grants from the Swiss National Science Foundation, the ETH Zürich, and a Starting Grant Award (EURIBIO260676) from the European Research Council.

References

- Allmang C, Kufel J, Chanfreau G, Mitchell P, Petfalski E, Tollervey D (1999) Functions of the exosome in rRNA, snoRNA and snRNA synthesis. *EMBO J* 18:5399–5410
- Allmang C, Mitchell P, Petfalski E, Tollervey D (2000) Degradation of ribosomal RNA precursors by the exosome. *Nucleic Acids Res* 28:1684–1691
- Altwater M, Chang Y, Melnik A, Occhipinti L, Schutz S, Rothenbusch U, Picotti P, Panse VG (2012) Targeted proteomics reveals compositional dynamics of 60S pre-ribosomes after nuclear export. *Mol Syst Biol* 8:628
- Armache JP, Jarasch A, Anger AM, Villa E, Becker T, Bhushan S, Jossinet F, Habeck M, Dindar G, Franckenberg S, Marquez V, Mielke T, Thomm M, Berninghausen O, Beatrix B, Soding J, Westhof E, Wilson DN, Beckmann R (2010) Cryo-EM structure and rRNA model of a translating eukaryotic 80S ribosome at 5.5-Å resolution. *Proc Natl Acad Sci U S A* 107:19748–19753
- Ban N, Beckmann R, Cate JH, Dinman JD, Dragon F, Ellis SR, Lafontaine DL, Lindahl L, Liljas A, Lipton JM, McAlear MA, Moore PB, Noller HF, Ortega J, Panse VG, Ramakrishnan V, Spahn CM, Steitz TA, Tchorzewski M, Tollervey D, Warren AJ, Williamson JR, Wilson D, Yonath A, Yusupov M (2014) A new system for naming ribosomal proteins. *Curr Opin Struct Biol*
- Bargis-Surgey P, Lavergne JP, Gonzalo P, Vard C, Filhol-Cochet O, Reboud JP (1999) Interaction of elongation factor eEF-2 with ribosomal P proteins. *Eur J Biochem FEBS* 262:606–611
- Bassler J, Grandi P, Gadal O, Lessmann T, Petfalski E, Tollervey D, Lechner J, Hurt E (2001) Identification of a 60S preribosomal particle that is closely linked to nuclear export. *Mol Cell Biol* 21:517–529
- Bassler J, Kallas M, Pertschy B, Ulbrich C, Thoms M, Hurt E (2010) The AAA-ATPase Real drives removal of biogenesis factors during multiple stages of 60S ribosome assembly. *Mol Cell* 38:712–721
- Bassler J, Klein I, Schmidt C, Kallas M, Thomson E, Wagner MA, Bradatsch B, Rechberger G, Strohmaier H, Hurt E, Bergler H (2012) The conserved Bud20 zinc finger protein is a new component of the ribosomal 60S subunit export machinery. *Mol Cell Biol* 32:4898–4912
- Baudin-Baillieu A, Tollervey D, Cullin C, Lacroute F (1997) Functional analysis of Rrp7p, an essential yeast protein involved in pre-rRNA processing and ribosome assembly. *Mol Cell Biol* 17:5023–5032
- Bayliss R, Leung SW, Baker RP, Quimby BB, Corbett AH, Stewart M (2002) Structural basis for the interaction between NTF2 and nucleoporin FxFG repeats. *EMBO J* 21:2843–2853
- Becam AM, Nasr F, Racki WJ, Zagulski M, Herbert CJ (2001) Rialp (Ynl163c), a protein similar to elongation factors 2, is involved in the biogenesis of the 60S subunit of the ribosome in *Saccharomyces cerevisiae*. *Mol Gen Genomics* 266:454–462
- Ben-Shem A, Garreau de Loubresse N, Melnikov S, Jenner L, Yusupova G, Yusupov M (2011) The structure of the eukaryotic ribosome at 3.0 Å resolution. *Science* (New York, NY) 334:1524–1529
- Bernstein KA, Gallagher JE, Mitchell BM, Granneman S, Baserga SJ (2004) The small-subunit processome is a ribosome assembly intermediate. *Eukaryot Cell* 3:1619–1626
- Billy E, Wegierski T, Nasr F, Filipowicz W (2000) Rcl1p, the yeast protein similar to the RNA 3'-phosphate cyclase, associates with U3 snoRNP and is required for 18S rRNA biogenesis. *EMBO J* 19:2115–2126
- Borovjagin AV, Gerbi SA (1999) U3 small nucleolar RNA is essential for cleavage at sites 1, 2 and 3 in pre-rRNA and determines which

- rRNA processing pathway is taken in *Xenopus* oocytes. *J Mol Biol* 286:1347–1363
- Borovjagin AV, Gerbi SA (2001) *Xenopus* U3 snoRNA GAC-Box A' and Box A sequences play distinct functional roles in rRNA processing. *Mol Cell Biol* 21:6210–6221
- Bradatsch B, Katahira J, Kowalinski E, Bange G, Yao W, Sekimoto T, Baumgartel V, Boese G, Bassler J, Wild K, Peters R, Yoneda Y, Sinning I, Hurt E (2007) Arx1 functions as an unorthodox nuclear export receptor for the 60S preribosomal subunit. *Mol Cell* 27:767–779
- Bradatsch B, Leidig C, Granneman S, Gnädig M, Tollervey D, Böttcher B, Beckmann R, Hurt E (2012) Structure of the pre-60S ribosomal subunit with nuclear export factor Arx1 bound at the exit tunnel. *Nat Struct Mol Biol* 19:1234–1241
- Briggs MW, Burkard KT, Butler JS (1998) Rrp6p, the yeast homologue of the human PM-Scl 100-kDa autoantigen, is essential for efficient 5.8S rRNA 3' end formation. *J Biol Chem* 273:13255–13263
- Britton RA (2009) Role of GTPases in bacterial ribosome assembly. *Annu Rev Microbiol* 63:155–176
- Bussiere C, Hashem Y, Arora S, Frank J, Johnson AW (2012) Integrity of the P-site is probed during maturation of the 60S ribosomal subunit. *J Cell Biol* 197:747–759
- Bywater MJ, Poortinga G, Sanij E, Hein N, Peck A, Cullinane C, Wall M, Cluse L, Drygin D, Anderes K, Huser N, Proffitt C, Bliesath J, Haddach M, Schwaebe MK, Ryckman DM, Rice WG, Schmitt C, Lowe SW, Johnstone RW, Pearson RB, McArthur GA, Hannan RD (2012) Inhibition of RNA polymerase I as a therapeutic strategy to promote cancer-specific activation of p53. *Cancer Cell* 22:51–65
- Champion EA, Lane BH, Jackrel ME, Regan L, Baserga SJ (2008) A direct interaction between the Utp6 half-a-tetratricopeptide repeat domain and a specific peptide in Utp21 is essential for efficient pre-rRNA processing. *Mol Cell Biol* 28:6547–6556
- Ciganda M, Williams N (2011) Eukaryotic 5S rRNA biogenesis. *Wiley Interdiscip Rev RNA* 2:523–533
- Clissold PM, Ponting CP (2000) PIN domains in nonsense-mediated mRNA decay and RNAi. *Curr Biol* 10:R888–R890
- Cole SE, LaRiviere FJ, Merrikh CN, Moore MJ (2009) A convergence of rRNA and mRNA quality control pathways revealed by mechanistic analysis of nonfunctional rRNA decay. *Mol Cell* 34:440–450
- Dalley JA, Selkirk A, Pool MR (2008) Access to ribosomal protein Rpl25p by the signal recognition particle is required for efficient cotranslational translocation. *Mol Biol Cell* 19:2876–2884
- De Keersmaecker K, Atak ZK, Li N, Vicente C, Patchett S, Girardi T, Gianfelici V, Geerdens E, Clappier E, Porcu M, Lahortiga I, Luca R, Yan J, Hulsemans G, Vranckx H, Vandepoel R, Sweron B, Jacobs K, Mentens N, Wlodarska I, Cauwelier B, Cloos J, Soulier J, Uytendaele A, Bagni C, Hassan BA, Vandenberghe P, Johnson AW, Aerts S, Cools J (2013) Exome sequencing identifies mutation in CNOT3 and ribosomal genes RPL5 and RPL10 in T-cell acute lymphoblastic leukemia. *Nat Genet* 45:186–190
- Decatur WA, Fournier MJ (2002) rRNA modifications and ribosome function. *Trends Biochem Sci* 27:344–351
- Decatur WA, Fournier MJ (2003) RNA-guided nucleotide modification of ribosomal and other RNAs. *J Biol Chem* 278:695–698
- Decatur WA, Liang XH, Piekna-Przybylska D, Fournier MJ (2007) Identifying effects of snoRNA-guided modifications on the synthesis and function of the yeast ribosome. *Methods Enzymol* 425:283–316
- Demoinet E, Jacquier A, Lutfalla G, Fromont-Racine M (2007) The Hsp40 chaperone Jjj1 is required for the nucleo-cytoplasmic recycling of preribosomal factors in *Saccharomyces cerevisiae*. *RNA (New York, NY)* 13:1570–1581
- Dez C, Houseley J, Tollervey D (2006) Surveillance of nuclear-restricted pre-ribosomes within a subnucleolar region of *Saccharomyces cerevisiae*. *EMBO J* 25:1534–1546
- Dosil M, Bustelo XR (2004) Functional characterization of Pwp2, a WD family protein essential for the assembly of the 90S pre-ribosomal particle. *J Biol Chem* 279:37385–37397
- Dragon F, Gallagher JE, Compagnone-Post PA, Mitchell BM, Porwancher KA, Wehner KA, Wormsley S, Settlege RE, Shabanowitz J, Osheim Y, Beyer AL, Hunt DF, Baserga SJ (2002) A large nucleolar U3 ribonucleoprotein required for 18S ribosomal RNA biogenesis. *Nature* 417:967–970
- Dunbar DA, Wormsley S, Agentis TM, Baserga SJ (1997) Mpp10p, a U3 small nucleolar ribonucleoprotein component required for pre-18S rRNA processing in yeast. *Mol Cell Biol* 17:5803–5812
- Ellis SR, Gleizes PE (2011) Diamond Blackfan anemia: ribosomal proteins going rogue. *Semin Hematol* 48:89–96
- Ellis SR, Lipton JM (2008) Diamond Blackfan anemia: a disorder of red blood cell development. *Curr Top Dev Biol* 82:217–241
- Englmeier L, Fomerod M, Bischoff FR, Petosa C, Mattaj IW, Kutay U (2001) RanBP3 influences interactions between CRM1 and its nuclear protein export substrates. *EMBO Rep* 2:926–932
- Faber AW, Van Dijk M, Raue HA, Vos JC (2002) Ngl2p is a Ccr4p-like RNA nuclease essential for the final step in 3'-end processing of 5.8S rRNA in *Saccharomyces cerevisiae*. *RNA (New York, NY)* 8:1095–1101
- Faza MB, Chang Y, Occhipinti L, Kemmler S, Panse VG (2012) Role of Mex67-Mtr2 in the nuclear export of 40S pre-ribosomes. *PLoS Genet* 8:e1002915
- Finch AJ, Hilcenko C, Basse N, Drynan LF, Goyenechea B, Menne TF, Gonzalez Fernandez A, Simpson P, D'Santos CS, Arends MJ, Donadieu J, Bellanne-Chantelot C, Costanzo M, Boone C, McKenzie AN, Freund SM, Warren AJ (2011) Uncoupling of GTP hydrolysis from eIF6 release on the ribosome causes Shwachman-Diamond syndrome. *Genes Dev* 25:917–929
- Freed EF, Prieto JL, McCann KL, McStay B, Baserga SJ (2012) NOL11, implicated in the pathogenesis of North American Indian childhood cirrhosis, is required for pre-rRNA transcription and processing. *PLoS Genet* 8:e1002892
- Fribourg S, Conti E (2003) Structural similarity in the absence of sequence homology of the messenger RNA export factors Mtr2 and p15. *EMBO Rep* 4:699–703
- Fromont-Racine M, Senger B, Saveanu C, Fasiolo F (2003) Ribosome assembly in eukaryotes. *Gene* 313:17–42
- Fujii K, Kitabatake M, Sakata T, Miyata A, Ohno M (2009) A role for ubiquitin in the clearance of nonfunctional rRNAs. *Genes Dev* 23:963–974
- Fujii K, Kitabatake M, Sakata T, Ohno M (2012) 40S subunit dissociation and proteasome-dependent RNA degradation in nonfunctional 25S rRNA decay. *EMBO J* 31:2579–2589
- Gadal O, Strauss D, Braspenning J, Hoepfner D, Petfalski E, Philippsen P, Tollervey D, Hurt E (2001a) A nuclear AAA-type ATPase (Rix7p) is required for biogenesis and nuclear export of 60S ribosomal subunits. *EMBO J* 20:3695–3704
- Gadal O, Strauss D, Kessl J, Trumpower B, Tollervey D, Hurt E (2001b) Nuclear export of 60s ribosomal subunits depends on Xpo1p and requires a nuclear export sequence-containing factor, Nmd3p, that associates with the large subunit protein Rpl10p. *Mol Cell Biol* 21:3405–3415
- Gallagher JE, Dunbar DA, Granneman S, Mitchell BM, Osheim Y, Beyer AL, Baserga SJ (2004) RNA polymerase I transcription and pre-rRNA processing are linked by specific SSU processome components. *Genes Dev* 18:2506–2517
- Garcia-Gomez JJ, Fernandez-Pevida A, Lebaron S, Rosado IV, Tollervey D, Kressler D, de la Cruz J (2014) Final pre-40S maturation depends on the functional integrity of the 60S subunit ribosomal protein L3. *PLoS Genet* 10:e1004205
- Geerlings TH, Vos JC, Raue HA (2000) The final step in the formation of 25S rRNA in *Saccharomyces cerevisiae* is performed by 5'→3' exonucleases. *RNA (New York, NY)* 6:1698–1703
- Geerlings TH, Faber AW, Bister MD, Vos JC, Raue HA (2003) Rio2p, an evolutionarily conserved, low abundant protein kinase essential for processing of 20S pre-rRNA in *Saccharomyces cerevisiae*. *J Biol Chem* 278:22537–22545

- Graille M, Seraphin B (2012) Surveillance pathways rescuing eukaryotic ribosomes lost in translation. *Nat Rev Mol Cell Biol* 13:727–735
- Grandi P, Rybin V, Bassler J, Petfalski E, Strauss D, Marzioch M, Schafer T, Kuster B, Tschochner H, Tollervey D, Gavin AC, Hurt E (2002) 90S pre-ribosomes include the 35S pre-rRNA, the U3 snoRNP, and 40S subunit processing factors but predominantly lack 60S synthesis factors. *Mol Cell* 10:105–115
- Granneman S, Gallagher JE, Vogelzangs J, Horstman W, van Venrooij WJ, Baserga SJ, Pruijn GJ (2003) The human Imp3 and Imp4 proteins form a ternary complex with hMpp10, which only interacts with the U3 snoRNA in 60–80S ribonucleoprotein complexes. *Nucleic Acids Res* 31:1877–1887
- Greber BJ, Boehringer D, Montellese C, Ban N (2012) Cryo-EM structures of Arx1 and maturation factors Reil and Jjj1 bound to the 60S ribosomal subunit. *Nat Struct Mol Biol* 19:1228–1233
- Hackmann A, Gross T, Baierlein C, Krebber H (2011) The mRNA export factor Npl3 mediates the nuclear export of large ribosomal subunits. *EMBO Rep* 12:1024–1031
- Hannan RD, Drygin D, Pearson RB (2013) Targeting RNA polymerase I transcription and the nucleolus for cancer therapy. *Expert Opin Ther Targets* 17:873–878
- Hampicharnchai P, Jakovljevic J, Horsey E, Miles T, Roman J, Rout M, Meagher D, Imai B, Guo Y, Brame CJ, Shabanowitz J, Hunt DF, Woolford JL Jr (2001) Composition and functional characterization of yeast 66S ribosome assembly intermediates. *Mol Cell* 8:505–515
- Hedges J, West M, Johnson AW (2005) Release of the export adapter, Nmd3p, from the 60S ribosomal subunit requires Rpl10p and the cytoplasmic GTPase Lsg1p. *EMBO J* 24:567–579
- Henry Y, Wood H, Morrissey JP, Petfalski E, Kearsley S, Tollervey D (1994) The 5' end of yeast 5.8S rRNA is generated by exonucleases from an upstream cleavage site. *EMBO J* 13:2452–2463
- Ho JH, Kallstrom G, Johnson AW (2000) Nascent 60S ribosomal subunits enter the free pool bound by Nmd3p. *RNA (New York, NY)* 6:1625–1634
- Hoelz A, Debler EW, Blobel G (2011) The structure of the nuclear pore complex. *Annu Rev Biochem* 80:613–643
- Hughes JM (1996) Functional base-pairing interaction between highly conserved elements of U3 small nucleolar RNA and the small ribosomal subunit RNA. *J Mol Biol* 259:645–654
- Hughes JM, Ares M Jr (1991) Depletion of U3 small nucleolar RNA inhibits cleavage in the 5' external transcribed spacer of yeast pre-ribosomal RNA and impairs formation of 18S ribosomal RNA. *EMBO J* 10:4231–4239
- Hung NJ, Johnson AW (2006) Nuclear recycling of the pre-60S ribosomal subunit-associated factor Arx1 depends on Reil in *Saccharomyces cerevisiae*. *Mol Cell Biol* 26:3718–3727
- Hung NJ, Lo KY, Patel SS, Helmke K, Johnson AW (2008) Arx1 is a nuclear export receptor for the 60S ribosomal subunit in yeast. *Mol Biol Cell* 19:735–744
- Hurt E, Hannus S, Schmelz B, Lau D, Tollervey D, Simos G (1999) A novel in vivo assay reveals inhibition of ribosomal nuclear export in ran-cycle and nucleoporin mutants. *J Cell Biol* 144:389–401
- Jakovljevic J, de Mayolo PA, Miles TD, Nguyen TM, Leger-Silvestre I, Gas N, Woolford JL Jr (2004) The carboxy-terminal extension of yeast ribosomal protein S14 is necessary for maturation of 43S pre-ribosomes. *Mol Cell* 14:331–342
- Kappel L, Loibl M, Zisser G, Klein I, Fruhmann G, Gruber C, Unterwieser S, Rechberger G, Pertschy B, Bergler H (2012) Rlp24 activates the AAA-ATPase Drg1 to initiate cytoplasmic pre-60S maturation. *J Cell Biol* 199:771–782
- Karbstein K, Jonas S, Doudna JA (2005) An essential GTPase promotes assembly of pre-ribosomal RNA processing complexes. *Mol Cell* 20:633–643
- Karl T, Onder K, Kodzius R, Pichova A, Wimmer H, Th r A, Hundsberger H, Loffler M, Klade T, Beyer A, Breitenbach M, Koller L (1999) GRC5 and NMD3 function in translational control of gene expression and interact genetically. *Curr Genet* 34:419–429
- Kemmler S, Occhipinti L, Veisu M, Panse VG (2009) Yvh1 is required for a late maturation step in the 60S biogenesis pathway. *J Cell Biol* 186:863–880
- Kiss T (2001) Small nucleolar RNA-guided post-transcriptional modification of cellular RNAs. *EMBO J* 20:3617–3622
- Klinge S, Voigts-Hoffmann F, Leibundgut M, Arpagaus S, Ban N (2011) Crystal structure of the eukaryotic 60S ribosomal subunit in complex with initiation factor 6. *Science (New York, NY)* 334:941–948
- Kos M, Tollervey D (2010) Yeast pre-rRNA processing and modification occur cotranscriptionally. *Mol Cell* 37:809–820
- Kressler D, Linder P, de La Cruz J (1999) Protein trans-acting factors involved in ribosome biogenesis in *Saccharomyces cerevisiae*. *Mol Cell Biol* 19:7897–7912
- Kressler D, Roser D, Pertschy B, Hurt E (2008) The AAA ATPase Rix7 powers progression of ribosome biogenesis by stripping Nsa1 from pre-60S particles. *J Cell Biol* 181:935–944
- Kressler D, Bange G, Ogawa Y, Stjepanovic G, Bradatsch B, Pratte D, Amlacher S, Strauss D, Yoneda Y, Katahira J, Sinning I, Hurt E (2012a) Synchronizing nuclear import of ribosomal proteins with ribosome assembly. *Science (New York, NY)* 338:666–671
- Kressler D, Hurt E, Bergler H, Bassler J (2012b) The power of AAA-ATPases on the road of pre-60S ribosome maturation—molecular machines that strip pre-ribosomal particles. *Biochim Biophys Acta* 1823:92–100
- Krogan NJ, Kim M, Ahn SH, Zhong G, Kobor MS, Cagney G, Emili A, Shilatifard A, Buratowski S, Greenblatt JF (2002) RNA polymerase II elongation factors of *Saccharomyces cerevisiae*: a targeted proteomics approach. *Mol Cell Biol* 22:6979–6992
- Kufel J, Dichtl B, Tollervey D (1999) Yeast Rnt1p is required for cleavage of the pre-ribosomal RNA in the 3' ETS but not the 5' ETS. *RNA (New York, NY)* 5:909–917
- LaCava J, Houseley J, Saveanu C, Petfalski E, Thompson E, Jacquier A, Tollervey D (2005) RNA degradation by the exosome is promoted by a nuclear polyadenylation complex. *Cell* 121:713–724
- Lafontaine DL (2010) A 'garbage can' for ribosomes: how eukaryotes degrade their ribosomes. *Trends Biochem Sci* 35:267–277
- Lafontaine DL, Tollervey D (2000) Synthesis and assembly of the box C+D small nucleolar RNPs. *Mol Cell Biol* 20:2650–2659
- Lafontaine D, Delcour J, Glasser AL, Desgres J, Vandenhautte J (1994) The DIM1 gene responsible for the conserved m(6)Am(6)A dimethylation in the 3'-terminal loop of 18S rRNA is essential in yeast. *J Mol Biol* 241:492–497
- Lafontaine DL, Preiss T, Tollervey D (1998) Yeast 18S rRNA dimethylase Dim1p: a quality control mechanism in ribosome synthesis? *Mol Cell Biol* 18:2360–2370
- Lamanna AC, Karbstein K (2009) Nob1 binds the single-stranded cleavage site D at the 3'-end of 18S rRNA with its PIN domain. *Proc Natl Acad Sci U S A* 106:14259–14264
- LaRiviere FJ, Cole SE, Ferullo DJ, Moore MJ (2006) A late-acting quality control process for mature eukaryotic rRNAs. *Mol Cell* 24:619–626
- Lebaron S, Schneider C, van Nues RW, Swiatkowska A, Walsh D, Bottcher B, Granneman S, Watkins NJ, Tollervey D (2012) Proofreading of pre-40S ribosome maturation by a translation initiation factor and 60S subunits. *Nat Struct Mol Biol* 19:744–753
- Lebreton A, Saveanu C, Decourty L, Rain JC, Jacquier A, Fromont-Racine M (2006) A functional network involved in the recycling of nucleocytoplasmic pre-60S factors. *J Cell Biol* 173:349–360
- Lee SJ, Baserga SJ (1999) Imp3p and Imp4p, two specific components of the U3 small nucleolar ribonucleoprotein that are essential for pre-18S rRNA processing. *Mol Cell Biol* 19:5441–5452
- Li Z, Lee I, Moradi E, Hung NJ, Johnson AW, Marcotte EM (2009) Rational extension of the ribosome biogenesis pathway using network-guided genetics. *PLoS Biol* 7:e1000213

- Lim YH, Charette JM, Baserga SJ (2011) Assembling a protein-protein interaction map of the SSU processome from existing datasets. *PLoS ONE* 6:e17701
- Lischwe MA, Ochs RL, Reddy R, Cook RG, Yeoman LC, Tan EM, Reichlin M, Busch H (1985) Purification and partial characterization of a nucleolar scleroderma antigen (Mr=34,000; pI, 8.5) rich in NG, NG-dimethylarginine. *J Biol Chem* 260:14304–14310
- Lo KY, Li Z, Wang F, Marcotte EM, Johnson AW (2009) Ribosome stalk assembly requires the dual-specificity phosphatase Yvh1 for the exchange of Mrt4 with P0. *J Cell Biol* 186:849–862
- Lo KY, Li Z, Bussièrè C, Bresson S, Marcotte EM, Johnson AW (2010) Defining the pathway of cytoplasmic maturation of the 60S ribosomal subunit. *Mol Cell* 39:196–208
- Lukowiak AA, Granneman S, Mattox SA, Speckmann WA, Jones K, Pluk H, Venrooij WJ, Terns RM, Terns MP (2000) Interaction of the U3-55k protein with U3 snoRNA is mediated by the box B/C motif of U3 and the WD repeats of U3-55k. *Nucleic Acids Res* 28:3462–3471
- Lupas AN, Martin J (2002) AAA proteins. *Curr Opin Struct Biol* 12:746–753
- Lygerou Z, Allmang C, Tollervey D, Seraphin B (1996) Accurate processing of a eukaryotic precursor ribosomal RNA by ribonuclease MRP in vitro. *Science (New York, NY)* 272:268–270
- Marmier-Gourrier N, Clery A, Schlotter F, Senty-Segault V, Branlant C (2011) A second base pair interaction between U3 small nucleolar RNA and the 5'-ETS region is required for early cleavage of the yeast pre-ribosomal RNA. *Nucleic Acids Res* 39:9731–9745
- Matsuo Y, Granneman S, Thoms M, Manikas RG, Tollervey D, Hurt E (2014) Coupled GTPase and remodelling ATPase activities form a checkpoint for ribosome export. *Nature* 505:112–116
- Melnikov S, Ben-Shem A, Garreau de Loubresse N, Jenner L, Yusupova G, Yusupov M (2012) One core, two shells: bacterial and eukaryotic ribosomes. *Nat Struct Mol Biol* 19:560–567
- Menne TF, Goyenechea B, Sanchez-Puig N, Wong CC, Tonkin LM, Ancliff PJ, Brost RL, Costanzo M, Boone C, Warren AJ (2007) The Shwachman-Bodian-Diamond syndrome protein mediates translational activation of ribosomes in yeast. *Nat Genet* 39:486–495
- Meyer AE, Hung NJ, Yang P, Johnson AW, Craig EA (2007) The specialized cytosolic J-protein, Jj1, functions in 60S ribosomal subunit biogenesis. *Proc Natl Acad Sci U S A* 104:1558–1563
- Meyer AE, Hoover LA, Craig EA (2010) The cytosolic J-protein, Jj1, and Rei1 function in the removal of the pre-60S subunit factor Arx1. *J Biol Chem* 285:961–968
- Meyer H, Bug M, Bremer S (2012) Emerging functions of the VCP/p97 AAA-ATPase in the ubiquitin system. *Nat Cell Biol* 14:117–123
- Milkereit P, Gadal O, Podtelejnikov A, Trumtel S, Gas N, Petfalski E, Tollervey D, Mann M, Hurt E, Tschochner H (2001) Maturation and intranuclear transport of pre-ribosomes requires Noc proteins. *Cell* 105:499–509
- Milkereit P, Strauss D, Bassler J, Gadal O, Kuhn H, Schutz S, Gas N, Lechner J, Hurt E, Tschochner H (2003) A Noc complex specifically involved in the formation and nuclear export of ribosomal 40 S subunits. *J Biol Chem* 278:4072–4081
- Mitchell P, Petfalski E, Shevchenko A, Mann M, Tollervey D (1997) The exosome: a conserved eukaryotic RNA processing complex containing multiple 3'→5' exoribonucleases. *Cell* 91:457–466
- Moy TI, Silver PA (2002) Requirements for the nuclear export of the small ribosomal subunit. *J Cell Sci* 115:2985–2995
- Ng CL, Waterman D, Koonin EV, Antson AA, Ortiz-Lombardia M (2005) Crystal structure of Mil (Mth680): internal duplication and similarity between the Imp4/Brix domain and the anticodon-binding domain of class IIa aminoacyl-tRNA synthetases. *EMBO Rep* 6:140–146
- Nissan TA, Bassler J, Petfalski E, Tollervey D, Hurt E (2002) 60S pre-ribosome formation viewed from assembly in the nucleolus until export to the cytoplasm. *EMBO J* 21:5539–5547
- Occhipinti L, Chang Y, Altvater M, Menet AM, Kemmler S, Panse VG (2013) Non-FG mediated transport of the large pre-ribosomal subunit through the nuclear pore complex by the mRNA export factor Gle2. *Nucleic Acids Res* 41:8266–8279
- Oeffinger M, Dlakic M, Tollervey D (2004) A pre-ribosome-associated HEAT-repeat protein is required for export of both ribosomal subunits. *Genes Dev* 18:196–209
- Oeffinger M, Zenklusen D, Ferguson A, Wei KE, El Hage A, Tollervey D, Chait BT, Singer RH, Rout MP (2009) Rrp17p is a eukaryotic exonuclease required for 5' end processing of pre-60S ribosomal RNA. *Mol Cell* 36:768–781
- Osheim YN, French SL, Keck KM, Champion EA, Spasov K, Dragon F, Baserga SJ, Beyer AL (2004) Pre-18S ribosomal RNA is structurally compacted into the SSU processome prior to being cleaved from nascent transcripts in *Saccharomyces cerevisiae*. *Mol Cell* 16:943–954
- Panse VG, Johnson AW (2010) Maturation of eukaryotic ribosomes: acquisition of functionality. *Trends Biochem Sci* 35:260–266
- Panse VG, Kressler D, Pauli A, Petfalski E, Gnadig M, Tollervey D, Hurt E (2006) Formation and nuclear export of preribosomes are functionally linked to the small-ubiquitin-related modifier pathway. *Traffic (Copenhagen, Denmark)* 7:1311–1321
- Peng WT, Robinson MD, Mnaimneh S, Krogan NJ, Cagney G, Morris Q, Davierwala AP, Grigull J, Yang X, Zhang W, Mitsakakis N, Ryan OW, Datta N, Jovic V, Pal C, Canadian V, Richards D, Beattie B, Wu LF, Altschuler SJ, Roweis S, Frey BJ, Emili A, Greenblatt JF, Hughes TR (2003) A panoramic view of yeast noncoding RNA processing. *Cell* 113:919–933
- Perez-Fernandez J, Roman A, De Las Rivas J, Bustelo XR, Dosil M (2007) The 90S preribosome is a multimodular structure that is assembled through a hierarchical mechanism. *Mol Cell Biol* 27:5414–5429
- Perez-Fernandez J, Martin-Marcos P, Dosil M (2011) Elucidation of the assembly events required for the recruitment of Utp20, Imp4 and Bms1 onto nascent pre-ribosomes. *Nucleic Acids Res* 39:8105–8121
- Pertschy B, Saveanu C, Zisser G, Lebreton A, Teng M, Jacquier A, Liebming E, Nobis B, Kappel L, van der Klei I, Hogenauer G, Fromont-Racine M, Bergler H (2007) Cytoplasmic recycling of 60S preribosomal factors depends on the AAA protein Drg1. *Mol Cell Biol* 27:6581–6592
- Pertschy B, Schneider C, Gnadig M, Schafer T, Tollervey D, Hurt E (2009) RNA helicase Prp43 and its co-factor Pfa1 promote 20 to 18 S rRNA processing catalyzed by the endonuclease Nob1. *J Biol Chem* 284:35079–35091
- Phipps KR, Charette J, Baserga SJ (2011) The small subunit processome in ribosome biogenesis—progress and prospects. *Wiley Interdiscip Rev RNA* 2:1–21
- Punta M, Coghill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, Pang N, Forslund K, Ceric G, Clements J, Heger A, Holm L, Sonnhammer EL, Eddy SR, Bateman A, Finn RD (2012) The Pfam protein families database. *Nucleic Acids Res* 40:D290–D301
- Rabl J, Leibundgut M, Ataïde SF, Haag A, Ban N (2011) Crystal structure of the eukaryotic 40S ribosomal subunit in complex with initiation factor 1. *Science (New York, NY)* 331:730–736
- Raices M, D'Angelo MA (2012) Nuclear pore complex composition: a new regulator of tissue-specific and developmental functions. *Nat Rev Mol Cell Biol* 13:687–699
- Rudra D, Mallick J, Zhao Y, Warner JR (2007) Potential interface between ribosomal protein production and pre-rRNA processing. *Mol Cell Biol* 27:4815–4824
- Russell DW, Spremulli LL (1979) Purification and characterization of a ribosome dissociation factor (eukaryotic initiation factor 6) from wheat germ. *J Biol Chem* 254:8796–8800
- Samanta MP, Liang S (2003) Predicting protein functions from redundancies in large-scale protein interaction networks. *Proc Natl Acad Sci U S A* 100:12579–12583
- Santos-Rosa H, Moreno H, Simos G, Segref A, Fahrenkrog B, Pante N, Hurt E (1998) Nuclear mRNA export requires complex formation

- between Mex67p and Mtr2p at the nuclear pores. *Mol Cell Biol* 18: 6826–6838
- Schafer T, Strauss D, Petfalski E, Tollervey D, Hurt E (2003) The path from nucleolar 90S to cytoplasmic 40S pre-ribosomes. *EMBO J* 22: 1370–1380
- Schafer T, Maco B, Petfalski E, Tollervey D, Bottcher B, Aebi U, Hurt E (2006) Hrr25-dependent phosphorylation state regulates organization of the pre-40S subunit. *Nature* 441:651–655
- Segref A, Sharma K, Doye V, Hellwig A, Huber J, Luhrmann R, Hurt E (1997) Mex67p, a novel factor for nuclear mRNA export, binds to both poly(A)+ RNA and nuclear pores. *EMBO J* 16:3256–3271
- Senay C, Ferrari P, Rocher C, Rieger KJ, Winter J, Platel D, Bourne Y (2003) The Mtr2-Mex67 NTF2-like domain complex. Structural insights into a dual role of Mtr2 for yeast nuclear export. *J Biol Chem* 278:48395–48403
- Senger B, Lafontaine DL, Graindorge JS, Gadal O, Camasses A, Sanni A, Garnier JM, Breitenbach M, Hurt E, Fasiolo F (2001) The nucleolar Tif6p and Efl1p are required for a late cytoplasmic step of ribosome synthesis. *Mol Cell* 8:1363–1373
- Sengupta J, Bussiere C, Pallesen J, West M, Johnson AW, Frank J (2010) Characterization of the nuclear export adaptor protein Nmd3 in association with the 60S ribosomal subunit. *J Cell Biol* 189:1079–1086
- Shajani Z, Sykes MT, Williamson JR (2011) Assembly of bacterial ribosomes. *Annu Rev Biochem* 80:501–526
- Shimizu K, Kawasaki Y, Hiraga S, Tawaramoto M, Nakashima N, Sugino A (2002) The fifth essential DNA polymerase phi in *Saccharomyces cerevisiae* is localized to the nucleolus and plays an important role in synthesis of rRNA. *Proc Natl Acad Sci U S A* 99:9133–9138
- Stage-Zimmermann T, Schmidt U, Silver PA (2000) Factors affecting nuclear export of the 60S ribosomal subunit in vivo. *Mol Biol Cell* 11:3777–3789
- Strasser K, Bassler J, Hurt E (2000) Binding of the Mex67p/Mtr2p heterodimer to FXFG, GLFG, and FG repeat nucleoporins is essential for nuclear mRNA export. *J Cell Biol* 150:695–706
- Strawn LA, Shen T, Wentz SR (2001) The GLFG regions of Nup116p and Nup100p serve as binding sites for both Kap95p and Mex67p at the nuclear pore complex. *J Biol Chem* 276:6445–6452
- Strub BR, Eswara MB, Pierce JB, Mangroo D (2007) Utp8p is a nucleolar tRNA-binding protein that forms a complex with components of the nuclear tRNA export machinery in *Saccharomyces cerevisiae*. *Mol Biol Cell* 18:3845–3859
- Strunk BS, Loucks CR, Su M, Vashisth H, Cheng S, Schilling J, Brooks CL 3rd, Karbstein K, Skiniotis G (2011) Ribosome assembly factors prevent premature translation initiation by 40S assembly intermediates. *Science (New York, NY)* 333:1449–1453
- Strunk BS, Novak MN, Young CL, Karbstein K (2012) A translation-like cycle is a quality control checkpoint for maturing 40S ribosome subunits. *Cell* 150:111–121
- Tafforeau L, Zorbas C, Langhendries JL, Mullineux ST, Stamatopoulou V, Mullier R, Wacheul L, Lafontaine DL (2013) The complexity of human ribosome biogenesis revealed by systematic nucleolar screening of Pre-rRNA processing factors. *Mol Cell* 51:539–551
- Tanaka N, Smith P, Shuman S (2011) Crystal structure of Rcl1, an essential component of the eukaryal pre-rRNA processosome implicated in 18S rRNA biogenesis. *RNA (New York, NY)* 17:595–602
- Taura T, Krebber H, Silver PA (1998) A member of the Ran-binding protein family, Yrb2p, is involved in nuclear protein export. *Proc Natl Acad Sci U S A* 95:7427–7432
- Thomson E, Tollervey D (2010) The final step in 5.8S rRNA processing is cytoplasmic in *Saccharomyces cerevisiae*. *Mol Cell Biol* 30:976–984
- Tollervey D, Lehtonen H, Carmo-Fonseca M, Hurt EC (1991) The small nucleolar RNP protein NOP1 (fibrillarin) is required for pre-rRNA processing in yeast. *EMBO J* 10:573–583
- Trapman J, Retel J, Planta RJ (1975) Ribosomal precursor particles from yeast. *Exp Cell Res* 90:95–104
- Tuck AC, Tollervey D (2013) A transcriptome-wide atlas of RNP composition reveals diverse classes of mRNAs and lncRNAs. *Cell* 154: 996–1009
- Udem SA, Warner JR (1972) Ribosomal RNA synthesis in *Saccharomyces cerevisiae*. *J Mol Biol* 65:227–242
- Ulbrich C, Diepholz M, Bassler J, Kressler D, Pertschy B, Galani K, Bottcher B, Hurt E (2009) Mechanochemical removal of ribosome biogenesis factors from nascent 60S ribosomal subunits. *Cell* 138:911–922
- Valenzuela DM, Chaudhuri A, Maitra U (1982) Eukaryotic ribosomal subunit anti-association activity of calf liver is contained in a single polypeptide chain protein of Mr=25,500 (eukaryotic initiation factor 6). *J Biol Chem* 257:7712–7719
- van Hoof A, Lennertz P, Parker R (2000) Three conserved members of the RNase D family have unique and overlapping functions in the processing of 5S, 5.8S, U4, U5, RNase MRP and RNase P RNAs in yeast. *EMBO J* 19:1357–1365
- Vanrobays E, Gelugne JP, Gleizes PE, Caizergues-Ferrer M (2003) Late cytoplasmic maturation of the small ribosomal subunit requires RIO proteins in *Saccharomyces cerevisiae*. *Mol Cell Biol* 23:2083–2095
- Venema J, Tollervey D (1995) Processing of pre-ribosomal RNA in *Saccharomyces cerevisiae*. *Yeast (Chichester, England)* 11:1629–1650
- Venema J, Vos HR, Faber AW, van Venrooij WJ, Raue HA (2000) Yeast Rrp9p is an evolutionarily conserved U3 snoRNP protein essential for early pre-rRNA processing cleavages and requires box C for its association. *RNA (New York, NY)* 6:1660–1671
- Vos HR, Bax R, Faber AW, Vos JC, Raue HA (2004) U3 snoRNP and Rrp5p associate independently with *Saccharomyces cerevisiae* 35S pre-rRNA, but Rrp5p is essential for association of Rok1p. *Nucleic Acids Res* 32:5827–5833
- Warner JR (1999) The economics of ribosome biosynthesis in yeast. *Trends Biochem Sci* 24:437–440
- Watkins NJ, Bohnsack MT (2012) The box C/D and H/ACA snoRNPs: key players in the modification, processing and the dynamic folding of ribosomal RNA. *Wiley Interdiscip Rev RNA* 3:397–414
- Watkins NJ, Segault V, Charpentier B, Nottrott S, Fabrizio P, Bachi A, Wilm M, Rosbash M, Branlant C, Luhrmann R (2000) A common core RNP structure shared between the small nucleolar box C/D RNPs and the spliceosomal U4 snRNP. *Cell* 103:457–466
- Wegierski T, Billy E, Nasr F, Filipowicz W (2001) Bms1p, a G-domain-containing protein, associates with Rcl1p and is required for 18S rRNA biogenesis in yeast. *RNA (New York, NY)* 7:1254–1267
- Wehner KA, Gallagher JE, Baserga SJ (2002) Components of an interdependent unit within the SSU processome regulate and mediate its activity. *Mol Cell Biol* 22:7258–7267
- Wentz SR, Rout MP (2010) The nuclear pore complex and nuclear transport. *Cold Spring Harb Perspect Biol* 2:a000562
- West M, Hedges JB, Chen A, Johnson AW (2005) Defining the order in which Nmd3p and Rpl10p load onto nascent 60S ribosomal subunits. *Mol Cell Biol* 25:3802–3813
- Wild T, Horvath P, Wyler E, Widmann B, Badertscher L, Zemp I, Kozak K, Csucs G, Lund E, Kutay U (2010) A protein inventory of human ribosome biogenesis reveals an essential function of exportin 5 in 60S subunit export. *PLoS Biol* 8:e1000522
- Wong CC, Traynor D, Basse N, Kay RR, Warren AJ (2011) Defective ribosome assembly in Shwachman-Diamond syndrome. *Blood* 118: 4305–4312
- Woolford JL Jr, Baserga SJ (2013) Ribosome biogenesis in the yeast *Saccharomyces cerevisiae*. *Genetics* 195:643–681
- Wu P, Brockenbrough JS, Metcalfe AC, Chen S, Aris JP (1998) Nop5p is a small nucleolar ribonucleoprotein component required for pre-18S rRNA processing in yeast. *J Biol Chem* 273:16453–16463
- Yao W, Roser D, Kohler A, Bradatsch B, Bassler J, Hurt E (2007) Nuclear export of ribosomal 60S subunits by the general mRNA export receptor Mex67-Mtr2. *Mol Cell* 26:51–62

- Yao Y, Demoinet E, Saveanu C, Lenormand P, Jacquier A, Fromont-Racine M (2010) Ecm1 is a new pre-ribosomal factor involved in pre-60S particle export. *RNA* (New York, NY) 16:1007–1017
- Yip WS, Vincent NG, Baserga SJ (2013) Ribonucleoproteins in archaeal pre-rRNA processing and modification. *Archaea* (Vancouver, BC) 2013:614735
- Zemp I, Wild T, O'Donohue MF, Wandrey F, Widmann B, Gleizes PE, Kutay U (2009) Distinct cytoplasmic maturation steps of 40S ribosomal subunit precursors require hRio2. *J Cell Biol* 185:1167–1180
- Zhang J, Harnpicharnchai P, Jakovljevic J, Tang L, Guo Y, Oeffinger M, Rout MP, Hiley SL, Hughes T, Woolford JL Jr (2007) Assembly factors Rpf2 and Rrs1 recruit 5S rRNA and ribosomal proteins rpL5 and rpL11 into nascent ribosomes. *Genes Dev* 21:2580–2592
- Zwarger M, Medalia O (2012) Unravelling the lamina network. *Nat Rev Mol Cell Biol* 13:140