A set of new and cross-amplifying microsatellite loci for conservation genetics of the endangered stone crayfish (Austropotamobius torrentium)

Journal Article

Author(s): Vorburger, Christoph; Rhyner, Nicola; Hartikainen, Hanna; Jokela, Jukka

Publication date: 2014

Permanent link: https://doi.org/10.3929/ethz-b-000088701

Rights / license: In Copyright - Non-Commercial Use Permitted

Originally published in: Conservation Genetics Resources 6(3), <u>https://doi.org/10.1007/s12686-014-0157-0</u>

MICROSATELLITE LETTERS

A set of new and cross-amplifying microsatellite loci for conservation genetics of the endangered stone crayfish (*Austropotamobius torrentium*)

Christoph Vorburger • Nicola Rhyner • Hanna Hartikainen • Jukka Jokela

Received: 30 January 2014/Accepted: 1 February 2014/Published online: 7 February 2014 © Springer Science+Business Media Dordrecht 2014

Abstract Like other indigenous crayfish species in Europe, the stone crayfish (*Austropotamobius torrentium*) is endangered due to habitat degradation, competition with invasive American crayfish and a deadly disease (crayfish plague) vectored by these invaders. To study the genetic diversity and connectivity of the remnant stone crayfish populations, eight new microsatellite loci were developed and supplemented by four cross-amplifying loci developed for related species. These twelve loci were tested for polymorphism in 130 stone crayfish from five streams in Switzerland. Variability was low with 1–7 alleles per locus, but the markers revealed strong genetic differentiation among streams. Use of these microsatellites will provide important information for the conservation and management of this endangered species.

Keywords Austropotamobius torrentium · Habitat fragmentation · Microsatellites · Stone crayfish

Microsatellite sequences were isolated by ecogenics GmbH (Switzerland) using the high-throughput genomic sequencing approach detailed in Abdelkrim et al. (2009). Genomic DNA (11 μ g) of a single stone crayfish was analyzed on a Roche

454 GS-FLX platform (Roche, Switzerland) using a 1/16th run and the GS FLX Titanium reagents. The 50,529 reads obtained had an average length of 257 bp. Microsatellite inserts suitable for primer design with a tetra- or a trinucleotide motif of at least 6 units or a dinucleotide motif of at least 11 units were present in 137 sequences. Primers were designed for 30 loci and tested in 15 unrelated stone crayfish. Only four primer pairs produced interpretable and polymorphic amplicons of the expected size. The remaining sequences obtained from ecogenics GmbH were therefore subjected to another search using the QDD software (Meglécz et al. 2010) and we designed and tested an additional 23 primer pairs on the same individuals. Of these, again four produced scorable products of variable size. We also tested 13 primer pairs developed for the congeneric species A. pallipes and A. italicus (Pedraza-Lara et al. 2011; Gouin et al. 2000), of which four yielded interpretable amplification products. The 12 usable microsatellite loci so obtained were then grouped into three multiplex reactions according to annealing temperatures and product sizes to genotype 130 stone crayfish from five streams in northeastern Switzerland (Table 1). Three streams were in close proximity of each other (<1 km) but separated by man-made migration barriers, the other two streams were more distant (20-50 km) and located in different drainages. DNA was extracted from clipped crayfish pleopods using a high salt protocol (Sunnucks and Hales 1996). PCRs were performed in 11 µl final volumes, using the Qiagen multiplex kit and the following thermal profile: 95 °C for 15 min, 35 cycles at 94 °C for 30 s, 1:30 min annealing at 52, 61 or 60 °C (for multiplex I, II or III, respectively) and 1:30 min at 72 °C, and a final extension at 60 °C for 30 min. The forward primer of each pair was labeled with a fluorophore as detailed in Table 1, PCR products were visualized on an ABI 3130 Genetic Analyzer and alleles were sized using GeneMapper[®] (Applied Biosystems).

C. Vorburger · J. Jokela

Institute of Integrative Biology, ETH Zürich, Universitätstrasse 16, 8092 Zurich, Switzerland

C. Vorburger $(\boxtimes) \cdot N$. Rhyner $\cdot H$. Hartikainen $\cdot J$. Jokela EAWAG, Swiss Federal Institute of Aquatic Science and Technology, Überlandstrasse 133, 8600 Dübendorf, Switzerland e-mail: christoph.vorburger@eawag.ch

H. Hartikainen

Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

Table 1 (individual:	Characterization of eight novel micr s from five streams in northeastern S	osatellite loci for stone crayfish (Austropotamobius torren witzerland	ntium) and four cross-	amplifying lo	ci from	congeneric species in a	a sample	of 130
Locus	GenBank accession no./reference	Primer sequence $(5'-3')$ [fluorophore]	Motif	Multiplex	$N_{\rm A}$	Allele size range (bp)	$H_{\rm O}$	$H_{\rm E}$
Newly dev AT1	veloped loci			=	6	108 210	0 503	0 506
	7 (70) (1 X	R: CAAGTAAGGGCCGGGTGAG	9(200)8(000)	=	n	017-071		
AT9	KJ150253	F: ACGTGTGTCTGTCTCTATAAATTC [ATT0565]	$(CTGT)_7$	Ш	2	204-217	0.210	0.213
		R: ATTAATTACGAGGAGGAACAAGAG						
AT22	KJ150254	F: GCACCCCGTTTTCTATGCG [YYE]	$(AAT)_8$	II	2	241-244	0.225	0.226
		R: AGCAAGGTACACCTCTCGG						
AT23	KJ150255	F: CTGTTTCTTGGGTCTCGGC [ATT0565]	(TGC) ₈	II	з	154-169	0.422	0.423
		R: CTCACCGACACCATACCCG						
AT37	KJ150256	F: ACTATCCGACCGAACGAACC [FAM]	(TAACC) ₁₂	Ш	7	287–338	0.489	0.558
		R: ACAGAACCGATTCTTGGCAT						
AT40	KJ150257	F: ATGGGTGTAGACCACGAACG [ATT0550]	(AT) ₁₃	II	7	113-115	0.273	0.258
		R: CGTGCTAGAATCATTGATCTTACCTT						
AT43	KJ150258	F: TTTCCGAATTTCAATCTGCTT [ATTO565]	$(AAAC)_5(AC)_{15}$	I	2	119–134	0.381	0.371
		R: CTGCCTTCTCCCTTAACGTG						
AT52	KJ150259	F: AACGAAGGTTTCAGAACGCC [YYE]	(AG) ₉	I	7	104-108	0.254	0.240
		R: CGCTGTTTCTCGGTTCTTGT						
Cross-amp	lifying loci							
AP1	Gouin et al. (2000)	F: TCTTGGGGATTGGCTAGTTG [YYE]		I	7	145-147	0.516	0.448
		R: CCTGAACTAAAAGGTGCTTTGG						
AP6	Gouin et al. (2000)	F: GCTGTGGGGATGGAGGT [FAM]		II	1	342	0.000	0.000
		R: CACTAGCGTATTCAAGCAACT						
Aitali3	Pedraza-Lara et al. (2011)	F: ATATCATGCGCGTCATCTCC [ATTO565]		II	9	249–308	0.579	0.582
		R: CACACGAGCACCAGACATTG						
Aitali4	Pedraza-Lara et al. (2011)	F: CGTTGATGGTAGGGGGGAAG [FAM]		I	1	196	0.000	0.000
		R: CGTCACCGACCATATAAAGTG						
N _A numbe	r of different alleles, Ho observed h	eterozygosity, $H_{\rm E}$ expected heterozygosity						

630

Polymorphism was relatively low in these samples. Two cross-amplifying loci were not polymorphic, the mean allele number was 2.75 and the mean observed heterozygosity 0.321 (Table 1). No significant deviations from Hardy-Weinberg and linkage equilibria were detected. Genetic differentiation among stone crayfish populations from different drainages was strong and highly significant (F_{ST} values between 0.171 and 0.364). Even the populations from the three closely adjacent but artificially fragmented streams were significantly differentiated (pairwise $F_{\rm ST}$ between 0.054 and 0.101, all P < 0.005). These markers will thus be useful to reconstruct historical population structure as well as recent fragmentation effects, despite their limited variability in these remnant populations at the edge of the species' distribution. The microsatellites will also be used to inform management of this endangered crayfish and to identify suitable source populations for reintroduction attempts.

Acknowledgments We thank ecogenics GmbH for help with marker development and Michael Kugler from the Office for Nature, Hunt and Fisheries of Canton St. Gallen for catching permits and support.

References

- Abdelkrim J, Robertson BC, Stanton JAL, Gemmell NJ (2009) Fast, cost-effective development of species-specific microsatellite markers by genomic sequencing. Biotechniques 46(3):185–191
- Gouin N, Grandjean F, Souty-Grosset C (2000) Characterization of microsatellite loci in the endangered freshwater crayfish Austropotamobius pallipes (Astacidae) and their potential use in other decapods. Mol Ecol 9(5):636–637
- Meglécz E, Costedoat C, Dubut V, Gilles A, Malausa T, Pech N, Martin J-F (2010) QDD: a user-friendly program to select microsatellite markers and design primers from large sequencing projects. Bioinformatics 26(3):403–404. doi:10.1093/bioinfor matics/btp670
- Pedraza-Lara C, Gonzalez EG, Bloor P, Doadrio I (2011) Isolation and characterization of novel polymorphic microsatellite loci for the white-clawed crayfish, *Austropotamobius italicus* (Faxon 1914). Mol Ecol Res 11(3):586–589. doi:10.1111/j.1755-0998. 2011.03004.x
- Sunnucks P, Hales DF (1996) Numerous transposed sequences of mitochondrial cytochrome oxidase I-II in aphids of the genus *Sitobion* (Hemiptera: Aphididae). Mol Biol Evol 13(3):510–524